The Activity of Certain Enzymes in Cadmium-Poisoned Chicks

By J. E. KENCH and P. J. D. GUBB. [*Medical Research Council (South Africa) Protein Research Unit, Department of Chemical Pathology, University of Cape Town, Cape Town, Republic of South Africa*]

Cadmium poisoning of men and animals gives rise to a serum albumin of low molecular weight (minialbumin), immunologically indistinguishable from normal albumin but lacking in tryptophan and with a deficit of lysine (Kench, Gain & Sutherland, 1965; Kench & Sutherland, 1966, 1967).

Activities of a number of enzymes in the liver and other tissues of newly hatched cadmium-poisoned chicks have now been compared with those of normal controls before and after incubation with Cd^{2+} at a concentration similar to that present *in vivo*. The purified enzyme, when available, was incubated with a range of concentrations of Cd^{2+} . For the experiment *in vivo* the usual treatment was a single injection of 12–15 μg of Cd^{2+} (as $\text{CdCl}_2 \cdot 2\frac{1}{2}\text{H}_2\text{O}$, approx. 0.11–0.13 μmol) into a superficial blood vessel of the chick embryo 7 days before hatching. The liver of newly hatched chicks was rapidly homogenized and fractionated by differential centrifugation in mannitol–sucrose–EDTA medium (Johnson & Lardy, 1965). Concentrations of Cd^{2+} in the various cellular fractions were determined, after wet oxidation, by atomic absorption spectrophotometry (P. J. D. Gubb & J. E. Kench, unpublished work).

On the basis of their reactivity with Cd^{2+} the ten enzymes examined could be grouped as follows: (1) inhibited both *in vitro* and *in vivo*, lipamide dehydrogenase, δ -aminolaevulate synthetase and xanthine dehydrogenase; (2) inhibited *in vitro* but not *in vivo*, cytochrome oxidase and succinate dehydrogenase; (3) resistant *in vitro* but biosynthesis impaired *in vivo*, catalase; (4) stimulated *in vitro* and *in vivo*, tryptophan oxygenase in liver cytosol of chicks, which was stimulated *in vitro* ($<0.2\text{ mM}$ - Cd^{2+}) and *in vivo* (70 μM - Cd^{2+}), though inhibited by Cd^{2+} in concentrations exceeding 0.5 mM; rat liver tryptophan oxygenase behaves in a similar manner (Sutherland & Kench, 1969); δ -aminolaevulate dehydratase, erythrocytic adenosine triphosphatase and the malate dehydrogenase isoenzyme in liver cytosol were other enzymes whose activity was enhanced by Cd^{2+} at certain concentrations; (5) was relatively insensitive to Cd^{2+} both *in vitro* and *in vivo*, mitochondrial malate dehydrogenase; inhibition by Cd^{2+} of the purified enzyme from pig heart was additive to inactivation by iodine, the two inhibitors not competing for sites on the enzyme.

Our findings are in accord with selective reactivity of thiol groups in different proteins (Wallenfels & Eisele, 1970; Varrone, Consiglio & Covelli, 1970). Interaction of Cd^{2+} with enzymes may provide information on the localization of enzymes within mitochondria and other cellular structures.

Johnson, D. & Lardy, H. (1965). In *Methods in Enzymology*, vol. 10, p. 94. Ed. by Estabrook, R. W. & Pullman, M. E. New York: Academic Press Inc.

Kench, J. E., Gain, A. C. & Sutherland, E. M. (1965). *S. Afr. med. J.* **39**, 1191.

Kench, J. E. & Sutherland, E. M. (1966). *S. Afr. med. J.* **40**, 1109.

Kench, J. E. & Sutherland, E. M. (1967). *Br. J. ind. Med.* **24**, 326.

Sutherland, E. M. & Kench, J. E. (1969). *Proc. S. Afr. Soc. Pathls*, July issue.

Varrone, S., Consiglio, E. & Covelli, I. (1970). *Eur. J. Biochem.* **13**, 305.

Wallenfels, K. & Eisele, B. (1970). *Biochem. J.* **117**, 48P.

Anticonvulsant Drugs and Xylose Uptake by Cerebral-Cortex Slices

By P. GRAY and J. C. GILBERT. (*Department of Biochemistry, University of St Andrews, St Andrews, Fife, U.K.*)

Xylose and glucose may be transported into brain by a common mechanism (Gilbert, 1965; Joanny, Corriol & Hillman, 1969), and for permeability studies xylose has the advantage of not being metabolized by brain.

Studies of the effects of anticonvulsants on sugar uptake may throw light on the mechanism of action of these drugs, because glucose is the principal oxidizable substrate utilized by brain, and it may also influence the activity of cerebral enzymes and the stability of cerebral membranes.

In the present studies the uptakes of xylose by guinea-pig cerebral-cortex slices incubated at 37°C in oxygenated bicarbonate media, containing pyruvate as nutrient and raffinose as extracellular space marker, were measured under a variety of conditions. The slices were always preincubated for 30 min at 37°C in the absence of xylose before transfer to a similar medium containing xylose. Any drugs and inhibitors tested were dissolved in both preincubation and incubation media.

The uptake of xylose (50 mM) by the non-raffinose compartment of the slices over a 9 min period was significantly stimulated by acetazolamide (20 mM) and by ethosuximide (500 μM), whereas, as reported by Gilbert, Ortiz & Millichap (1966), diphenylhydantoin was without significant effect. Initial-velocity studies indicated that the apparent K_m and V_{max} for xylose were both increased by

acetazolamide and were both decreased by ethosuximide (an effect previously observed with phenobarbitone). The thiol-group inhibitors iodoacetamide (100 μ M) and 5,5'-dithiobis-(2-nitrobenzoic acid) (100 μ M) did not alter the rate of xylose uptake over a 9 min incubation period. However, both inhibitors abolished the stimulatory effects of acetazolamide and ethosuximide.

The significance of these results will be discussed.

Gilbert, J. C. (1965). *Nature, Lond.*, **205**, 87.

Gilbert, J. C., Ortiz, W. R. & Millichap, J. G. (1966). *J. Neurochem.* **13**, 247.

Joanny, P., Corriol, J. & Hillman, H. (1969). *Biochem. J.* **112**, 367.

Microbial Metabolism of Alkylbenzenesulphonates

By A. J. WILLETTTS and R. B. CAIN. (*Microbiology Group, Department of Botany, University of Newcastle upon Tyne, Newcastle upon Tyne NE2 7RU, U.K.*)

Alkylbenzenesulphonates are a group of detergent compounds in common domestic and industrial usage. Several reports (Nelson, McKinney, McAteer & Konecky, 1961; Huddleston & Allred, 1963; Swisher, 1963) have concluded that this group of substituted aromatic compounds is biodegraded, but previous research has almost exclusively been limited to whole-cell studies with a mixed microflora degrading, in many instances, an isomeric mixture of alkylbenzenesulphonates. This communication reports the enzymology of (1-phenylundecyl)benzene-*p*-sulphonate degradation by a pure culture of a *Bacillus* sp. elected for growth on alkylbenzenesulphonates as the major source of carbon and sulphur.

Continuous sampling experiments, after inoculation of the pure strain of *Bacillus* species into alkylbenzenesulphonate medium, indicated that the sequence of biochemical events was: (1) removal of the sulphonate grouping, probably at the cell surface, leaving free sulphite and a *p*-hydroxyphenylalkanoic acid in the medium; (2) assimilation of sulphite and its subsequent intracellular oxidation to sulphate plus reduction to sulphide; (3) assimilation of the *p*-hydroxyphenylalkanoic acid, which was subsequently degraded intracellularly by classical β -oxidation of the alkyl side chain followed by 'ortho' cleavage of the aromatic nucleus (Ornston & Stanier, 1964).

The specific activity of the desulphonating enzyme, which was assayed by following sulphite production (Robinson, 1965), was increased 100-fold by growth on (1-phenylundecyl)benzene-*p*-sulphonate over that observed in succinate-grown cells. The enzyme was repressed but not inhibited

by sulphite in the medium. Substrate-specificity studies implicated isosteric activation of the enzyme. Sulphite-cytochrome *c* oxidoreductase activities were also elevated sixfold after growth on alkylbenzenesulphonates, but adenosine phosphosulphate-cytochrome *c* reductase was constitutive in this *Bacillus*.

Enzymes associated with β -oxidation were recorded in alkylbenzenesulphonate-grown cells. The acyl-CoA acyltransferase present was active with a range of detergent isomers and β -hydroxyacyl-CoA dehydrogenase was also detected. The activity of isocitrate lyase, an enzyme characteristically induced in metabolic sequences producing acetyl-CoA, was 90-fold higher in alkylbenzenesulphonate-grown than in succinate-grown cells.

The aromatic nucleus produced by biodegradation of alkylbenzenesulphonate homologues containing an even number of carbon atoms in the side chain was identified by g.l.c. and t.l.c. as *p*-hydroxyphenylacetate. The equivalent aromatic nucleus from homologues with an odd number of carbon atoms in the side chain was identified as *p*-hydroxybenzoate. All the enzymes to degrade the appropriate aromatic by a classical 'ortho' pathway to succinate and acetate were induced to high activities in *Bacillus* after growth on the appropriate alkylbenzenesulphonate isomer.

This work was supported by the Science Research Council and The Royal Society.

Huddleston, R. L. & Allred, R. C. (1963). *Devs ind. Microbiol.* **4**, 24.

Nelson, J. F., McKinney, R. E., McAteer, J. H. & Konecky, M. S. (1961). *Devs ind. Microbiol.* **2**, 93.

Ornston, L. N. & Stanier, R. Y. (1964). *Nature, Lond.*, **204**, 1279.

Robinson, H. C. (1965). *Biochem. J.* **94**, 687.

Swisher, R. D. (1963). *J. Am. Oil Chem. Soc.* **40**, 648.

Metabolism of Aromatic Compounds by Fungi: Conversion of β -Carboxymuconolactone into 3-Oxoadipate in *Aspergillus niger*

By D. R. THATCHER and R. B. CAIN. (*Microbiology Group, Department of Botany, University of Newcastle upon Tyne, Newcastle upon Tyne NE1 7RU, U.K.*)

The conversion of protocatechuate into 3-oxoadipate was first shown to proceed via β -carboxymuconolactone [3-carboxy-4-hydroxyhex-2-enedioic acid (1 $\rightarrow$ 4)-lactone] in *Neurospora crassa* (Gross, Gafford & Tatum, 1956) and later incriminated as the lactone intermediate characteristic of fungal aromatic metabolism by Cain, Bilton & Darrah (1968). Prokaryotic organisms neither form β -carboxymuconolactone nor do they