- Lenton, J. R., Bowen, M. R. & Saunders, P. F. (1968). *Nature, Lond.*, **220**, 86.
- Milborrow, B. V. (1967a). *Planta*, **76**, 93.
- Milborrow, B. V. (1967b). *Proc. 6th int. Conf. Plant Growth Substances, Ottawa*. (In the Press.) Ed. by Wightman, F. Noddle, R. C. & Robinson, D. R. (1969). *Biochem. J.* **112**, 547.
- Ohkuma, K., Addicott, F. T., Smith, O. E. & Thiessen, W. E. (1965). *Tetrahedron Lett.* p. 2529.
- Ohkuma, K., Lyon, J. L., Addicott, F. T. & Smith, O. E. (1963). *Science*, **142**, 1592.
- Robinson, D. R. (in Preparation)
- Rothwell, K. & Wain, R. L. (1964). *Proc. 5th int. Conf. Natural Plant Growth Regulators*, p. 363. Ed. by Nitsch, J. P.
- Taylor, H. F. & Smith, T. A. (1967). *Nature, Lond.*, **215**, 1514.
- Varner, J. E. & Johri, M. M. (1967). *Proc. 6th int. Conf. Plant Growth Substances, Ottawa*: (in the Press). Ed. by Wightman, F.
- Villiers, T. A. (1968). *Planta*, **82**, 342.

Gibberellins and their Biosynthesis

By CHARLES A. WEST. (*Division of Biochemistry, Department of Chemistry, University of California, Los Angeles, Calif., U.S.A.*)

The gibberellins were first discovered in culture filtrates of the phytopathological fungus *Gibberella fujikuroi* as the products that stimulated internode elongation in rice seedlings. Subsequent investigations of the physiological activities of the gibberellins have revealed a wide range of developmental processes in flowering plants, including germination, dormancy, mobilization of food reserves in the endosperm, stem elongation, leaf growth and morphology, floral induction, parthenocarp and fruit growth, that can be influenced under appropriate circumstances by exogenous gibberellins. A number of field applications of the gibberellins in controlling the growth of economic plants are in use.

The gibberellins have been shown to occur naturally in a wide variety of angiosperms as well as in the fungus. These findings have led to the generally held view that the gibberellins are natural growth regulators or hormones in flowering plants. To date, 24 different gibberellins of defined structure have been isolated from *G. fujikuroi* culture filtrates and angiosperm sources, and it seems certain from both published and unpublished reports that additional members of this family will be identified. All of these gibberellins are diterpenoids based on the *enantio*-gibberellane carbon skeleton (gibbane nucleus).

The production of gibberellins in plants, and hence the course of plant development, appears to be under the control of a number of environmental as well as genetic factors. A detailed understanding

of the biosynthetic pathway and the characteristics of the enzymes involved in gibberellin production should thus provide a basis for investigations of natural regulation of some aspects of plant development. It may also provide a more rational basis for the development of methods of artificial control of these processes.

The results of radioactive-tracer studies with [$1\text{-}^{14}\text{C}$]acetate and [$2\text{-}^{14}\text{C}$]mevalonate (Birch, Richards & Smith, 1958) confirmed the suggested isoprenoid origin of the gibberellins. The co-occurrence of (–)-kaurene and gibberellins in culture filtrates and the conversion of (–)-[$17\text{-}^{14}\text{C}$]-kaurene into [$17\text{-}^{14}\text{C}$]gibberellic acid in *G. fujikuroi* cultures (Cross, Galt & Hanson, 1964) established the intermediary role of this diterpene hydrocarbon in the biosynthetic pathway. Information about intermediates and the enzymes participating in the biosynthesis and further metabolism of (–)-kaurene has come from studies of higher-plant cell-free systems from wild cucumber (*Echinocystis macrocarpa*) endosperm and castor bean (*Ricinus communis*) seedlings. Soluble enzymes catalyse the synthesis of the acyclic diterpene precursor *trans*-geranylgeranyl pyrophosphate from mevalonate (Oster & West, 1968). Another soluble enzyme system mediates the cyclization of *trans*-geranylgeranyl pyrophosphate to (–)-kaurene (Upper & West, 1967). A bicyclic intermediate, *trans*-copalyl pyrophosphate, has recently been shown to be an intermediate in this conversion. (–)-Kaurene is then converted successively into a series of oxidation products (–)-kauren-19-ol, (–)-kauren-19-al, (–)-kauren-19-oic acid and (–)-7 β -hydroxy-kauren-19-oic acid (Dennis & West, 1967). These reactions are catalysed by a particulate cell fraction requiring NADPH and oxygen, and have the general characteristics of mixed-function oxidases.

Some suggestions about intermediates and the sequence of reactions in later stages of gibberellin biosynthesis have come from tracer studies in *G. fujikuroi* cultures (see, e.g., Cross, Norton & Stewart, 1968; Verbiscar, Cragg, Geissman & Phinney, 1967).

The enzymic site of action of several plant-growth retardants that have been shown to interfere with the overall synthesis of gibberellins (Kende, Ninnemann & Lang, 1963; Ninnemann, Zeevaert, Kende & Lang, 1964; Dennis, Upper & West, 1965) have been investigated in cell-free systems. In extracts of *E. macrocarpa* endosperm, *R. communis* seedlings and *G. fujikuroi* mycelia 2'-isopropyl-4'-(trimethylammonium chloride)-5'-methylphenyl piperidine-1-carboxylate (Amo 1618) selectively inhibits, among the reactions between mevalonate and (–)-kaurene, the conversion of *trans*-geranylgeranyl pyrophosphate into copalyl pyrophosphate. β -Chloroethyltrimethylammonium chloride (CCC)

inhibits the same step in the last-named two extracts, but is ineffective in *E. macrocarpa* endosperm. Tributyl-2,4-dichlorobenzylphosphonium chloride (Phosfon), on the other hand, was less selective in that it inhibited both steps of the cyclization of geranylgeranyl pyrophosphate to (-)-kaurene in the *G. fujikuroi* extracts. Phosfon was also shown to block the formation of other cyclic diterpene hydrocarbons from geranylgeranyl pyrophosphate in *R. communis* seedling extracts.

- Birch, A. J., Richards, R. W. & Smith, H. (1958). *Proc. chem. Soc.* p. 192.
 Cross, B. E., Galt, R. H. B. & Hanson, J. R. (1964). *J. chem. Soc.* p. 295.
 Cross, B. E., Norton, K. & Stewart, J. C. (1968). *J. chem. Soc. C*, p. 1054.
 Dennis, D. T., Upper, C. D. & West, C. A. (1965). *Plant Physiol.* **40**, 948.
 Dennis, D. T. & West, C. A. (1967). *J. biol. Chem.* **242**, 3293.
 Kende, H., Ninnemann, H. & Lang, A. (1963). *Naturwissenschaften*, **50**, 599.
 Ninnemann, H., Zeveaart, J. A. D., Kende, H. & Lang, A. (1964). *Planta*, **61**, 229.
 Oster, M. O. & West, C. A. (1968). *Arch. Biochem. Biophys.* **127**, 112.
 Upper, C. D. & West, C. A. (1967). *J. biol. Chem.* **242**, 3285.
 Verbisar, A. J., Cragg, G., Geissman, T. A. & Phinney B. O. (1967). *Phytochemistry*, **6**, 807.

The Wilting Toxins of *Fusicoccum amygdali* Del.

By K. D. BARROW and E. B. CHAIN. (*Department of Biochemistry, Imperial College of Science and Technology, London S.W.7*)

Cultures of *Fusicoccum amygdali* Del., a wilting pathogen of peach and almond in Southern Europe, have been shown to produce a number of phytotoxic metabolites termed the fusicoccins (Ballio *et al.* 1964; Barrow *et al.* 1968a). Strain selection and modifications of earlier media have raised the total fusicoccin yields to 250–350 µg./ml. in both shake flasks and stirred fermenters. The increased yield has led to a simplified isolation and separation procedure (G. T. Banks, E. B. Chain & R. Thomas, unpublished work).

Wilting is induced in plant tissues by the major metabolite fusicoccin A at concentrations as low as 0.1 µg./ml.

The structure of fusicoccin A is shown (I); it was determined by chemical degradation and extensive use of physical methods (Barrow *et al.* 1968a,b), and by an X-ray structure determination of a suitable heavy-atom derivative of the fusicoccin aglycone (Hough, Hursthouse, Neidle & Rogers, 1968).

(I) Me, methyl; Ac, acetyl

The empirical formula of fusicoccin A is $C_{36}H_{56}O_{12}$. It contains one methoxyl and two acetoxy groups, forms a dihydro derivative, $C_{36}H_{58}O_{12}$, on catalytic hydrogenation and a penta-acetate, $C_{42}H_{62}O_{15}$, on acetylation. Treatment of fusicoccin A with acid liberates (per mole) 1 mole of glucose and 1,1-dimethylallyl alcohol, which dehydrates to isoprene. At alkaline pH fusicoccin A is transformed into the metabolites fusicoccin B ($C_{36}H_{56}O_{12}$) and fusicoccin C ($C_{34}H_{54}O_{11}$) and finally into fusicoccin D (deacetyl fusicoccin, $C_{32}H_{52}O_{10}$). The distribution of acetoxy groups between the aglycone and sugar residues was readily deduced from mass-spectral measurements. Fusicoccin A and B have one acetate group on the sugar and one on the aglycone, fusicoccin C has one acetate group on the aglycone, and the penta-acetate has three acetate groups on the sugar and two on the aglycone. The glucose must therefore be disubstituted and the consumption of 2 moles of periodate by fusicoccin D establishes the 1,6-disubstitution. Two of the products from the reaction of fusicoccin A with acid, fusicoccin E ($C_{29}H_{46}O_{11}$) and fusicoccin F ($C_{27}H_{44}O_{10}$), are glucosides, and the group attached to position 6 of the glucose is therefore 1,1-dimethylallyl alcohol. The aglycone ($C_{21}H_{34}O_5$) can be prepared in low yield by acid hydrolysis, but more efficient methods are the Smith degradation (Abdel-Akher, Hamilton, Montgomery & Smith, 1952) (yield 65–75%) or reaction of the product from the periodate oxidation of fusicoccin D with strong base (Dugan & de Mayo, 1965) (yield 30–40%). Extensive n.m.r. and mass-spectral studies on the aglycone and its derivatives, including a number of selectively oxidized products, were used to elucidate the structure. The reaction of the aglycone with mercuric acetate followed by treatment with bromide ions gave a suitable derivative for the X-ray structure (Hough *et al.* 1968). A similar derivative was also prepared from