Hunter, unpublished work). The sensitivity of tissue homogenates to urea, phenol and some proteolytic action suggests involvement of protein, but the reverse is suggested by the stability to heat and formalin. Similarly the sensitivity of scrapie to periodate is suggestive of polysaccharide involvement, but unfortunately methods of isolation that utilize phenol or urea inactivate.

The association of scrapie activity with plasma membrane and endoplasmic-reticulum fragments has been discussed by Hunter (1969). This substantiates hypotheses proposed by Gibbons & Hunter (1967) and Adams & Field (1968) that both the disease process and replication of the agent are located in a steric rearrangement of the component layers in a cellular membrane. Broadly considered as a protein-phospholipid double-layered structure with attached oligosaccharide residues, this conception allows for the variable effects of chemical treatment. Variations in permeability in different cell populations can explain the pathological changes in brain tissue compared with other tissues and also the probable differences in scrapie replication rate in the various organs after scrapie inoculation. Formation of membrane in the presence of a foreign component might well require an induced enzyme system whose own synthesis would be under genetic control.

The mode of replication of a modified membrane structure is problematical, but the stage has often been reached when malfunction has been postulated in a cellular process whose normal function, structure and differentiated synthesis is, itself, as yet not fully understood.

- Adams, D. H. & Caspary, E. A. (1967). *Brit. med. J.* **3**, 173.
 Adams, D. H., Caspary, E. A. & Field, E. J. (1969). *J. gen. Virol.* **4**, 89.
 Adams, D. H. & Field, E. J. (1968). *Lancet*, ii, 714.
 Alper, T., Cramp, W. A., Haig, D. A. & Clarke, M. C. (1967). *Nature, Lond.*, **214**, 764.
 Brotherston, J. G., Renwick, C. C., Stamp, J. T., Zlotnik, I. & Pattison, I. H. (1968). *J. comp. Path.* **78**, 9.
 Caspary, E. A. & Sewell, F. M. (1968). *Experientia*, **24**, 793.
 Dickinson, A. G., Meikle, V. M. H. & Fraser, H. (1968). *J. comp. Path.* **78**, 293.
 Dickinson, A. G., Stamp, J. T., Renwick, C. C. & Rennie, J. C. (1968). *J. comp. Path.* **78**, 313.
 Gajdusek, D. C. (1967). *New Engl. J. Med.* **276**, 392.
 Gibbons, R. A. & Hunter, G. D. (1967). *Nature, Lond.*, **215**, 1041.
 Hunter, G. D. (1969). *Biochem. J.* (in the Press).
 Hunter, G. D., Gibbons, R. A., Kimberlin, R. H. & Millson, G. C. (1969). *J. comp. Path.* **79**, 101.
 Kimberlin, R. H. (1969). *Biochem. J.* (in the Press).
 Kimberlin, R. H. & Hunter, G. D. (1967). *J. gen. Virol.* **1**, 115.

COLLOQUIUM ON 'STRUCTURE, FUNCTION AND BIOSYNTHESIS OF SMALL RIBONUCLEIC ACID SPECIES'

Fractionation of Low-Molecular-Weight Ribonucleic Acid

By U. E. LOENING. (*Department of Botany, University of Edinburgh*)

The fractionation of RNA in the size range from t-RNA* to the smaller ribosomal RNA will be considered. In general, methods that fractionate according to the composition of the RNA have been used for t-RNA, and methods dependent on molecular weight have been used for the others. This is largely because of the present stage of knowledge of the heterogeneity of t-RNA and the need to prepare pure t-RNA fractions, and the state of ignorance of the significance or function of the other low-molecular-weight RNA components.

Examples will be given of the fractionation of t-RNA by countercurrent distribution, by methylated albumin-kieselguhr and reversed-phase chromatography, and by DEAE-cellulose, DEAE-Sephadex and related columns. These methods

have shown differences in the t-RNA distribution in different tissues and t-RNA unique to the mitochondria and chloroplasts. Rapidly labelled low-molecular-weight RNA also shows fractions obtained by methylated albumin-kieselguhr chromatography that are distinct from the t-RNA and 5s RNA.

Fractionation by molecular weight has shown the existence of several types of RNA distinct from t-RNA. Column chromatography by methylated albumin-kieselguhr or by gel filtration, sedimentation, and gel electrophoresis will be compared. The gel-filtration methods depend almost entirely on molecular-weight differences and are not much influenced by composition or conformation. Gel electrophoresis gives the greatest resolution, but the column methods have the advantage that large amounts of RNA can be fractionated.

Examples of the gel-electrophoretic analysis of RNA will be shown to illustrate the following findings.

(a) Several homogeneous types of RNA other than t-RNA, 5s RNA and 7s RNA can be found.

* Abbreviation: t-RNA, transfer RNA.

Some of these represent a class of stable low-molecular-weight RNA in the nuclei of animal cells. There are differences in the weights of these in different species of mammal. The chloroplasts of higher plants also contain a range of low-molecular-weight RNA types, and a '5s' RNA that is distinct from the '5s' RNA of the cytoplasm.

(b) Ribosomal RNA tends to be degraded at specific points in the chain, leading to the formation of homogeneous low-molecular-weight fractions. These are not easy to distinguish from the RNA described above, and it is therefore important to establish that a low-molecular-weight component is not ribosomal. In general, the high resolution of gel electrophoresis, and the possibility of detecting trace amounts of RNA, has shown that it is almost impossible to prepare cell fractions in which degradation of RNA has been avoided. It is possible that part of the breakdown occurs *in vivo* in old ribosomes; a clear example of this in the case of chloroplast ribosomes will be shown.

(c) When the low-molecular-weight stable fractions and the ribosomal-RNA breakdown products can be identified or eliminated, it should be possible to detect messenger RNA. In the favourable cases in which a cell is synthesizing largely one or a few polypeptide chains, it should be possible to detect the messenger RNA by u.v. absorption, and its molecular weight determined by gel electrophoresis should correlate with the peptide chain length. The messenger RNA for haemoglobin from reticulocytes has been fairly well characterized. Attempts in this laboratory to isolate the messenger RNA for the light and heavy antibody chains will be described.

Primary Sequences and their Determination

By G. G. BROWNEE. (*Medical Research Council Laboratory of Molecular Biology, Cambridge*)

The methods that have recently been developed for determining the sequence of low-molecular-weight RNA will be reviewed. Emphasis will be placed on the relative simplicity of the radioactivity approach, which uses microgram quantities of nucleic acid, in contrast with methods relying on the detection of the ultraviolet absorption of the base. The fundamental approach in finding the sequence of a nucleic acid, which is the same whether the RNA is radioactive or not, will be illustrated by referring to the methods used to find the sequence of uniformly ^{32}P -labelled 5s RNA of *Escherichia coli*. The extent to which these radioactivity techniques are applicable to larger RNA molecules will depend, primarily, on the ease of the fractionation and on the yields that may be obtained of partial fragments, say 50–100 residues long, isolated from them. For many large RNA species it may be possible to isolate such

fragments from partial enzymic digests and by finding their sequence to reconstruct long stretches of the molecule.

The radioactivity approach is not limited to nucleic acid labelled *in vivo*, and attempts are now being made to find the sequence of fragments of non-radioactive RNA, which are labelled at their 5'-hydroxyl end with [^{32}P]phosphate *in vitro*. This may be achieved by using a specific virus-induced phosphokinase and [γ - ^{32}P]ATP. This may be the method of choice for RNA which cannot be conveniently labelled *in vivo*, such as mammalian RNA.

The sequence of 13 transfer RNA molecules is now known, about half determined by classical and half by radioactivity methods. In all cases an anticodon triplet has been located in a specific region of the molecule which is of opposite polarity and which forms base pairs with the predicted codon. A likely secondary-structure model can be predicted, both from the base sequence and from the fact that some regions of the molecule are particularly resistant to enzymic digestion. All transfer RNA species have some sequences in common, which could either represent common features necessary for the maintenance of the tertiary structure, or represent common functional regions. Attempts at correlating specific functions with specific regions of the molecule will be discussed by referring to results obtained with various transfer RNA molecules.

A Molecular Model for Transfer Ribonucleic Acid

By W. FULLER, S. ARNOTT and J. CREEK. [*Department of Biophysics and Medical Research Council Biophysics Research Unit, King's College (University of London), 26–29 Drury Lane, London W.C.2*]

Physical, chemical and biological studies on transfer RNA will be discussed in the context of a model that we have built for the three-dimensional structure of the molecule. With respect to its base-pairing this model is based on the clover-leaf structure. It has the T ψ C arm stacked between the amino acid and anticodon arms so that these three helical regions are coaxial. The T ψ C loop is assumed to have a conformation like that proposed by Fuller & Hodgson (1967) for the anticodon loop. The features of the structure (i.e. the dihydrouracil arm and the extra loop) that vary with the different transfer RNA species came close to each other in our model, forming a protuberance or knob on the helical stem formed by the amino acid, T ψ C and anticodon arms. The conformation of the stem is similar to that observed in crystalline fibres of extensive, regular, two-stranded RNA and is the