of the androgen metabolites ranged from 7 to 40 min. as monitored by a flame ionization detector.

In the detector hydrogen is burned in a stream of air, producing water. The metabolites emerging from the column pass into the flame and are oxidized during detection. If steroid oxidation is complete the only products are water and carbon dioxide. It is therefore possible by condensing the water vapour and absorbing the carbon dioxide emerging from the detector to monitor the ^3H or ^{14}C associated with each detected peak.

The 'dead space' of the detector was reduced to a minimum by a polytetrafluoroethylene insert. Effluent gases emerged via a narrow thin-walled polytetrafluoroethylene tube, which passed through an ice bath in order to condense the water vapour. The condensate was collected in scintillation-counting vials containing 7 ml. of scintillator (Bray, 1960). Radioactivity recoveries of $81 \pm 5\%$ have been obtained, indicating that oxidation is complete and that within this closed system trapping is efficient. During eight replicate experiments radioactivity determinations, of individual metabolites, were within $\pm 5\%$.

Belham, J. E., Neal, G. E. & Williams, D. C. (1968). *Biochem. J.* **109**, 33 p.

Bray, G. A. (1960). *Analyt. Biochem.* **1**, 278.

The Reception of Androgens in the Rat Ventral Prostate

By J. E. BELHAM, G. E. NEAL and D. C. WILLIAMS.
(*Marie Curie Memorial Foundation, Caterham, Surrey*)

Adult castrated rats were injected intraperitoneally with [^3H]testosterone *in vivo*. After 40 min. the animals were killed and prostatic nuclear and cytoplasmic fractions were prepared. Androgens were isolated by coupled t.l.c. and g.l.c. Labelling was predominantly associated with dihydrotestosterone. Anderson & Liao (1968) reported similar findings for prostatic nuclei, but in their experiments labelling of dihydrotestosterone in the cytoplasmic fraction was less predominant. The administration *in vivo* of cyproterone acetate, an anti-androgen whose action has been shown to be on the target organ (Neumann & Berswordt-Wallrabe, 1966), resulted in marked decrease in the amounts of labelled dihydrotestosterone recovered from both cytoplasmic and nuclear fractions after injection of [^3H]testosterone *in vivo*.

Studies *in vitro* with isolated prostatic nuclear and cytoplasmic fractions showed that [^3H]testosterone is converted into a range of metabolites, given in decreasing order of labelling: androstenediol, dihydrotestosterone, androstenedione, andro-

sterone and androstenedione. Addition of cyproterone acetate to the incubations indicated that the anti-androgen does not fundamentally alter the pattern of metabolism of testosterone.

These results could indicate that, of the metabolites formed *in vivo*, dihydrotestosterone is selectively retained within the prostate in both the nucleus and the cytoplasm.

Experiments *in vitro*, in which [^3H]testosterone and dihydrotestosterone were added to nuclear and high-speed-supernatant fractions obtained from castrated or cyproterone-treated castrated rats, indicated that cyproterone antagonizes the uptake of label *in vitro* by both these subcellular fractions. The high-speed-supernatant fractions obtained from prostates after injection of [^3H]testosterone *in vivo* were subjected to Sephadex G-100 column chromatography. High labelling in a substance tentatively identified as dihydrotestosterone was observed in the column-excluded fraction. Treatment of the rat with cyproterone acetate *in vivo* was found to decrease labelling in this high-molecular-weight fraction.

These results suggest that cytoplasmic receptors for androgens are present in the rat ventral prostate, and they are similar to those observed by Toft & Gorski (1966) for oestrogens in the rat uterus. In contrast with the uterine oestrogen receptors, however, it appears that in the rat ventral prostate metabolism of testosterone to dihydrotestosterone is a necessary prerequisite for attachment to the receptors.

It is a pleasure to thank Miss G. M. Baker for performing the animal experiments and Miss M. Reid for her technical assistance. We also acknowledge the generous gift of cyproterone acetate by Schering Chemicals Ltd.

Anderson, K. M. & Liao, S. (1968). *Nature, Lond.*, **219**, 277.
Neumann, F. & Berswordt-Wallrabe, R. von (1966). *J. Endocrin.* **35**, 363.

Toft, D. & Gorski, J. (1966). *Proc. nat. Acad. Sci., Wash.*, **55**, 1574.

The Properties of Fractions of Streptolysin S

By A. G. TAYLOR. [*Department of Pathology, Institute of Orthopaedics (University of London), Stanmore, Middlesex*]

Three biological activities have been attributed to streptolysin S. Originally described as a haemolytic toxin (Todd, 1938), it has since been shown to release enzymes from lysosomes (Weissmann, Keiser & Bernheimer, 1963) and also to induce blast transformation in a high percentage of cells when added to lymphocyte cultures (Hirschhorn, Schreiber, Verbo & Gruskin, 1964). It has been suggested by Taranta, Cuppari & Quagliata (1968),