glycolysis. In this investigation it has been assumed that the maximum activities of the regulatory enzymes of both glycolysis and the cycle will reflect the maximum rates of these two processes; and therefore maximum activities of α -glycerophosphate oxidase (EC 1.1.2.1) and phosphofructokinase (or hexokinase plus phosphorylase) have been measured in crude muscle extracts.

For insect flight muscles the maximum activities of these two enzymes were almost identical (expressed as μ moles of triose phosphate produced or oxidized/min./g. wet wt. of muscle), whereas in vertebrate muscle phosphofructokinase activity was about 100-fold higher than the activity of the oxidase: the activities were in blowfly (*Calliphora erythrocephala*) 90 and 110, in bumble bee (*Bombus hortorum*) 50 and 54, in cockroach (*Periplaneta americana*) 48 and 96, in locust (*Locusta migratoria*) 33 and 32, in butterfly (*Vanessa urticae*) 24 and 24, in the water bug (*Lethocerus cordofanus*) 8 and 8, in pheasant pectoral muscle 2.8 and 280, in pigeon pectoral muscle 1.2 and 36, in rabbit semitendinosus 0.2 and 16, in rabbit adductor magnus 0.8 and 60, in dogfish red muscle 0.1 and 24, and in dogfish white muscle 0.6 and 62 for the oxidase and phosphofructokinase respectively. It is concluded that for insect flight muscle the α -glycerophosphate cycle is of sufficient activity to account for reoxidation of all the glycolytic NADH, whereas in vertebrate muscle this cycle cannot account for more than about 2% of the required rate of NADH oxidation. In vertebrate white (anaerobic) muscle the conversion of pyruvate into lactate by lactate dehydrogenase is probably the most important mechanism for reoxidation of NADH. However, in red (aerobic) muscle, in which pyruvate is oxidized by the mitochondria, mechanisms other than the conversion of pyruvate into lactate or the α -glycerophosphate cycle must exist for reoxidation of glycolytic NADH.

Klingenberg, M. & Bücher, Th. (1960). *Annu. Rev. Biochem.* **29**, 669.

Pette, D. (1966). In *Regulation of Metabolic Processes in Mitochondria*, p. 28. Ed. by Tager, J. M., Papa, S., Quagliariello, E. & Slater, E. C. Amsterdam: Elsevier Publishing Co.

Sacktor, B. (1961). *Annu. Rev. Ent.* **6**, 103.

Sacktor, B. (1965). In *Physiology of Insecta*, vol. 2, p. 483. Ed. by Rockstein, M. New York and London: Academic Press.

The Effects of Calcium Ions on the Activities of Hexokinase, Phosphofructokinase and Fructose 1,6-Diphosphatase from Vertebrate and Insect Muscles

By H. VAUGHAN and E. A. NEWSHOLME. (*Agricultural Research Council Unit of Insect Physiology, Department of Zoology, University of Oxford*)

The coupling between nervous stimulation and mechanical activity in muscle is considered to be mediated by release of Ca^{2+} from the sarcoplasmic reticulum (for review see Weber, 1966): the increased concentration of Ca^{2+} in the sarcoplasm stimulates myofibrillar adenosine triphosphatase activity, which causes contraction; relaxation is brought about by reabsorption of the Ca^{2+} by the sarcoplasmic reticulum. The dependence of muscular contraction on a supply of ATP from catabolic processes (particularly glycolysis) suggested the possibility that the increased sarcoplasmic concentration of Ca^{2+} might stimulate the activities of the regulatory enzymes of glycolysis, or modify the basic control mechanism of glycolysis. A study was made of the effects of Ca^{2+} on the activities of hexokinase, phosphofructokinase and fructose 1,6-diphosphatase, and their inhibitions by glucose 6-phosphate, ATP and AMP respectively. As regulation of intracellular Ca^{2+} concentration may differ between red muscle and white muscle and between insect fibrillar muscle and non-fibrillar muscle (see Smith, 1966), enzymes from these four types of muscle have been used in this investigation.

The results show that increasing the Ca^{2+} concentration from 0.001 to $10 \mu\text{M}$ (the range that activates myofibrillar adenosine triphosphatase) has no effect on the activities of these enzymes from all types of muscle; nor does it change the inhibitions of hexokinase by glucose 6-phosphate, phosphofructokinase by ATP and fructose 1,6-diphosphatase by AMP. However, high concentrations of Ca^{2+} ($1000 \mu\text{M}$) inhibited these enzymes, although the inhibitions by glucose 6-phosphate, ATP and AMP were not changed.

Margreth, Cantani & Schiaffino (1967) have suggested that inhibition of frog leg muscle phosphofructokinase by Ca^{2+} might be an important mechanism of glycolytic control, as this enzyme might be located within the sarcoplasmic reticulum. An investigation of the possible location of hexokinase and phosphofructokinase by differential centrifugation showed that in vertebrate muscle a proportion of these enzymes sedimented with the reticulum fraction. However, with insect muscles the activities of these enzymes were found exclusively in the supernatant fraction. Thus in various muscles there is no correlation between the intracellular distribution of these two enzymes and their

sensitivities to Ca^{2+} inhibition; the physiological importance of the latter is therefore questionable.

Margreth, A., Cantani, C. & Schiaffino, S. (1967). *Biochem. J.* **102**, 35c.

Smith, D. S. (1966). *Progr. Biophys. molec. Biol.* **16**, 107.

Weber, A. (1966). In *Current Topics in Bioenergetics*, vol. 1, p. 203. Ed. by Sanadi, D. R. New York and London: Academic Press.

Studies on Equine Trypsinogen and Trypsin

By C. I. HARRIS* and T. HOFMANN.† (*Department of Biochemistry, University of Sheffield*)

Equine trypsinogen was prepared by extraction of minced pancreas with 0.25 M- H_2SO_4 , fractionation with $(\text{NH}_4)_2\text{SO}_4$ and chromatography on cellulose citrate (Whatman CT-60) at pH 6.0. Inactive components were also removed from commercial bovine trypsin by this chromatographic system. Equine trypsin was prepared from trypsinogen by autocatalytic activation in the presence of 0.05 M- CaCl_2 .

Purified trypsinogen and trypsin appeared homogeneous on ultracentrifugation and by disc electrophoresis (Reisfeld, Lewis & Williams, 1962). A molecular weight of 25600 ± 1000 for equine trypsin was determined by the high-speed equilibrium method of Yphantis (1964).

The *N*-terminal residues of trypsinogen and trypsin were found to be serine and isoleucine respectively after carbamoylation (Stark & Smyth, 1963; Gertler & Hofmann, 1967) and these were confirmed by dansylation (Gray, 1967). Identical *C*-terminal sequences (Gln, Thr, Ile, Ala)-Ala-Asn were found in carboxypeptidase A digests of *S*-sulpho-trypsinogen and *S*-sulpho-trypsin.

The structure of the activation peptide was found to be Ser-Ser-Thr-Asp₄-Lys. This is similar to the activation octapeptides from pig and sheep trypsinogens but differs markedly in the *N*-terminal two residues of the sequence. The amino acid composition of equine trypsin showed general similarities to the trypsins from cow, pig and sheep (see Schyns, Bricteux-Gregoire & Florkin, 1969) but differed in detail in the content of methionine (1 residue/mol.) and lysine and arginine (both 7 residues/mol.).

The common features of *N*-terminal isoleucine and *C*-terminal asparagine in the trypsins from cow and pig have thus been extended to equine trypsin. Similarly, the common sequence of Asp₄-Lys reported to be in the activation peptides from cow,

pig and sheep trypsinogens has also been found in this study. The functional significance of this sequence has been reported by Abita, Delaage, Lazdunski & Savrda (1969).

Abita, J. P., Delaage, M., Lazdunski, M. & Savrda, J. (1969). *Europ. J. Biochem.* **8**, 314.

Gertler, A. & Hofmann, T. (1967). *J. biol. Chem.* **242**, 2522.

Gray, W. (1967). In *Methods in Enzymology*, vol. 11, p. 149. Ed. by Hirs, C. H. W. New York: Academic Press Inc.

Reisfeld, R. A., Lewis, U. J. & Williams, D. E. (1962). *Nature, Lond.*, **195**, 281.

Schyns, R., Bricteux-Gregoire, S. & Florkin, M. (1969). *Biochim. biophys. Acta*, **175**, 97.

Stark, G. R. & Smyth, D. G. (1963). *J. biol. Chem.* **238**, 214.

Yphantis, D. A. (1964). *Biochemistry*, **3**, 297.

Zinc-Dependent Conformational Stability in the Alkaline Phosphatase of *Escherichia coli*

By C. N. A. TROTMAN and C. GREENWOOD. (*School of Biological Sciences, University of East Anglia, Norwich*)

Alkaline phosphatase of *Escherichia coli* has been found variously to contain 4, 3 or 2 g.atoms of Zn/mole (Simpson & Vallee, 1968; Reynolds & Schlesinger, 1967; Plocke, Levinthal & Vallee, 1962); at least part of the zinc content is essential to activity, and its absence through chelation or genetic alteration (Schlesinger, 1966) renders the enzyme inactive. Zinc has also been shown (Reynolds & Schlesinger, 1967) to be essential for subunit re-association but not necessary for refolding of the subunit from an extended-coil conformation.

In the present work ultraviolet circular dichroism (CD) has been employed to detect major conformational phenomena related to the presence of zinc, with wild-type enzyme from *E. coli* K 12 and a zinc deficient enzyme from *E. coli* C₁₀F₁₈.

The CD extrema near 221 and 207 nm. remained essentially unchanged when *E. coli* wild-type enzyme was suspended in high urea concentrations; the 192 nm. peak became inaccessible. *E. coli* wild-type enzyme depleted of zinc by dialysis against EDTA behaved quite differently; substantial deviation from natural conformation occurred at above 2 M-urea, characterized by loss of the 221 nm. and 207 nm. peaks and the appearance of a larger peak near 202 nm. Sedimentation-velocity analysis of EDTA-treated enzyme in 6 M-urea revealed that dissociation into subunits had occurred; the low sedimentation coefficient, $S_{20,w}$ 1.7s, was consistent with an extremely extended conformation.

E. coli C₁₀F₁₈ inactive enzyme behaved similarly to EDTA-treated *E. coli* wild-type enzyme when treated with urea solutions. *E. coli* C₁₀F₁₈ enzyme incubated with Zn^{2+} and phosphate with partial

* Present address: Department of Biochemistry, University of Birmingham.

† Present address: Department of Biochemistry, University of Toronto, Toronto, Ont., Canada.