is added), followed by a return to the slow steady rate of swelling.

Antimycin, added at the point of maximum swelling, prevented the reversal phase, and slow steady swelling resulted. Addition of oligomycin, 2,4-dinitrophenol or carbonyl cyanide *p*-trifluoromethoxyphenylhydrazone at the same point did not prevent reversal of swelling, which thus appears to be dependent on respiration but not high-energy intermediates.

Addition of ethylenedioxybis(ethyleneamine)-tetra-acetic acid before arsenate prevented significant swelling, and addition after arsenate apparently caused a return to the original volume. In the latter case this was accompanied by a decrease in the rate of oxygen uptake and restoration of some respiratory control.

These results are interpreted as follows. During the phase of accelerating swelling and respiration Ca^{2+} is being released and β -hydroxybutyrate equilibrates across the membrane. When respiration reaches a very high rate, product can leave faster than β -hydroxybutyrate can enter, possibly because the membrane is damaged, allowing outward leakage of K^+ . This results in reversal of swelling, superseded by a slow degradation of the membrane. Respiration can, however, still maintain some electrochemical gradient between the inside of the mitochondrion and the medium. When respiration ceases this gradient disappears and there is a small burst of swelling due to re-equilibration of ions across the membrane.

Hunter, F. E. & Ford, L. (1955). *J. biol. Chem.* **216**, 357.

Packer, L. (1961). *J. biol. Chem.* **236**, 214.

Ter Welle, H. F. & Slater, E. C. (1967). *Biochim. biophys. Acta*, **143**, 1.

Proton Transport in Mitochondrial Oxidative Phosphorylation

By M. J. SELWYN and A. P. DAWSON. (*School of Biological Sciences, University of East Anglia, Norwich*)

At physiological pH values phosphorylation of ADP produces approximately equimolecular amounts of OH^- ion. Since the inner mitochondrial membrane is not highly permeable to H^+ or OH^- ions, this internal production of OH^- ions must be compensated by some associated process. P_i is effectively exchanged for two OH^- ions (Chappell & Crofts, 1965), thus overcompensating for the internal production of OH^- ions. If the ATP-ADP exchange-diffusion carrier exchanged ATP^{3-} for ADP^{3-} the overall process would be balanced, but this mechanism would be thermodynamically un-

satisfactory since a pH differential that favoured P_i uptake would also favour retention of ATP.

If the ATP-ADP exchange-diffusion carrier exchanged ATP^{4-} for ADP^{3-} and overcompensation by the $\text{P}_i\text{-OH}^-$ ion exchange were compensated by a respiration-driven proton pump, then conditions of raised internal pH and negative internal electrical potential would favour uptake of P_i and exchange of internal ATP^{4-} for external ADP^{3-} .

This suggests a mechanism of oxidative phosphorylation in which production of non-phosphorylated high-energy intermediates and proton pumping proceed in parallel. Such a scheme has features that contrast with those of the chemical and chemiosmotic mechanisms proposed for oxidative phosphorylation.

Production of the chemical intermediate and the electrochemical potential difference at a particular phosphorylation site could result from the same electron-transfer step, but could also result from neighbouring electron transfers.

The magnitude of the electrochemical potential difference across the mitochondrial membrane would be small, except in the presence of substances, such as Ca^{2+} ions, that discharge the non-phosphorylated high-energy intermediate, when all the energy from electron transfer becomes available for generating electrochemical potential differences.

Uncouplers have to produce both proton permeability in the membrane and discharge of the non-phosphorylated high-energy intermediate, unless some other process can compensate for the proton pumping. 2,4-Dinitrophenol affects the activity of the mitochondrial adenosine triphosphatase coupling factor (Pullman, Penefsky, Datta & Racker, 1960; Selwyn, 1968) and also renders lipid membranes permeable to protons (Hopfer, Lehninger & Thompson, 1968). Ca^{2+} ions uncouple only in the presence of ions such as acetate, which can discharge the proton gradient (Lehninger, Carafoli & Rossi, 1967). Acetate ions alone do not uncouple and are not accumulated, since they do not discharge the chemical intermediate.

Chappell, J. B. & Crofts, A. R. (1965). *Biochem. J.* **95**, 393.

Hopfer, U., Lehninger, A. L. & Thompson, T. E. (1968).

Proc. nat. Acad. Sci., Wash., **59**, 484.

Lehninger, A. L., Carafoli, E. & Rossi, C. S. (1967). *Advanc. Enzymol.* **29**, 259.

Pullman, M. E., Penefsky, H. S., Datta, A. & Racker, E. (1960). *J. biol. Chem.* **235**, 3322.

Selwyn, M. J. (1968). *FEBS Lett.* **1**, 247.