The Electron-Transport Chain of *Tetrahymena pyriformis* Strain ST

By G. TURNER, D. LLOYD and B. CHANCE. (*Medical Research Council Group for Microbial Structure and Function, Department of Microbiology, University College of South Wales and Monmouthshire, Cardiff, and Johnson Research Foundation, Department of Biophysics, University of Pennsylvania, Philadelphia, Pa., U.S.A.*)

Previous studies with cell-free preparations and isolated mitochondria of *Tetrahymena pyriformis*, an obligate aerobic ciliate protozoon, have suggested that this organism has no cytochrome oxidase, as there is no absorption peak in difference spectra at 605nm. and reduced mammalian cytochrome *c* is not reoxidized by such preparations (Eichel, 1955; Kobayashi, 1965; Van de Vijver, 1966).

The present work demonstrates that the organism does possess a cytochrome component, which reacts with oxygen more rapidly than cytochromes *b* and *c*. Examination of difference spectra (oxidized—anaerobic) of isolated mitochondria revealed a peak of absorption at 620nm. but not at 605nm. (Turner & Lloyd, 1968). The velocities of reoxidation of cytochromes *b*, *c* and the 620nm.-absorbing component were measured by mixing an anaerobic suspension of mitochondria with oxygen in the regenerative-flow apparatus of Chance (1954). The values for the pseudo-first-order reaction-velocity constants and the corresponding half-times at 25°, with 16 μ M-oxygen and 5mM-succinate, were as follows: *b* (430–410nm.) $k_1 = 55 \text{ sec.}^{-1}$, $t_{1/2} = 13 \text{ msec.}$; *c* (553–540nm.) $k_1 = 120 \text{ sec.}^{-1}$, $t_{1/2} = 6 \text{ msec.}$; *a* (620–650nm.) $k_1 = 180 \text{ sec.}^{-1}$, $t_{1/2} = 4 \text{ msec.}$

Lowering of the oxygen concentration to 1 μ M did not appear to affect the reaction velocities of these components, suggesting high oxygen affinity of the oxidase.

It has previously been noted that there is a considerable variation in sensitivity to cyanide among different strains of *Tetrahymena* (Van de Vijver, 1966). Oxidation of α -oxoglutarate and of succinate by strain ST is 90–100% inhibited in the presence of 1mM-KCN.

Reoxidation of flavoprotein components of anaerobic mitochondria was studied in the regenerative-flow apparatus, with 16 μ M-oxygen, both spectrophotometrically (465–510nm., $t_{1/2} = 400 \text{ msec.}$) and fluorimetrically (435→520nm., $t_{1/2} < 100 \text{ msec.}$; 460→520nm., $t_{1/2} < 100 \text{ msec.}$; 475→520nm., $t_{1/2} = 1.0 \text{ sec.}$).

Although cytochrome *b* is completely reduced in state 5 by succinate, only 60% reduction occurs with α -oxoglutarate; cytochrome *c* and cytochrome *c* oxidase are completely reducible by each of these substrates. Antimycin A inhibits the reoxidation of cytochrome *b* by cytochrome *c*.

Although isolated mitochondria do not oxidize externally added NADH, a rotenone-sensitive NADH-cytochrome *c* oxidoreductase activity is present. Endogenous NAD⁺ is readily reducible on addition of α -oxoglutarate, but not of succinate.

G. T. holds a Medical Research Council Research Studentship and thanks the Medical Research Council for a travel grant.

Chance, B. (1954). *Disc. Faraday Soc.* **17**, 120.

Eichel, H. J. (1955). *Fed. Proc.* **14**, 43.

Kobayashi, S. (1965). *J. Biochem., Tokyo*, **58**, 444.

Turner, G. & Lloyd, D. (1968). *Abstr. 5th Meet. Fed. Europ. biochem. Soc., Prague*, A-822.

Van de Vijver, G. (1966). *Enzymologia*, **31**, 382.

DEMONSTRATION

The Dialysis of Small Volumes (5–50 μ l.) of Fluid

By N. G. HEATLEY. (*Sir William Dunn School of Pathology, University of Oxford*)

The fluid to be dialysed is placed on a disc (7.5mm. diam.) of cellophan or other semi-permeable membrane, which is held by slight suction to the upper end of a vertical glass tube containing water (or buffer etc.).

Two forms of the apparatus have been made:

(a) when the diffusate is needed, and (b) for removal of unwanted diffusible material; in this model the under side of the disc is irrigated continually with fresh dialysing fluid. (Heatley, 1969.)

In both models the disc and droplet may be covered with a dome containing desiccant. This reduces the volume of the drop and greatly accelerates dialysis; it also forces more of the total diffusible material into the diffusate.

Heatley, N. G. (1969). *Analyst*, (in the Press).