which is converted into Factor XIIa on exposure to a foreign surface, has been purified in the activated form by Shoenmakers, Matze, Haanen & Zilliken (1965), who have found that it has estereolytic activity that, like its coagulant activity, is inhibited by di-isopropyl phosphorofluoridate.

Thus at the present time it seems that there are two types of coagulation factors: those that are zymogens and become converted into enzymes, and those factors that potentiate the activity of the enzymic factors.

- Barton, P. G. (1967). *Nature, Lond.*, **215**, 1508.
 Barton, P. G., Jackson, C. M. & Hanahan, D. J. (1967). *Nature, Lond.*, **214**, 293.
 Davie, E. W. & Ratnoff, O. D. (1964). *Science*, **145**, 1310.
 Esnouf, M. P. & Williams, W. J. (1962). *Biochem. J.* **84**, 62.
 Gladner, J. A. & Laki, K. (1958). *J. Amer. chem. Soc.* **80**, 1263.
 Hanahan, D. J. & Papahadjopoulos, D. (1965). *Thromb. Diath. Haemorrh.* Suppl. no. 17, p. 71.
 Hemker, H. C., Esnouf, M. P., Hemker, P. W., Swart, A. C. W. & Macfarlane, R. G. (1967). *Nature, Lond.*, **215**, 248.
 Hemker, H. C. & Khan, J. C. (1967). *Nature, Lond.*, **215**, 1201.
 Jackson, C. M. & Hanahan, D. J. (1968). *Biochemistry*, **7**, 4506.
 Jobin, F. & Esnouf, M. P. (1967). *Biochem. J.* **102**, 666.
 Lechner, K. & Deutsch, E. (1965). *Thromb. Diath. Haemorrh.* **13**, 314.
 Leveson, J. E. & Esnouf, M. P. (1969). *Brit. J. Haemat.* **17**, 173.
 Macfarlane, R. G. (1964). *Nature, Lond.*, **202**, 498.
 Milstone, J. H. (1964). *Fed. Proc.* **23**, 742.
 Milstone, J. H., Oulianoff, N. & Milstone, V. K. (1963). *J. gen. Physiol.* **47**, 315.
 Morawitz, P. (1905). *Ergebn. Physiol.* **4**, 307.
 Papahadjopoulos, D. & Hanahan, D. J. (1964). *Biochim. biophys. Acta*, **90**, 436.
 Schoenmakers, J. G. G., Matze, R., Haanen, C. & Zilliken, F. (1965). *Biochim. biophys. Acta*, **101**, 166.
 Williams, W. J. & Esnouf, M. P. (1962). *Biochem. J.* **84**, 52.

The Chemistry of the Activation of Prothrombin

By STAFFAN MAGNUSSON. (*Medical Research Council Laboratory of Molecular Biology, Cambridge CB2 2QH*)

Bovine prothrombin prepared from oxalated plasma by precipitation of euglobulins at low ionic strength (Mellanby, 1930) followed by adsorption on magnesium hydroxide and elution with carbon dioxide (Fuchs, 1930), ammonium sulphate fractionation, isoelectric fractionation (Seegers, 1952) and equilibrium chromatography on cation-exchange resin (Amberlite IRC-50) (Miller, 1958) appears to be homogeneous during ultracentrifugation (Lamy & Waugh, 1958) and

has a molecular weight about 68 000. One mole of *N*-terminal alanine/mole (Magnusson, 1958; Miller, 1958) and about 11.5% carbohydrate, namely 4.3% sialic acid, 5.4% neutral sugar (galactose and mannose in a 1:1 ratio) and 1.8% glucosamine (Magnusson, 1965b), have been found. Neuraminidase degradation (Schwick & Schultze, 1959; Tishkoff, Pechet & Alexander, 1960) shows that sialic acid is terminal. Partial acid hydrolysis (Magnusson, 1965b) indicates that the oligosaccharide sequences are sialic acid-galactose-mannose-glucosamine-peptide. Autocatalytic activation in 25% sodium citrate (Seegers, 1949) and activation in the presence of thromboplastin, calcium chloride and Factor V have been studied by *N*-terminal amino acid analysis (Magnusson, 1958, 1965a) and both found to be proteolytic in nature. In both types of activation mixture 2 moles of *N*-terminal isoleucine/mole of *N*-terminal alanine were found at the peak of thrombin activity. Carboxypeptidase A and B degradation released only arginine and lysine from citrate-activation fragments, indicating that the activating enzyme has trypsin-like specificity (Magnusson & Steele, 1965). Miller & van Vunakis (1956) showed that di-isopropyl phosphorofluoridate inhibits thrombin. When they attempted citrate activation of prothrombin in the presence of di-isopropyl phosphorofluoridate no thrombin was formed and no prothrombin consumed. They also found that a mixture of thromboplastin, calcium chloride and dilute serum could be preincubated with di-isopropyl phosphorofluoridate without losing its prothrombin-activating capacity. When activation of prothrombin with thromboplastin, Factor V and calcium chloride was attempted in the presence of di-isopropyl phosphorofluoridate (Magnusson, 1965a) no thrombin was formed and no new *N*-terminal residues appeared. It is thus clear that both types of activation depend on a serine proteinase with trypsin-like specificity that is present in an inactive form in this type of prothrombin preparation. It seems unlikely at present that this enzyme is thrombin. A second enzyme activity is formed during citrate activation (Kowarzyk & Marciniak, 1961). It has esterase activity towards α -*N*-toluene-*p*-sulphonyl-L-arginine methyl ester but no thrombin activity and it can accelerate prothrombin activation. The stoichiometric relation of this activity to prothrombin is not clear. If, on the one hand, it turns out to represent only a very small amount of protein relative to the amount of thrombin it could either represent impurity in the prothrombin preparation, which seems unlikely, or it could be the product of a small proportion of prothrombin molecules, activated in an alternative way to that which produces thrombin. In the latter case one would expect it to contain the same active-

centre sequence as thrombin. If, on the other hand, it turns out to be present in stoichiometric proportion to thrombin it could be part of the covalent structure of prothrombin.

It is clear (Magnusson, 1965*d*) that thrombin formed in both citrate activation and thromboplastin activation contains the *N*-terminal sequence Ile-Val-Glu-Gly-. Recent evidence for the amino acid sequence (S. Magnusson & B. S. Hartley, unpublished work) of the B-chain of bovine thrombin isolated from thromboplastin activation mixtures (Magnusson, 1965*c*) shows sufficient sequence homology with the pancreatic serine proteinases for one to feel confident that the activation mechanism is very similar to that of chymotrypsinogen (Sigler, Blow, Matthews & Henderson, 1968), the proteolytic cleavage of the peptide bond to produce the *N*-terminal isoleucine being the critical step that produces active thrombin. Since the amino acid composition of prothrombin is very similar to that of thrombin and the molecular weight nearly twice that of thrombin, we have tried to obtain evidence that prothrombin of molecular weight 68000 is a dimer. Reduction and carboxymethylation, even if followed by maleylation, failed to produce material of smaller molecular weight as evidenced by gel filtration (S. Magnusson & B. S. Hartley, unpublished work). Unless one invokes some kind of covalent bond other than a disulphide bond to hold two polypeptide chains together, the best hypothesis at the moment seems to be that prothrombin consists of only one polypeptide chain with the B-chain of thrombin (*C*-terminal serine) constituting the *C*-terminal part and the A-chain constituting part of the polypeptide chain between the *N*-terminal alanyl sequence and the B-chain. This leaves about 250 amino acid residues and about 75% of the carbohydrate of prothrombin to be accounted for as 'non-thrombin' or 'pro' material. This, incidentally, is sufficient polypeptide material to include a second serine proteinase. The key questions at the moment are whether the second proteinase is in fact part of the covalent structure of prothrombin or, as has been suggested by others (Seegers, 1969), prothrombin is an aggregate of two different proteins held together by non-covalent forces. The second important question is whether the second proteinase can be activated by some other means than proteolysis, which could be the common denominator of all the different systems that are known to trigger off prothrombin activation, such as concentrated salt solutions, phospholipids and basic polyamino acids.

Bovine thrombin and prothrombin used in this study is prepared at the Chemistry Department II, Karolinska Institutet, Stockholm, Sweden. Support from the Swedish Medical Research Council and the U.S. National Institutes of Health is gratefully acknowledged.

- Fuchs, H. J. (1930). *Biochem. Z.* **222**, 470.
 Kowarzyk, H. & Marciniak, E. (1961). *Pol. Tyg. lek.* **16**, 1641.
 Lamy, F. & Waugh, D. F. (1958). *Thromb. Diath. Haemorrh.* **2**, 12.
 Magnusson, S. (1958). *Acta chem. scand.* **12**, 355.
 Magnusson, S. (1965*a*). *Ark. Kemi*, **23**, 271.
 Magnusson, S. (1965*b*). *Ark. Kemi*, **23**, 285.
 Magnusson, S. (1965*c*). *Ark. Kemi*, **24**, 349.
 Magnusson, S. (1965*d*). *Ark. Kemi*, **24**, 375.
 Magnusson, S. & Steele, B. (1965). *Ark. Kemi*, **24**, 359.
 Mellanby, J. (1930). *Proc. Roy. Soc. B*, **107**, 221.
 Miller, K. D. (1958). *J. biol. Chem.* **231**, 987.
 Miller, K. D. & van Vunakis, H. (1956). *J. biol. Chem.* **223**, 227.
 Schwick, G. & Schultze, H. E. (1959). *Clin. chim. Acta*, **4**, 26.
 Seegers, W. H. (1949). *Proc. Soc. exp. Biol., N.Y.*, **72**, 677.
 Seegers, W. H. (1952). *Rec. chem. Progr. Kresge-Hooker Sci. Lib.* **13**, 143.
 Seegers, W. H. (1969). *Annu. Rev. Physiol.* **31**, 269.
 Sigler, P. B., Blow, D. M., Matthews, B. W. & Henderson, R. (1968). *J. molec. Biol.* **35**, 143.
 Tishkoff, G. H., Pechet, L. & Alexander, B. (1960). *Blood*, **15**, 778.

Prochymosin and Chymosin (Prorennin and Rennin)

By B. FOLTMANN. (*The Protein Laboratory, University of Copenhagen, Denmark*)

The milk-clotting enzyme from the fourth stomach of the calf has for many years been called rennin in the English-speaking countries. Confusion very often occurs, however, between rennin (EC 3.4.4.3) from the calf stomach and renin (EC 3.4.4.15) from the kidneys. The milk-clotting enzyme was called chymosin 50 years before the term rennin was introduced (Deschamp, 1840; Lea & Dickinson, 1890), and to avoid further misunderstanding I return to the old nomenclature. The chemistry of the enzyme has been reviewed (Foltmann, 1966).

Like other acidic proteases from the gastric juice, chymosin is secreted as an inactive precursor, prochymosin. The precursor is stable at weakly alkaline pH whereas the active enzyme is rapidly denatured at pH values above 7. Below pH 5 prochymosin is converted into chymosin by a limited proteolysis during which a peptide segment is cleaved from the *N*-terminus.

If the increase of milk-clotting activity is plotted against time, the course of the activation at pH 4.5–5 appears as more or less sigmoid curves, indicating that the activation is at least partly autocatalytic. Experiments carried out at lower pH show a different course of reaction. The lower the pH the higher is the initial rate of reaction and the sigmoid shape of the curve disappears.