Species Body Size and Liver Glutamine Synthetase, Glutaminase and Glutamine Aminotransferase

By E. V. ROWSELL, P. J. BUTTERY* and J. A. CARNIE. (*Department of Biochemistry, University of Manchester Institute of Science and Technology, Manchester 1*)

Previous work on various mammalian species has shown that as adult body size diminishes certain liver enzymes concerned with amino acid metabolism increase greatly in activity: pyruvate aminotransferases (Lin, Rivlin & Knox, 1959; Rowsell & Carnie, 1964), L-alanine-2-oxoglutarate aminotransferase (Cornelius, 1963) and serine dehydratase (Rowsell, Carnie & Wahbi, 1965). We are surveying for other enzymes in liver homogenates with the prospect that more will be found having a similar dependence on body size; the significance of this phenomenon may then become clearer.

Glutamine synthetase was assayed as described by Elliott (1955), by calculating rates from Wu's (1963) extinction coefficient value for the Fe^{+++} complex of L-glutamyl- γ -hydroxamate; glutaminase in 0.05M-potassium phosphate buffer, pH 7.8, with 0.02M-L-glutamine, the glutamate liberated being measured in heat-inactivated incubation samples with ninhydrin after chromatographic separation; and glutamine amino-transferase as described by Buttery, Rowsell & Carnie (1969). In all cases initial rates were obtained from linear progress curves under conditions where rate was proportional to liver concentration.

Assays with fresh liver from the local abattoir were included and there were delays in transport (never more than 45 min.) before these could be started. Some attempt was made to gauge what losses in activity this might involve. When freshly obtained chopped rat liver was stored ice-cold in the appropriate buffer for homogenization (as in conveying abattoir samples), the activities measured after 2 hr. for glutamine synthetase, glutaminase and glutamine aminotransferase were respectively (compared with immediate assay) 87%, 77% and 98%.

Glutamine aminotransferase activities [μ moles of 2-oxoglutarate/hr/g. of liver $\pm$ S.E.M. (no. of animals)] were: mouse, 67 ± 2 (12); red-backed vole, 75 and 77; hamster, 26 ± 2 (7); rat, 18 ± 1 (18); guinea pig, 3.1 ± 0.5 (5); rabbit, 3.3 ± 1.6 (4); cat, 4.7 ± 0.4 (5); pig, sheep and ox, 1.6-2.8 (each 2). There is a fairly systematic acute rise in activity with diminishing body size.

This was not found with glutaminase activities (μ moles of glutamate/hr./g.): mouse, 319 ± 43 (8);

rat, 244 ± 6 (6); guinea pig, 190 ± 13 (5); cat, 380 ± 26 (4); pig, 245 and 113.

For glutamine synthetase it appears that activity (μ moles of glutamyl- γ -hydroxamate/hr./g.) is of a higher order in the smaller mammals: mouse, 517 ± 18 (4); red-backed vole, 224 ± 23 (4); rat, 398 ± 38 (6); guinea pig, 18 ± 3 (3); rabbit, 24 ± 2 (3); cat, 46 ± 2 (4); sheep, 52 ± 4 (3); goat, 58 ± 7 (3); pig 65 ± 16 (3); ox, 36 ± 9 (5).

The significance of these findings will be discussed in relation to the variation of basal metabolism with adult body size.

Buttery, P. J., Rowsell, E. V. & Carnie, J. A. (1969). *Biochem. J.* **115**, 42.

Cornelius, C. E. (1963). *Nature, Lond.*, **200**, 580.

Elliott, W. H. (1955). In *Methods in Enzymology*, vol. 2, p. 337. Ed. by Colowick, S. P. & Kaplan, N. O. New York: Academic Press Inc.

Lin, E. C. C., Rivlin, R. S. & Knox, W. E. (1959). *Amer. J. Physiol.* **196**, 303.

Rowsell, E. V. & Carnie, J. A. (1964). *Abstr. 6th int. Congr. Biochem., New York*, IV-144.

Rowsell, E. V., Carnie, J. A. & Wahbi, S. D. (1965). *Biochem. J.* **96**, 13P.

Wu, C. (1963). *Comp. Biochem. Physiol.* **8**, 335.

Investigation of the Cathepsins of Human Gastric Carcinomata

By D. J. ETHERINGTON* and W. H. TAYLOR. (*Department of Chemical Pathology, The United Liverpool Hospitals, Liverpool L3 5RT*)

Extracts of gastric carcinomata are devoid of pepsin (Greenstein & Stewart, 1942), but proteolytic activity near pH 3.5 has been demonstrated (Taylor, 1960a) that appears similar to the proteolytic activity of normal intestinal mucosa (Taylor, 1960b).

The construction of pH-activity curves with tumour extracts, by a modification of the Anson (1938) method of assay, gave a maximum ranging from pH 3.3 to 3.8 and curves that were asymmetrical. Electrophoresis of the extracts on agar gel at pH 8.2 revealed a cationic and at least one anionic proteolytic zone at positions corresponding respectively to cathepsins D and E of rabbit tissues (Lapresle & Webb, 1960, 1962).

Chromatography of one extract on DEAE-cellulose (Lapresle & Webb, 1962) gave three peaks of activity. The first peak (proteinase 1) eluted from the column was chromatographically and electrophoretically similar to rabbit cathepsin D. The pH-activity curve for this proteinase was symmetrical, with the maximum at pH 3.7. The second and third peaks (proteinases 2a and 2b) were only partially resolved, and emerged from the column

* Present address: Agricultural Research Council Meat Research Institute, Langford, Bristol BS18 7DY.

* Present address: Department of Applied Biochemistry and Nutrition, University of Nottingham.

at an equivalent position to cathepsin E. Asymmetrical, similar, pH-activity curves were obtained for both these proteinases, with the maxima at pH 3.4 to 3.5. After agar-gel electrophoresis, both proteinases were located towards the anode with proteinase 2b slightly ahead of proteinase 2a. Proteinase 1 was resolved further into two major (1a and 1b) and two minor peaks when chromatographed on CM-cellulose.

The sites of action of proteinases 1a and 1b on the B-chain of oxidized insulin were investigated at pH 3.5. Both cathepsins cleaved to completion bonds Tyr₍₁₆₎-Leu₍₁₇₎ and Phe₍₂₄₎-Phe₍₂₅₎ and both exhibited moderate action at bonds Leu₍₁₅₎-Tyr₍₁₆₎ and Phe₍₂₅₎-Tyr₍₂₆₎. Bonds Phe₍₁₎-Val₍₂₎, Glu₍₁₃₎-Ala₍₁₄₎ and Ala₍₁₄₎-Leu₍₁₅₎ were partially cleaved by proteinase 1a and only weakly attacked by proteinase 1b, whereas the bond at Gly₍₂₃₎-Phe₍₂₄₎ was more susceptible to hydrolysis by proteinase 1b.

Proteinases 1a and 1b have therefore similar specificities, and in their action they resemble two cathepsins D that have been previously investigated from bovine lung (Dannenberg & Smith, 1955) and swine thyroid (Kress, Peanasky & Klitgaard, 1966). Proteinases 2a and 2b have not been purified sufficiently for an investigation of their respective specificities.

Anson, M. L. (1938). *J. gen. Physiol.* **22**, 79.

Dannenberg, A. M. & Smith, E. L. (1955). *J. biol. Chem.* **215**, 55.

Greenstein, J. P. & Stewart, H. L. (1942). *J. nat. Cancer Inst.* **2**, 631.

Kress, L. F., Peanasky, R. J. & Klitgaard, H. M. (1966). *Biochim. biophys. Acta*, **113**, 375.

Lapresle, C. & Webb, T. (1960). *Biochem. J.* **76**, 538.

Lapresle, C. & Webb, T. (1962). *Biochem. J.* **84**, 455.

Taylor, W. H. (1960a). *J. clin. Path.* **13**, 349.

Taylor, W. H. (1960b). *Bull. Soc. Chim. biol., Paris*, **42**, 1313.

Regulation of Penicillinase Synthesis in *Bacillus licheniformis*: Binding of Inducer to Cells, and the Half-Life of the Messenger Ribonucleic Acid

By J. W. DAVIES. (*Department of Molecular Biology, University of Edinburgh, King's Buildings, Edinburgh 9*)

When cells of *Bacillus licheniformis* 749 are induced with penicillin, there is a slow increase in the differential rate of synthesis of penicillinase, the maximum rate being reached 60 min. or more after induction. This phenomenon, and the very slow deadaptation rate on removal of inducer, have encouraged speculation that the regulation mechanism may differ from other inducible enzymes.

Experiments interpreted as evidence for a long-lived messenger RNA led Harris & Sabath (1964) to suggest that the scheme proposed by Jacob & Monod (1961) is not applicable to the regulation of penicillinase in *Bacillus cereus*. An alternative model has been proposed (Csanyi, Jacobi & Straub, 1967) involving stabilization of the messenger after addition of inducer. It has been suggested that such a model can apply also to *B. licheniformis* (see Richmond, 1969) and that the slow deadaptation of penicillinase synthesis is a consequence of the long life of stabilized message. The binding of the inducer, [¹⁴C]penicillin, to *B. licheniformis* cells has been measured and the results showed a specific, irreversible, binding as in *B. cereus* (Pollock & Perret, 1951; Duerksen, 1963) and *Staphylococcus aureus* (Rogers, 1967), although 20–30-fold higher (approx. 5000 molecules/cell). Deadaptation by resuspending washed cells in a fresh medium does not remove the bound inducer, and thus the slow deadaptation can be explained without involving long-lived messenger. The existence of long-lived messenger has also been challenged. The enzyme-forming capacity of *B. licheniformis* cells was measured after stopping transcription with actinomycin D or rifampicin (1–5 µg./ml.). The half-life of the penicillinase messenger was estimated to be 4.7 min. for induced or constitutive cells. Although this value is twice the average half-life for total messenger RNA in this organism, there was no indication of an unusually long-lived messenger.

It is suggested that regulation of penicillinase production in *B. licheniformis* does not differ fundamentally from the β-galactosidase system of *Escherichia coli*. However, since the binding of inducer may be at the cell surface, it may require a secondary internal inducer to remove repressor from the DNA. Alternatively, if penicillin is the primary inducer, there may be a physical separation between inducer and repressor, which could explain the induction kinetics. A model suggesting this type of regulation, based on experiments with *B. cereus*, has recently been proposed (Imsande, 1969) and would seem to apply to *B. licheniformis*.

This work was supported by a Medical Research Council grant to the research group on Bacterial Enzyme Variation.

Csanyi, V., Jacobi, G. & Straub, B. F. (1967). *Biochim. biophys. Acta*, **145**, 470.

Duerksen, J. D. (1963). *Biochim. biophys. Acta*, **87**, 123.

Harris, H. & Sabath, L. D. (1964). *Nature, Lond.*, **202**, 1078.

Imsande, J. (1969). *J. Bact.* (in the Press).

Jacob, F. & Monod, J. (1961). *J. molec. Biol.* **3**, 318.

Pollock, M. R. & Perret, C. J. (1951). *Brit. J. exp. Path.* **32**, 387.

Richmond, M. H. (1969). In *Essays in Biochemistry*, vol. 4, p. 105. Ed. by Campbell, P. N. & Greville, G. D. London: Academic Press (Inc.) Ltd.

Rogers, H. J. (1967). *Biochem. J.* **103**, 90.