precipitated from liver and spleen is unknown. The values obtained with cortex and muscle probably represent the lowest sensitivity of the method with tissue extracts, as no specific binding of [6,7-³H]oestradiol-17 β to cortex was detected by gradient centrifugation; anterior hypothalamus, contained a 9S binding protein and, with it, more ³H radioactivity was precipitated by protamine (25%) than with cortex. After labelling of the 10⁵g supernatant from muscle with 1.0nM-[7 α -³H]-dehydroepiandrosterone sulphate or [7 α -³H]testosterone sulphate, 5% of the ³H radioactivity was precipitated by protamine, indicating the virtual absence of non-specific interaction between the polycation and steroid sulphates. After equilibration of 1.0nM-[6,7-³H]oestradiol-17 β with albumin, serum or phosphovitin, less than 5% of the ³H radioactivity was precipitated by protamine.

King, R. J. B., Gordon, J. & Steggles, A. W. (1969). *Biochem. J.* **114**, 649.

Scatchard, G. (1949). *Ann. N.Y. Acad. Sci.* **51**, 660.

Acetoacetate Formation in Bovine Tissues

By G. D. BAIRD and K. G. HIBBITT. (*Department of Biochemistry, Agricultural Research Council Institute for Research on Animal Diseases, Compton, Newbury, Berks.*)

Studies with preparations *in vitro* have demonstrated that the liver and the epithelial lining of the rumen wall are important sites of ketogenesis in the healthy and the ketotic ruminant (see Hibbitt & Baird, 1967). In the cow it has also been claimed, on the basis of observations made in the intact animal, that the mammary gland may also be ketogenic, at least during ketosis (Kronfeld, 1963).

Krebs, Wallace, Hems & Freedland (1969) have shown, in rat liver, that measurement of ketone-body production from slices may not give a true indication of ketogenic potential. An alternative to the use of slices and tissue pieces is to assay the activity of the four enzymes involved in acetoacetate formation from acetyl-CoA (acetoacetyl-CoA thiolase, HMG-CoA* synthase, HMG-CoA lyase and acetoacetyl-CoA deacylase) in homogenates of the tissues concerned. In the present study these enzymes were assayed by the methods of Williamson, Bates & Krebs (1968) in homogenates of bovine liver, rumen epithelium, mammary gland and kidney.

HMG-CoA lyase was assayed in only liver and rumen epithelium from healthy cows. The other three enzymes were assayed in each of the four tissues derived from healthy lactating cows and

from cows suffering from spontaneous ketosis. No significant differences were found in the activities of these three enzymes in the ketotic tissues as compared with the healthy tissues.

The activities of the enzymes in bovine liver were similar to those reported for the corresponding enzymes in rat liver, indicating that in the cow, as in the rat, acetoacetate formation proceeds mainly via the HMG-CoA pathway.

Rumen epithelium was found to contain thiolase, HMG-CoA synthase and HMG-CoA lyase, but no deacylase. It is therefore clear that acetoacetate formation must take place exclusively by way of HMG-CoA in this tissue. This observation is of some interest and demonstrates that the HMG-CoA pathway for acetoacetate synthesis is not confined to the liver in ruminants, as it appears to be in non-ruminants.

Thiolase and deacylase were both present in bovine kidney tissue, but not HMG-CoA synthase. Neither HMG-CoA synthase nor acetoacetyl-CoA deacylase was present in bovine mammary gland, although there was some thiolase activity.

Hibbitt, K. G. & Baird, G. D. (1967). *Vet. Rec.* **81**, 511.

Krebs, H. A., Wallace, P. G., Hems, R. & Freedland, R. A. (1969). *Biochem. J.* **112**, 595.

Kronfeld, D. S. (1963). *Ann. N.Y. Acad. Sci.* **104**, 799.

Williamson, D. H., Bates, M. W. & Krebs, H. A. (1968). *Biochem. J.* **108**, 353.

Lipolytic Activity in the Pancreas

By W. R. MILLER (introduced by D. A. HALL). (*Department of Medicine, University of Leeds, The General Infirmary, Leeds LS1 3EX*)

The major lipolytic activity of the pancreas is attributed to the pancreatic lipase, but the organ also contains a relatively high content of lipoprotein lipase, which preferentially degrades the triglyceride moiety of lipoproteins, in contrast with pancreatic lipase, which acts on emulsified triglyceride. Although different reactivity to inhibitors has been reported for the two enzymes, LeVeen, Giordano & Zinberg (1961) and Kessler, Finkel, Dreiling & Janowitz (1963) have been unable to ascertain whether the lipoprotein lipase activity of the pancreas is a secondary property of the conventional lipase or is due to the presence of a separate enzyme.

In the present study the lipolytic activities of two different preparations from the pancreas have been measured by using as substrates a commercial triglyceride emulsion both in the free and albumin-activated state for determination of triglyceridase and lipoprotein lipase activity respectively.

Each preparation exhibited the properties of both enzymes, being inhibited by NaF and activated by

*Abbreviation: HMG-CoA, β -hydroxy- β -methylglutaryl-CoA.

bile salts (pancreatic lipase characteristics) and being inhibited by protamine and pyrophosphate (lipoprotein lipase characteristics). 1M-NaCl either activated, inhibited or had no effect, depending on the conditions of assay. By the use of specific inhibitors for one or other enzyme, the differing optimum pH values for each of the two types of lipolytic activity were demonstrable. From these observations it might be assumed that both preparations consisted of a mixture of two separate enzymes. However, incubation with NaF and protamine in combination did not result, as would be expected in this case, in complete inhibition, but merely in the degree of inhibition equal to that caused by protamine alone. Similarly, in the presence of Ca^{2+} and albumin, where 1M-NaCl gave 40% inhibition (characteristic of lipoprotein lipase) and fluoride 65% inhibition (characteristic of pancreatic lipase), the inhibition in the combined presence of NaF and 1M-NaCl was only 69%. These observations suggest that the two types of lipolytic activity are mediated by a single enzyme.

Further evidence for this was obtained by using Sephadex chromatography. Each pancreatic preparation gave a single chromatographic peak with a molecular weight of 800 000 when run through Sephadex G-200, but after acetone treatment a delipidized product in each case gave a second peak of much smaller molecular weight (15 000–20 000). This molecule also demonstrated the mixed lipolytic activities associated with the parent molecule.

Kessler, J. I., Finkel, M., Dreiling, D. A. & Janowitz, H. D. (1963). *Proc. Soc. exp. Biol., N.Y.*, **113**, 127.

LeVeen, H. H., Giordano, P. & Zinberg, S. (1961). *Physiologist*, **4**, 65.

Fat-Body Lipogenesis during Development of the Adult Male Desert Locust

By P. R. WALKER and E. BAILEY. (*Department of Biochemistry, University of Sheffield*)

Changes in fat-body lipogenesis measured by acetate incorporation into lipid and changes in associated enzymes have been followed during development of the adult male desert locust (*Schistocerca gregaria*). Lipid accumulates in the fat-body during the first 8 days after emergence, the major period of accumulation occurring between the sixth and the eighth day. There is a rapid increase in $[\text{U-}^{14}\text{C}]$ acetate incorporation into lipid between the fifth and seventh day after a period of intensive feeding between emergence and 5 days. Similar results were obtained for $[\text{U-}^{14}\text{C}]$ glucose, $[\text{U-}^{14}\text{C}]$ trehalose and $[1,5\text{-}^{14}\text{C}_2]$ citrate incorporation into lipid. The incorporation of $[1,5\text{-}^{14}\text{C}_2]$ citrate into lipid implies the functioning of the citrate-

cleavage pathway in the conversion of carbohydrate into lipid.

'Malic' enzyme, which is considered to be important in the production of NADPH for lipogenesis in mammalian tissues (Kornacker & Ball, 1965), increases in activity between emergence and 4 days, exhibiting a pronounced peak at 4 days. This pattern is similar to those of phosphofructokinase and pyruvate kinase, which both increase in activity after emergence, and phosphofructokinase in particular has a pronounced peak of activity at 4 days. Isocitrate dehydrogenase (NADP-dependent) and glucose 6-phosphate dehydrogenase, which may also be important in the production of NADPH for lipogenesis, increase in activity between emergence and 2 days, the increased activity being maintained until after 8 days. ATP citrate lyase is the key enzyme in the citrate-cleavage pathway. The activity of this enzyme is constant between emergence and 4 days and then increases to give a peak of activity between the sixth and the eighth day. The developmental pattern for this enzyme is very similar to the pattern of acetate incorporation into lipid.

The scheme of Kornacker & Ball (1965) involving coupling of ATP citrate lyase and 'malic' enzyme requires parallel changes in activity of these enzymes during changes in lipogenesis. Since there is no correlation between the changes in activity of ATP citrate lyase and 'malic' enzyme during development it is unlikely that this pathway operates in the locust. 'Malic' enzyme may function to produce malate from pyruvate during increased glycolysis. Intramitochondrial oxaloacetate derived from the malate could be used to condense with acetyl-CoA to provide citrate for lipogenesis. NADPH for lipogenesis may be provided by glucose 6-phosphate dehydrogenase and isocitrate dehydrogenase.

The authors thank the Medical Research Council for a grant to P.R.W.

Kornacker, M. S. & Ball, E. G. (1965). *Proc. nat. Acad. Sci., Wash.*, **54**, 899.

Rat Liver β -Xylosidase, a Lysosomal Membrane Enzyme

By D. FISHER* and P. W. KENT. (*Department of Biochemistry, University of Oxford*)

β -Xylosidase activity, as determined with *o*-nitrophenyl and *p*-nitrophenyl β -D-xylopyranosides, is widely distributed in animal tissues

* Present address: Department of Biochemistry, Royal Free Hospital School of Medicine, London W.C.1.