radioactive peak was most readily obtained when Nonidet (NP 40) or BRIJ 58 was used to disrupt the mitochondria after pulse-labelling, but similar ribosome-like particles were also obtained by use of sodium deoxycholate or Triton X-100. Mitochondria disrupted in very low Mg^{2+} concentrations showed radioactivity to about 50S. The results are entirely consistent with the assumption that animal mitochondria contain mitoribosomes sedimenting at less than 80S. Experiments in which radioactive mitoribosomes were mixed with unlabelled *E. coli* ribosomes and sedimented in a sucrose density gradient containing 10mM- Mg^{2+} showed that the two peaks (radioactivity and extinction) were close together whereas *E. coli* ribosomes were readily separable under these conditions from rat liver cytoribosomes and Krebs-cell cytoribosomes. These results reinforce the suggestion that in protein synthesis mitochondria resemble bacteria much more closely than they do the eukaryotic cytoplasmic ribosomes.

O'Brien, T. W. & Kalf, G. W. (1967a). *J. biol. Chem.* **242**, 2172.

O'Brien, T. W. & Kalf, G. W. (1967b). *J. biol. Chem.* **242**, 2180.

Work, T. S., Coote, J. L. & Ashwell, M. A. (1968). *Fedn Proc. Fedn Am. Soc. exp. Biol.* **27**, 1174.

Factors Affecting Protein Synthesis in Yeast Mitochondria

By L. A. GRIVELL and D. B. ROODYN. (*Department of Biochemistry, University College of London, London W.C.1, U.K.*)

Mitochondria isolated essentially free of micro-organisms from protoplast lysates of *Saccharomyces carlsbergensis* have been found to incorporate [^{14}C]leucine into protein during incubation *in vitro* (Grivell, 1967). Incorporation takes place mainly into an RNA-rich fraction, possibly located within the inner mitochondrial membrane, and is sensitive to chloramphenicol and erythromycin, but resistant to cycloheximide and ribonuclease. Activity requires K^+ , Mg^{2+} and phosphate and has a broad optimum at pH 6.7; Li^+ and Na^+ are inhibitory. Response to Mg^{2+} is dependent on the phosphate concentrations and the adenine nucleotide present. The system requires ATP, which may be exogenous or generated by respiratory phosphorylation in the presence of oxidizable substrate and ADP; the phosphoenolpyruvate-pyruvate kinase ATP-regenerating system is fully effective only when succinate is present. Experiments with oligomycin suggest that there is not an obligatory participation of the hypothetical high-energy intermediates of oxidative phosphorylation.

Transport of [^{14}C]leucine has also been measured by a centrifugation-filtration technique; there is approximately twofold concentration of amino acid in a process that appears to be complete within 20s of addition of mitochondria to the incubation medium and that is not markedly affected by the addition of oxidizable substrate, ATP, oligomycin or antimycin. It is concluded that there is not a simple relationship between the ability of the mitochondrion to accumulate amino acids and to synthesize protein.

Direct measurement of the changes occurring in the concentrations of mitochondrial amino acids during incubation reveal discrepancies between radioactive and chemical methods of estimation of concentration ratios; equilibration between [^{14}C]leucine and the endogenous pool is not complete even after 30min. Although the size of the endogenous leucine pool does not increase during incubation, there is an approximately threefold increase in exogenous leucine. The difficulties of assessing true rates of protein synthesis in a system where the amino acid pool is not homogeneous and increases during incubation are to be stressed.

The authors are grateful to the Science Research Council for financial support and for the award of a Research Scholarship to L. A. G.

Grivell, L. A. (1967). *Biochem. J.* **105**, 44c.

Ribosome Particles in Yeast Mitochondria

By P. V. VIGNAIS and J. HUET. (*Biochimie, Faculté de Médecine et Centre d'Études Nucléaires, 38-Grenoble, France*) and J. ANDRÉ (introduced by D. E. GRIFFITHS). (*Biologie Cellulaire 4, Faculté des Sciences, 91-Orsay, France*)

Evidence for the existence of ribosomes within mitochondria from yeast cells (*Candida utilis*, C.B.S. 1516) has been obtained from examination of electron micrographs of isolated mitochondria, and also from the characterization of the mitoribosomes after their isolation from mitochondria.

Electron micrographs of thin sections of yeast mitochondria stained with uranyl acetate and lead citrate showed rows of particles with diameters of 180 Å attached along the matrix side of the inner membrane and of the cristae. Free mitoribosomes were obtained by disruption of mitochondria with deoxycholate and purified by centrifugation on a linear 5–20% (w/v) sucrose density gradient in 5mM- $MgCl_2$ –50mM-KCl–10mM-tris–HCl buffer, pH 7.6. The sedimentation profile of mitoribosomes obtained under the above conditions is characterized by a major peak at 53–55S and a smaller one at 76–78S. Increasing the $MgCl_2$ concentration in the sucrose density gradient above 20mM results in an increase of the 76–78S peak with a slight shift to