PROCEEDINGS OF THE BIOCHEMICAL SOCIETY

The 483rd Meeting of the Society was held at the University College of North Wales, Bangor, on Thursday, Friday and Saturday, 20, 21 and 22 June, when the following papers were presented:

COLLOQUIUM ON 'SOME ASPECTS OF SOIL BIOCHEMISTRY'

The Nature of Soil Humic Acid

By R. D. HAWORTH (*Department of Chemistry, University of Sheffield*)

The results of spectroscopic and chemical degradation studies on acid-boiled humic acid (ABHA) will be discussed.

The electron-spin-resonance spectra of ABHA in alkaline solution are of two main types corresponding to the pH of the soil from which the acid is derived, and although structural deductions cannot be made the spectra have useful ecological applications. The free radical responsible for the electron-spin-resonance spectra has the properties of a semiquinone ion radical, and the remarkable stability is best explained by its association with extensive conjugation.

Degradation experiments involving distillation with zinc dust gave small yields of an oil shown by u.v. and mass spectrometry to contain numerous aromatic hydrocarbons, of which coronene was the most complex representative detected.

Reduction of ABHA with hydroiodic acid gave an alicyclic oil which on dehydrogenation with palladium-charcoal gave a mixture containing pyrene, a methylpyrene and perylene.

Permanganate oxidation of reduced and methylated ABHA gave an acid mixture, which was converted by a series of reactions into a neutral product containing a methylpyrene (identified by u.v. and mass spectra).

It is suggested that humic acid is composed of an aromatic 'core' responsible for the electron-spin-resonance spectra, to which polysaccharides, proteins, metals and relatively simple phenols are attached by chemical or physical means or both.

Some observations concerning the protein linkages and the mechanism of formation of humic acid will be made.

Transformations of some Labelled Carbohydrates in Soil

By M. V. CHESHIRE, C. M. MUNDIE and H. SHEPHERD (*Macaulay Institute for Soil Research, Aberdeen*)

The sugar composition of the polysaccharide fraction of soil organic matter varies little with soil type (Gupta, Sowden & Stobbe, 1963) and is not much affected by incubation under laboratory conditions. It is not known whether it represents mainly resistant plant residues or material newly synthesized by microbial action.

To investigate this point, radioactive starch or glucose, sufficient to raise the glucose content two- or three-fold, was incubated with moist soil for several months, and the transfer of label from glucose to other sugars studied. For this purpose it was necessary to devise methods by which the greater part of the carbohydrate could be extracted from soil (Cheshire & Mundie, 1966), and the constituent sugars separated for measurement of specific radioactivities (Mundie, Cheshire & Bacon, 1968). Attention was concentrated on the major sugar constituents: glucose, galactose, mannose, xylose, arabinose, fucose and rhamnose.

After 2 weeks, only 20% of the added radioactivity could be recovered in the soil hydrolysate, and in 5 weeks the carbohydrate content of the soil returned almost to its initial value. Glucose remained the most highly labelled sugar even after 12 weeks; significant amounts of radioactivity were present in the other hexoses (including 6-deoxyhexoses), but very little in the pentoses.

More soils will have to be examined, and incubation conditions varied, before it can be concluded that the soil pentoses are not of microbial origin.

Cheshire, M. V. & Mundie, C. M. (1966). *J. Soil Sci.* **17**, 372.
Gupta, U. C., Sowden, F. J. & Stobbe, P. C. (1963). *Proc. Soil Sci. Soc. Amer.* **27**, 380.
Mundie, C. M., Cheshire, M. V. & Bacon, J. S. D. (1968). *Biochem. J.* in the Press.