sap, whereas it stimulated when ascites ribosomes were used in the presence of their own sap. When reticulocyte sap was incubated with ascites cell ribosomes EMC-RNA had no significant effect.

One interpretation of these results is that ascites-cell sap contains a factor absent from reticulocyte-cell sap which is necessary for the translation of the added EMC-RNA. It seems that reticulocyte ribosomes are able to combine with added RNA in such a way that incorporation on the endogenous message is decreased.

We thank the Science Research Council for a Studentship for M. B. M., Dr A. T. H. Burness and the Medical Research Council for the facilities to grow the virus, and the British Empire Cancer Campaign for Research for financial support.

Kerr, I. M., Cohen, N. & Work, T. S. (1966). *Biochem. J.* **98**, 826.

Variations in the Microstructure of Histones during Early Cleavage Stages in *Echinus*

By MARGERY G. ORD and L. A. STOCKEN (*Department of Biochemistry, University of Oxford*)

Earlier work on nuclear proteins from sea-urchins indicated that in the mature unfertilized egg the thiol group(s) on the histone(s) was present in the disulphide form and that an increase occurred in the proportion of thiol after fertilization, reaching a peak just before cleavage (Ord & Stocken, 1968). These findings have now been confirmed and extended by using *Echinus esculentus* and histone extracted from washed nuclei in ethanol-1.25M-HCl (4:1, v/v) (Johns, 1964). Fertilization and subsequent incubation of the eggs were performed at 8°; the first furrow appeared at 180min. after fertilization. The eggs were collected at 10-20min. intervals from fertilization to the second cleavage. The packed eggs were stored at -15° for about a fortnight before extracting the histones. The acid-ethanol extracts were precipitated immediately with 5vol. of acetone-HCl and were dialysed against 50mM-HCl and water at 2° before analysis. Only traces of lysine-rich histones are present in *Echinus* in early cleavage stages; the material analysed contained components migrating on electrophoresis on polyacrylamide gel in positions similar to that of the arginine-rich histone F3 and of the F2A group. Neither phosphate nor thiol is present in the latter histones so that the SH+S·S and phosphate contents are predominantly accounted for by histone F3 (Ord & Stocken, 1967).

The total SH+S·S content of the histones analysed did not vary (69.3 ± 13.0 nmol/mg. of protein), but the proportion of thiol increased from less than 10 to more than 50% at 140, 180 and 275min. after fertilization. From work on thymi-

dine uptake into *Strongylocentrotus purpuratus* at 15° (Hinegardner, Rao & Feldman, 1964), these times probably correspond to periods of DNA synthesis. As reported earlier there is also a sharp peak in the percentage of SH 30min. after fertilization that is not found in subsequent cycles. The phosphate content of the proteins (53.6 ± 5.9 ng.atoms of P/mg. of protein) remained constant for 10-90min. after fertilization; preliminary results indicate it may then vary reciprocally to the proportion of thiol.

Our thanks are due to the Director, Mr R. I. Currie, and his colleagues at the Scottish Marine Biological Research Station, Millport, where the material was collected.

Hinegardner, R. T., Rao, B. & Feldman, D. E. (1964). *Exp. Cell Res.* **36**, 53.

Johns, E. W. (1964). *Biochem. J.* **92**, 55.

Ord, M. G. & Stocken, L. A. (1967). *Biochem. J.* **102**, 631.

Ord, M. G. & Stocken, L. A. (1968). *Biochem. J.* **107**, 403.

Histone Phosphorylation and Cell Division

By W. S. STEVELY and L. A. STOCKEN (*Department of Biochemistry, University of Oxford*)

An approximate temporal correlation between the phosphorylation of histone F1 and DNA synthesis in regenerating rat liver has been reported by Ord & Stocken (1967). More detailed work has now shown that the increase in phosphorylation of histone F1 precedes the incorporation of ^{32}P into DNA by approx. 1hr.

Partially hepatectomized and control sham-operated animals were injected with $100\mu\text{C}$ of [^{32}P]phosphate/100g. body wt. intramuscularly 30min. before death. Groups of three or four rats were killed at $\frac{1}{2}$ hr. intervals over the period 15-19hr. after hepatectomy. Histone F1 was obtained from washed nuclei by extraction with perchloric acid (Johns, 1964) and its phosphate content determined by the method of Bartlett (1959). ^{32}P incorporation into DNA was measured by the method of DeLuca, Rossiter & Strickland (1953). Tissue phosphate soluble in 20% trichloroacetic acid was used as a reference standard for determining relative specific radioactivities.

A further indication of a link between DNA synthesis and phosphorylation of histone F1 comes from the finding that 1000 rads total body γ -irradiation at 12hr. after partial hepatectomy prevents the phosphorylation of histone F1 in a manner similar to the inhibition of DNA synthesis (Holmes & Mee, 1955).

Unilaterally nephrectomized rats have also been found to give increased phosphorylation of histone F1 24hr. after the operation when DNA synthesis is known to be taking place. 1000 rads total body