phate pathway in pancreatic β -cells may be related to the secretion of insulin. Thus a high activity of the enzymes of this pathway has been demonstrated in islet cells (Kissane, Lacy, Brolin & Smith, 1964). Further studies from this Laboratory (Montague, Howell & Taylor, 1967) have shown that certain pentoses and pentose derivatives, known to be metabolized through the pentose phosphate pathway, are as effective as glucose in promoting insulin secretion.

The mechanism by which glucose may release insulin has now been investigated by measuring the concentrations of glucose 6-phosphate and 6-phosphogluconate in islet cells actively secreting in response to various agents.

Isolated islets of Langerhans prepared by collagenase digestion of rat pancreas (Howell & Taylor, 1968) were incubated with various agents known to affect insulin secretion. After 30 min. incubation, the medium was removed from the islets and its insulin content determined by immunoassay. The islets were rapidly cooled to -76° and then disrupted ultrasonically after the addition of perchloric acid. The concentrations of glucose 6-phosphate and 6-phosphogluconate in the perchloric extracts were estimated by using the enzymic cycling procedure described by Lowry, Passonneau, Hasselberger & Schultz (1964).

When the glucose concentration of the incubation medium was raised from 2 to 20 mM, the concentrations of glucose 6-phosphate and 6-phosphogluconate in the islets were markedly increased. In addition, a number of other agents also raised the concentration of 6-phosphogluconate. These included tolbutamide, citrate, glucagon and theophylline, all agents known to promote insulin secretion. In addition, adrenaline ($1\mu\text{M}$) as well as inhibiting the secretion of insulin in response to 20 mM-glucose, also caused a significant decrease in the concentration of 6-phosphogluconate. Citrate and tolbutamide were also found to increase the concentration of glucose 6-phosphate in the islets.

Thus there seems to be a close parallelism between the concentration of 6-phosphogluconate and the secretion of insulin by islets. It is suggested that glucose is normally metabolized through the pentose phosphate pathway in islet cells in order to stimulate the secretion of insulin.

Howell, S. L. & Taylor, K. W. (1968). *Biochem. J.* **108**, 17.
Kissane, J. M., Lacy, P. E., Brolin, S. E. & Smith, C. H. (1964). In *The Structure and Metabolism of the Pancreatic Islets*, p. 281. Ed. by Brolin, S. E., Hellman, B. & Knutson, H. Oxford, Pergamon Press Ltd.

Lowry, O. H., Passonneau, J. V., Hasselberger, F. X. & Schultz, O. W. (1964). *J. biol. Chem.* **239**, 18.
Montague, W., Howell, S. L. & Taylor, K. W. (1967). *Nature, Lond.*, **215**, 1088.

Insulin Biosynthesis: Studies of Ribonucleic Acid Metabolism *in vitro* in Isolated Surviving Mammalian and Fish Islets

By N. S. TRACK, R. J. JARRETT and H. KEEN
(Department of Medicine, Guy's Hospital Medical School, London, S.E. 1)

The present studies were designed to elucidate the role of RNA in insulin biosynthesis in rat islets prepared by microdissection after the pancreatic ducts had been ligated and atrophy of the acinar tissue had been allowed to occur (Keen, Sells & Jarrett, 1965), and in cod islets prepared by excision of the distinct islet organ (Grant & Reid, 1968).

Glucose concentration has a stimulatory effect on the incorporation of labelled precursors into rat islet RNA (Jarrett, Keen & Track, 1967); this effect was not found in cod islet experiments with various glucose concentrations. Rat but not cod islet RNA synthesis was inhibited by actinomycin D. Sucrose-gradient-ultracentrifugation analysis of the rat islet RNA demonstrated that incorporated radioactivity was present in the low-molecular-weight fraction. To establish the *S* value of this labelled material samples of both rat- and cod-islet RNA were separated on polyacrylamide gels. The labelled RNA had a mobility on the gels that indicated that the corresponding *S* value had a range 3.6–4.0S.

The structure of insulin permits calculation of the hypothetical molecular weight of the messenger-RNA (m-RNA) that would 'code' for either the two chains separately or one single polypeptide chain. The isolation (Steiner & Oyer, 1967; Steiner, Cunningham, Spigelman & Aten, 1967; Tung & Yip, 1968; Grant & Reid, 1968) and sequence determination (Chance & Ellis, 1968) of an insulin precursor molecule support the single-polypeptide-chain mode of synthesis, which would require only one m-RNA. As the precursor seems to be composed of approximately 80 amino acids, it would require a m-RNA with an *S* value larger than that of the labelled material found in both rat and cod experiments.

These results suggest that the insulin m-RNA may have a long half-life and that transfer-RNA (t-RNA) modulation (Stent, 1964) may be one of the factors regulating insulin biosynthesis; this would explain the observed *S* values.

Monjardino & MacGillivray (1968) have shown that this type of mechanism seems to be operating in human lymphocytes.

Experiments are in progress with [^3H]methyl-methionine in an attempt to determine whether it is in fact t-RNA that is being labelled in the incubations.

N.S.T. is aided by a grant from the Foundation Fund of

the Canadian Diabetic Association. All the cod islet experiments were carried out in Dr P. T. Grant's Department.

- Chance, R. & Ellis, R. (1968). *Science* (in the Press).
 Grant, P. T. & Reid, K. M. B. (1968). *Biochem. J.* (accompanying communication).
 Jarrett, R. J., Keen, H. & Track, N. (1967). *Nature, Lond.*, **213**, 634.
 Keen, H., Sells, R. & Jarrett, R. J. (1965). *Diabetologia*, **1**, 28.
 Monjardino, J. P. P. V. & MacGillivray, A. J. (1968). *Biochem. J.* (in the Press).
 Steiner, D. F. & Oyer, P. E. (1967). *Proc. nat. Acad. Sci., Wash.*, **57**, 473.
 Steiner, D. F., Cunningham, D., Spigelman, L. & Aten, B. (1967). *Science*, **157**, 697.
 Stent, G. S. (1964). *Science*, **144**, 816.
 Tung, A. K. & Yip, C. C. (1968). *Diabetologia*, **4**, 68.

Biosynthesis of an Insulin Precursor by Cod Islet Tissue and the Occurrence of Intermediates in the Conversion to Insulin

By P. T. GRANT and K. B. M. REID (N.E.R.C. Fisheries Biochemical Research Unit, University of Aberdeen)

Cod insulin contains four proline residues (Grant & Reid, 1968). The incorporation of [U-¹⁴C]proline by slices of cod islet tissue at 15° into protein soluble in acid ethanol was approximately linear for 6 hr., and was inhibited by puromycin but slightly stimulated by actinomycin D. The radioactivity of protein fractions in these extracts was determined in view of the conclusion by Steiner, Cunningham, Spigelman & Aten (1967) that insulin is formed in rat and human islets from a single polypeptide chain precursor.

The radioactivity of an extract after 1 hr. was present in only one fraction (fraction P1). The molecular weight of this fraction was about 10200, and it was appreciably more basic than insulin as judged by gel filtration on Sephadex G50 and chromatography on DEAE-cellulose as described by Grant & Reid (1968). Longer periods of incubation increased the radioactivity of fraction P1 and after 1 hr. there was a detectable and progressive increase in the radioactivity both of insulin and of fractions intermediate in molecular size and basicity between fraction P1 and insulin.

Slices were incubated with [U-¹⁴C]proline for 1 hr. to label fraction P1, *O*-*p*-nitrophenyl *O*-ethyl phenylpropylphosphonate (Becker, Fukuto, Boone, Canham & Boger, 1963) was added, and the incubation was continued for a further 4 hr. This tryptic and chymotryptic inhibitor had little effect on the biosynthesis of fraction P1, but there was a marked decrease in the radioactivity associated with insulin and the intermediate fractions. In a similar experiment, puromycin or an excess of unlabelled

L-proline resulted in the radioactivity of fraction P1 being less than that associated with insulin and the intermediate fractions.

Treatment of a radioactive preparation of the single-chain polypeptide, fraction P1, with trypsin (150:1), resulted in a progressive increase of biological activity in the digest as measured by the rat fat-pad bioassay, and of the radioactivity in intermediates and insulin fractions separated on DEAE-cellulose.

These results are consistent with the concept that insulin formation involves the biosynthesis of a precursor polypeptide, but the conversion of precursor into product in cod islet tissue appears to involve the cleavage of peptide bonds, resulting in the formation of a number of intermediates that in the present procedures are poorly separated from one another and to a certain extent from insulin.

- Becker, E. L., Fukuto, T. R., Boone, B., Canham, D. C. & Boger, E. (1963). *Biochemistry*, **2**, 72.
 Grant, P. T. & Reid, K. B. M. (1968). *Biochem. J.* **106**, 531.
 Steiner, D. F., Cunningham, D., Spigelman, L. & Aten, B. (1967). *Science*, **157**, 697.

Rapid Effects of Testosterone on Prostatic Polyamine-Synthesizing Enzyme Systems

By A. E. PEGG and H. G. WILLIAMS-ASHMAN (Department of Pharmacology and Brady Urological Institute, Johns Hopkins School of Medicine, Baltimore, Md., U.S.A.)

The prostate glands and their secretions in many species contain high concentrations of spermidine and spermine (Mann, 1964; Tabor & Tabor, 1964). The ability of spermidine and spermine to interact with polynucleotides, ribosomes and phospholipids, and also to influence the enzymic synthesis of DNA, RNA and proteins (Tabor & Tabor, 1964; Raina & Telaranta, 1967) prompted us to examine the activity of some enzymes involved in polyamine biosynthesis in relation to androgen-induced growth of the ventral prostate in orchietomized rats. During the latter process, nuclear RNA polymerase activity is increased within 1 hr. after injection of testosterone (Liao, Leininger, Sagher & Barton, 1965), whereas DNA polymerase activity exhibits large but transitory elevations between the second and the fifth day after commencement of daily androgen treatments (Coffey, Shimazaki & Williams-Ashman, 1968).

Putrescine is formed in rat prostate by action of a pyridoxal phosphate-dependent L-ornithine decarboxylase (Pegg & Williams-Ashman, 1968a). Spermidine is synthesized from S-adenosyl-L-methionine (SAM) and putrescine. The decarboxylation of SAM by soluble prostatic extracts is markedly and specifically stimulated by putrescine;