Sporulation in *Bacillus subtilis*

MORPHOLOGICAL CHANGES

By D. KAY

Sir William Dunn School of Pathology, University of Oxford

AND S. C. WARREN*

Microbiology Unit, Department of Biochemistry, University of Oxford

(Received 25 March 1968)

1. When *Bacillus subtilis* was grown in a medium in which sporulation occurred well-defined morphological changes were seen in thin sections of the cells. 2. Over a period of 7.5 hr. beginning 2 hr. after the initiation of sporulation the following major stages were observed: axial nuclear-filament formation, spore-septum formation, release of the fore-spore within the cell, development of the cortex around the fore-spore, the laying down of the spore coat and the completion of the corrugated spore coat before release of the spore from the mother cell. 3. The appearance of refractile bodies and 2,6-dipicolinic acid and the development of heat-resistance began between 5 and 6.5 hr. after initiation of sporulation. 4. The appearance of 2,6-dipicolinic acid and the onset of refractility appeared to coincide with a diminution of electron density in the spore core and cortex. 5. Heat-resistance was associated with the terminal stage, the completion of the spore coat. 6. The spore coat was composed of an inner and an outer layer, each of which consisted of three or four electron-dense laminae. 7. Serial sections through cells at an early stage of sporulation showed that the membranes of each spore septum were always continuous with the membranes of a mesosome, which was itself in close contact with the bacterial or spore nucleoid. 8. These changes were correlated with biochemical events occurring during sporulation.

The principal stages in the development of the spore have been described for *Bacillus cereus* by Young & Fitz-James (1959), and for *Bacillus subtilis* by Ryter (1965). The process appears to be similar in both organisms and, for the latter, eight morphological stages have been described (Schaeffer, Ionesco, Ryter & Balassa, 1965), starting with the vegetative form (stage 0). Stage I consists of the formation of an axial chromatic filament, which is followed (stage II) by the development of the spore septum. During stage III the spore membrane develops and the fore-spore is released inside the mother cell. Stage IV is represented as the formation of the cortex and stage V as the development of the spore coat. During stage VI the spore matures to give an intact included spore, which is finally released as the mother cell autolyses (stage VII).

The present studies were undertaken in an attempt to reveal more of the detailed morphology of sporulation and also in the hope that it would be possible to determine an approximate time-scale

for the process. This would provide the basis for an empirical correlation between the sequence of biochemical events that occur during sporulation (Warren, 1968) and the morphological development of the spore (cf. Murrell, 1967).

MATERIALS AND METHODS

Organism and culture. *B. subtilis* (Marburg strain 168), which requires indole or tryptophan for growth, was used throughout. A loopful of spore suspension was inoculated into Difco Penassay Broth (Difco Antibiotic Medium 3) (1 ml.) to which additional glucose (0.4%, w/v) had been added, and the culture shaken for 6 hr. at 37°. A tenfold dilution of this culture (1 ml.) was used to inoculate 11. batches, in 51. conical flasks, of a liquid sporulation medium (Warren, 1968) that was a 2.5-fold concentration of the medium of Donnellan, Nags & Levinson (1964). These cultures were shaken overnight at 37° and 51. of the batches, whose bacterial densities varied in the range 0.40–0.475 mg. dry wt./ml., were pooled and distributed in 500 ml. lots in 31. conical flasks. These flasks were shaken at 37° and single flasks harvested at 1.5 hr. intervals over a period of 7.5 hr. Samples (2.5 ml.) were removed for electron microscopy and phase-contrast microscopy.

*Present address: Unilever Research Laboratories, Colworth House, Sharnbrook, Bedford.

Assay methods. Determination of DPA* and numbers of heat-resistant spores were made as described by Warren (1968) and the remainder of the culture was used for the biochemical assays reported in the preceding paper (Warren, 1968).

Electron microscopy. Samples (2.5 ml.) were taken from the flask and centrifuged, and the cells fixed by the method of Kellenberger, Ryter & Séchaud (1958). The pellet of cells was dehydrated in graded ethanol and embedded in Araldite.

Serial sections were cut from blocks that had been trimmed to give truncated pyramids about 0.1 mm. across. The ribbons were collected on Formvar-coated hexagonal grids (Athene; Smethurst High-Light Ltd., Bolton, Lancs.) to minimize the chance of the required areas of any section lying behind a grid bar. Up to 15 sections could be observed in the microscope and this was sufficient for particular cells to be followed right through.

Both single and serial sections were stained with lead citrate (Reynolds, 1963).

RESULTS

Spore formation is considered to begin at the end of the period of exponential growth and is indicated as t_0 in Fig. 3 in the preceding paper (Warren, 1968). Samples were taken at 1.5 hr. intervals beginning 2 hr. after initiation of sporulation, i.e. at t_2 .

The times of appearance of refractility, DPA and heat-resistance are shown in Table 1.

Over the 7.5 hr. period of observation most of the cells showed either complete or partial spore formation, but it was clear that the culture was far from synchronous in the initiation of spore formation. No spores were seen in the first samples (zero time) and the first appearance of pre-spores was observed in the 3 hr. specimens, but in this sample some cells had still only reached the stage of initial

*Abbreviation: DPA, 2,6-dipicolinic acid.

Table 1. *Appearance of refractile bodies, DPA and heat-resistance during sporulation*

Samples (50 ml.) of sporulating cultures were removed from flasks harvested at the times indicated and 1 drop of each was examined in the phase-contrast microscope. DPA and the number of heat-resistant cells were determined as described in the Materials and Methods section. N.D., Not detectable.

Sample	Time	Cells with refractive bodies (%)	DPA content (μ mole/mg.)	No. of heat-resistant cells
2	$t_{3.5}$	N.D.	N.D.	1.5×10^3
3	t_5	N.D.	N.D.	1.5×10^3
4	$t_{6.5}$	5	0.0126	1.5×10^8
5	t_8	70	0.0272	7.5×10^8
6	$t_{9.5}$	85	0.0419	1.5×10^9
	t_{24}	100	0.1388	5.0×10^9

septum formation. The cells in Plates 1 and 2 have been chosen to illustrate successive and well-marked morphological stages in spore formation, but they are not necessarily taken from samples obtained in chronological order.

Plate 1(a) shows the cell in the condition described as stage I by Schaeffer *et al.* (1965) in which the nuclear material (A) was formed into an axial filament. The first major structural change (stage II) in the cell is shown in Plate 1(b). A spore septum (B) has formed across the cell close to the pole. It is clearly distinguishable from the division septum (C), which is thicker and generally forms close to the midpoint of the long axis of the cell. A few cells showed the spore septa after 1.5 hr. The section in Plate 1(b) was cut parallel to the long axis of the cell but close to the cell wall and so no nuclear material is visible.

Stages in the formation of the spore septa are shown in Plates 3 and 4, where serial sections through three cells with five presumptive spore septa are illustrated in the order in which they were

EXPLANATION OF PLATES 1 AND 2

Stages in the development of spores in *B. subtilis* incubated in a defined medium (see the Materials and Methods section for conditions of growth and preparation of sections for electron microscopy. (a) One cell of a pair, showing the earliest sign of the onset of spore formation. The bacterial DNA (A) is concentrated axially. (b) Formation of the initial spore septum (B). This is thinner than the division septum (C). The mode of formation of the spore septum is shown in more detail in the serial sections (Plates 3 and 4). (c) The spore septum (B) is extending towards the centre of the cell. An area containing a mesosome (D) and some DNA has been cut off from the main body of the cell. (d) The spore membrane (E), now complete, is separated from the cytoplasmic membrane (F). A mesosome (D) is connected to the inner leaf of the spore membrane. Another mesosome (D1) is pressed against the end of the cell by the developing cortex of the spore. (e) The spore is now surrounded by the cortex (G) and the earliest trace of the outer membrane (H). The cytoplasmic ribosomes (I) are pressed back by the developing spore. The area occupied by the cortex (G) is becoming electron-lucent (i.e. does not stain) and may show that DPA is being laid down. The structures (J) inside the spore membrane are not understood and are seen only rarely. (f) The outer spore membrane (H) is now developing and shows a layered structure. The first indication of the inner layer (K) can be seen. (g) The outer spore membrane is now well developed and shows at least four layers. It is very electron-dense. The spore membrane (E) is now very indistinct and the area both inside it and immediately outside it poorly stained. (h) The spore is now complete but is still within the cell wall (L). Both the outer spore membrane (H) and the inner layer (K) have developed the corrugated appearance characteristic of the complete spore. The spore contents are ill-defined and do not take up electrons stains. The bars indicate 1μ . Inset (h) bar indicates 0.1μ .

cut. As it is known that no more than a single spore ever forms in one cell it must be assumed that not all the spore septa shown in this series will develop into mature spores. Nevertheless the five septa appear to be developing along the same lines and there is nothing to suggest which of them might eventually have become mature. The cells and septa are referred to in order from the left. Septa 1 and 2 are in cell 1, septum 3 is in cell 2 and septa 4 and 5 are in cell 3. As the width of the sectioned cells is about 6000 Å and it takes 12 sections to traverse a cell, each section must be about 500 Å thick.

All five septa have mesosomes lying near them (cf. Ellar, Lundgren & Slepecky, 1967), but examination of the appropriate section shows that the mesosomes are attached to the septa. The first septum (cell 1) is in a very early stage of development and consists merely of an elongation of the mesosomal membranes (Plate 4*g*–4*j*), which form an invagination of the cell semipermeable membrane. The second septum (cell 1) is complete across the cell and is bulging slightly towards the centre. The associated mesosome is attached to the cytoplasmic membrane of the cell at its pole (Plate 3*c* and 3*d*). The third septum (cell 2) is in an advanced stage of development and has formed a large bulge. The associated mesosome lies outside the developing spore and its point of attachment can be seen (Plate 3*c* and 3*d*). The fourth septum (cell 3) might be in a similar developmental stage to the first, but, as the point of contact of the mesosome with the cytoplasmic membrane was not visible in the available sections, this cannot be known with certainty. There are slight inward protrusions of the cytoplasmic membrane on opposite edges of the cell near the mesosome (Plate 4*h*–4*k*) and these might be involved in the initiation of formation of the spore septum. The fifth septum (cell 3) is formed completely across the cell and its associated mesosome, which appears to be an invagination of the septal membrane, is seen to be attached to it (Plate 4*j*).

The bacterial nucleoids are visible in cell 1 (Plate 3, *b*–*f*), cell 2 (Plates 3*d*–*f*, 4*g* and 4*h*) and cell 3 (Plates 3*f* and 4*g*–4*j*). The same Plates show that they are associated with mesosomes in cell 1 (Plate 3*d*–3*f*) and in cell 3 (Plate 4*h*–4*k*), but in cell 2 there is no apparent connexion. The mesosomes connected with the developing spore septa are themselves sometimes associated with parts of the bacterial nucleoid that have been severed from the main mass and have migrated into the polar parts of the cells. This can be seen in cell 1 (Plates 3*b*–3*f* and 4*g*) for the mesosome within the developing spore and in cell 3 (Plate 3*c*–3*e*). In cell 2 part of the nucleoid is visible inside the spore (Plates 3*c*–3*f*, 4*g* and 4*h*).

After the formation of the spore septum across the cell near the pole, during which a piece of the nucleoid either with or without a mesosome is separated from the main mass, the septum (B) bulges towards the centre of the cell as shown in Plate 1(*c*). The membrane continues to extend, and its original points of contact with the cytoplasmic membrane appear to approach one another until the ellipsoidal structure, known as the fore-spore, becomes detached within the cell. This stage (stage III) is shown in Plate 1(*d*). The mode of formation of the spore membrane and its continuity with the cytoplasmic membrane during growth suggest that both possess a common chemical composition and physical structure. The fore-spore enclosed within the complete spore membrane (E) contains a mesosome (D) attached to the membrane, and also some ribosomes. The enclosed portion of the bacterial nucleoid, presumably one chromosome, cannot be identified at this stage as it appears to have become dispersed throughout the spore. It is evident from the way in which the mesosome (D1) is pressed against the end of the cell that, outside the spore membrane, there is already a layer of material that is as yet indistinguishable from the cell cytoplasm. This probably represents the beginning of the cortex.

The next stage (stage IV), that of cortex formation, is shown in Plate 2(*e*). The cortex (G) lies outside the spore membrane, but within the thick multilayered spore coat that eventually forms. Inside the spore membrane no detail of nuclear or ribosomal material is now visible, but sometimes small vesicles (J) can be seen just within the membrane.

Outside the cortex the first signs of the spore coat (H) are evident as a thin double membrane with a granular layer inside it. A cell mesosome has become pressed against the cytoplasmic membrane by the developing spore. At this stage (Plate 2*f*) a decrease in the electron density of the spore occurs particularly in the cortex, which develops electron-lucent areas. Eventually the major part of the spore cortex and core becomes electron-lucent (Plate 2*g* and 2*h*). This is observed regularly and does not seem to be an artifact of shrinkage. It is believed to be related to a definite change in the chemical composition of the spore, possibly the deposition of DPA (see Table 1), which begins at $t_{6.5}$. If the development of the electron-lucent areas were due to the deposition of DPA, then they should not be observed earlier than sample no. 5 ($t_{6.5}$). This is found to be the case; no spore structures showing light areas were found except in specimens taken at t_8 and $t_{9.5}$.

In Plate 2*f* the spore coat (H) is developing as a coarse multilayered structure, which is particularly electron-dense (stage V), while just inside it a

finer inner coat (K) is appearing. These two structures are again evident in Plate 2(g), but now the spore coat is very electron-dense and consists of four dark layers separated by less dense layers and surrounded by a thinner dark line. The spore membrane (E) is still visible.

The penultimate stage (stage VI) of spore development is shown in Plate 2(h), where the spore outer coat (H) and the inner coat (K) have assumed the typical corrugated appearance of the mature spore. The corrugations appear to be formed as a result of the deposition of material at the periphery of the cortex in the form of ridges. The cortex (the light area) remains roughly ellipsoidal in shape and only the layers external to it assume the corrugations. The surface appearance of the spores from several members of the genus *Bacillus* have been studied by the replica method and have been shown to bear a pattern of ridges that is species-specific (Bradley & Williams, 1957). It is therefore considered that the corrugations are unlikely to be artifacts, but rather that they represent the ridges in cross-sections. The spore is still contained within the wall (L) of the mother cell.

It is clear from Plate 2(h) inset, not only that the spore coat is a multilayered structure, but that two different sets of layers are present, the fine inner coat (K) and the coarser outer coat (H) (see also Murrell, 1967). The centre-to-centre spacing of the individual lamellae is 120 Å for H and 40 Å for K.

Since the cultures from which the samples were taken were not synchronous, no precise time-scale can be attached to the series of cells shown in Plates 1 and 2. An estimate of the length of each stage might be made from the frequency with which cells in each condition were found, but sufficient cells to make this estimate were not examined.

DISCUSSION

The formation of spores in bacilli is accompanied by marked changes in the chemical composition, enzymic content and morphology of the cell. It is important to relate the first two to the structural changes that accompany them, and it has been part of the purpose of this series of papers to attempt this. Ideally the studies should be made on synchronous cultures where all cells enter spore formation at the same time, as suggested by Murrell (1967). This was not achieved in the experiments described here and therefore any precise time relationships between chemical and morphological alterations to the cells cannot be made. However, certain general observations can be made with confidence.

As shown by Fig. 3 in the preceding paper (Warren, 1968), by the time the initial sample was taken (t_2) the culture had developed its full complement of alkaline phosphatase, net protein synthesis

had almost stopped and synthesis of aconitase had begun. At this stage no morphological changes could be seen. In the second and third samples ($t_{3.5}$ and t_5) many cells had formed spore septa and the first complete spore membranes had appeared. At this time the aconitase activity was still increasing and glucose dehydrogenase had made its appearance. The connexion between these enzymes and the formation of the spore membrane, if any, is unknown. It is clear that the immature spore within its membrane, which is continuous with the bacterial membrane and therefore probably has the same composition, still lacks DPA and is neither refractile nor heat-resistant.

The stages of cortex development and its envelopment in at least two varieties of multi-layered structures (the outer and inner coats) appear to follow rapidly, while at the same time the spore core and cortex lose electron density. It is at this time, between t_5 and $t_{6.5}$, that the synthesis of DPA begins and the first refractile spores can be seen.

The information obtained from this study was combined with that obtained by Warren (1968) to produce the composite picture shown in Fig. 1.

Whether any component of the bacterial cell is dense or not in the electron microscope probably depends on the formation of covalent-linked osmium compounds produced by the interaction of the osmium tetroxide in the fixative with a variety of compounds present in the cell. A well-studied example is that of osmium tetroxide with methyl oleate (Korn & Weisman, 1966), in which bis(methyl 9,10-dihydroxystearate) osmate is formed. From the high electron density of the spore coat it would be expected that it would contain a high proportion of material reactive with osmium tetroxide. However, examination of the available analyses of the spore coat (Murrell, 1967) shows that very little lipid (2.84% of dry wt.) is present. No information on the composition of the lipid is available.

The developing spores become refractile at about t_6 . This change is presumably caused by the formation of a structure that has a higher refractive index than the cytoplasm of the cell and the surrounding medium. It coincides with the spore coat becoming highly electron-dense and is accompanied by the deposition of DPA, but the precise cause of the onset of refractility is unknown. Wise, Swanson & Halvorson (1967) have shown that mutants of *B. cereus* that are unable to synthesize DPA nevertheless form refractile spores, which, however, are not heat-resistant.

A number of hypotheses have been put forward to account for the heat-resistance of spores, including a protecting effect of DPA on spore proteins (Mishiro & Ochi, 1966), an association of spore enzymes with spore structures (Stewart &

EXPLANATION OF PLATES 3 AND 4

Serial sections (*a-k*) through cells of *B. subtilis* in an early stage of spore formation. These show a set of 11 serial sections passing through three cells of *B. subtilis*, in which five presumptive spores are developing. Counting from the left the first membrane in the first cell is at the very early stage of mesosome attachment to the cytoplasmic membrane (*g*) close to the end of the cell. The second spore membrane (in the first cell) is complete across the cell. Some DNA has been trapped in the developing spore and the mesosome is attached to the cytoplasmic membrane at the end of the cell (*c* and *d*). The third spore membrane (in the second cell) is well developed and is swelling out towards the middle of the cell. Some DNA has been trapped in the spore (*c-h*), but the mesosome is outside the spore. The fourth spore membrane (in the third cell) is present only as two opposing bulges in the cytoplasmic membrane (*g-k*). A mesosome (*i-k*) is probably associated with the bulges. The fifth spore membrane (in the third cell) is formed right across the cell. The mesosome, seen associated with some DNA, is attached to the spore membrane and lies within the spore (*j*). The bar indicates 1μ .

Fig. 1. Morphological and biochemical events during sporulation of *B. subtilis*. The data of Warren (1968) were used to construct a chart showing the points of rapid synthesis of enzymes and the appearance of refractility, DPA and heat-resistant spores. The morphological development of the spore during this time is indicated schematically at the top of the Figure.

Halvorson, 1954) and an intrinsic resistance of some spore proteins (see, e.g., Baillie & Norris, 1963) and, at suitable salt concentrations, of glucose dehydrogenase (Bach & Sadoff, 1962). An alternative suggestion is that the spore core is in a highly dehydrated state and that this increases the stability of proteins and nucleic acids to heat and radiation (Donnellan & Setlow, 1965; Smith & Yoshikawa, 1966). No satisfactory mechanism has so far been proposed to explain how this dehydrated state is achieved. It seemed to us that the formation of the calcium salt of DPA (see Powell, 1953) might lead to dehydration of the spore if it were deposited, as we have suggested, as a semi-solid matrix between the rigid inner spore coat and the cytoplasmic membrane. In this way the membrane might be compressed, as is in fact indicated by its diffuse outline in the mature spore (Plate 2*h*) and by the decrease in the volume that it encloses as the electron-lucent areas in the cortex increase in size (Plate 2*f* and 2*g*). Water, but probably not the high-molecular-weight components of the spore core, diffuse out through the layer of DPA and into the exterior of the spore, which appears to be freely permeable to water (Black & Gerhardt, 1962; Neihof, Thompson & Deitz, 1967).

A composite picture of the biochemistry and morphology of the developing spore can now be constructed as shown in Fig. 1.

Many aspects of spore formation remain vague, in particular the events leading to the condensation of the nucleoids into an axial filament, the attachment of the mesosomes to the cytoplasmic membrane near the poles and the exact location of DPA. It is also not clear to what extent the morphological changes are necessarily correlated with, or dependent on, the concurrent biochemical changes (see Fig. 1). It is possible that some indication of the interdependence might be obtained by carrying out further studies of this kind on mutants blocked at various stages in sporulation.

We acknowledge the skilled assistance of Mrs Wendy Brownsill for preparing the sections and thank Mrs Edith Goodman for expert technical assistance. This work was supported by a grant from the Science Research Council.

REFERENCES

- Bach, J. A. & Sadoff, H. L. (1962). *J. Bact.* **83**, 699.
- Baillie, A. & Norris, J. R. (1963). *J. appl. Bact.* **26**, 102.
- Black, S. H. & Gerhardt, P. (1962). *J. Bact.* **83**, 301.
- Bradley, D. E. & Williams, D. J. (1957). *J. gen. Microbiol.* **17**, 75.

- Donnellan, J. E., Nags, E. H. & Levinson, H. S. (1964). *J. Bact.* **87**, 332.
- Donnellan, J. E. & Setlow, R. B. (1965). *Science*, **149**, 308.
- Ellar, D., Lundgren, D. G. & Slepecky, R. A. (1967). *J. Bact.* **94**, 1189.
- Kellenberger, E., Ryter, A. & Séchaud, J. (1958). *J. biophys. biochem. Cytol.* **4**, 671.
- Korn, E. D. & Weisman, R. A. (1966). *Biochim. biophys. Acta*, **116**, 309.
- Mishiro, Y. & Ochi, M. (1966). *Nature, Lond.*, **211**, 1190.
- Murrell, W. G. (1967). *Advanc. microb. Physiol.* **1**, 133.
- Neihof, R., Thompson, J. K. & Deitz, V. R. (1967). *Nature, Lond.*, **216**, 1304.
- Powell, J. F. (1953). *Biochem. J.* **54**, 210.
- Reynolds, E. S. (1963). *J. Cell. Biol.* **17**, 208.
- Ryter, A. (1965). *Ann. Inst. Pasteur*, **108**, 40.
- Schaeffer, P., Ionesco, H., Ryter, A. & Balassa, G. (1965). *Colloq. int. Cent. nat. Rech. sci., Marseille*, p. 553.
- Smith, K. C. & Yoshikawa, H. (1966). *Photochem. Photobiol.* **5**, 777.
- Stewart, B. T. & Halvorson, H. O. (1954). *Arch. Biochem. Biophys.* **49**, 168.
- Warren, S. C. (1968). *Biochem. J.* **109**, 811.
- Wise, J., Swanson, A. & Halvorson, H. O. (1967). *J. Bact.* **94**, 2075.
- Young, E. I. & Fitz-James, P. C. (1959). *J. biophys. biochem. Cytol.* **6**, 467.