is concluded (a) that activation by NaCl can almost match the activation by tris and (b) that the effect is not on k_3 but on k_2 .

Fernley, H. N. & Walker, P. G. (1966). *Nature, Lond.*, **212**, 1435.

Heppel, L. A., Harkness, D. R. & Hilmo, R. J. (1962). *J. biol. Chem.* **237**, 841.

Trentham, D. R. & Gutfreund, H. (1968). *Biochem. J.* **106**, 455.

Wilson, I. B., Dayan, J. & Cyr, K. (1964). *J. biol. Chem.* **239**, 4182.

Shay Chloroma Alkaline Phosphatase

By J. DUNNE, J. J. FENNELLY, K. F. MCGEENEY.
(Department of Medicine and Therapeutics, University College, Dublin, Irish Republic)

The characterization of normal human and animal tissue alkaline phosphatase (orthophosphoric monoester phosphohydrolase, EC 3.1.3.1) has been described (Morton, 1955) and studied in great detail (Moss, Eaton, Smith & Whitby, 1967; Sussman, Small & Cotlove, 1968). Since such studies have not to date been reported for a tumour enzyme we studied the Shay Chloroma, a myeloblastic tumour in the rat (Moloney, Dorr, Dowd & Boschetti, 1962). This tumour contains an average of 8.0 i.u. of alkaline phosphatase/g. wet wt. of tumour. The enzyme was purified by butanol extraction, acetone fractionation, ammonium sulphate fractionation, gel filtration on Sephadex G-200 and chromatography on DEAE-cellulose DE52.

Enzyme activity was eluted in 3 peaks from DEAE-cellulose DE52 at 50, 100 and 250 mM-NaCl, designated respectively peaks A, B and C. The purification achieved was 200-fold for peak B and 90-fold and 120-fold for peaks A and C. Enzyme B was eluted in two peaks of activity from Sephadex G-200, an included and excluded peak, and enzymes A and C were both totally excluded from the gel.

The K_m values for the three enzymes with *p*-nitrophenyl phosphate as substrate were 0.46, 0.37 and 0.44 mM and the pH optimum was pH 10.2 in each case. Enzyme A was the most resistant to urea inactivation, 55% of the activity remaining in 3.0 M-urea compared with 20% with enzymes B and C. The urea inhibition was non-competitive. On polyacrylamide-gel electrophoresis, enzyme B moved as a single band and had the highest mobility, enzyme A separated into three bands and enzyme C into two bands of activity. One component of enzyme A and of enzyme B was retarded after neuraminidase treatment. Extensively dialysed samples of the three enzymes were activated by Mg^{2+} , all to different degrees. EDTA inactivated enzymes A, B and C, the inactivation being a function of time and concentration; the activity was only partially restored by addition of Mg^{2+} .

The three enzymes varied in their heat stability but were all inhibited to the same extent (20%) by L-phenylalanine. The ratio of alkaline phosphatase activity to inorganic pyrophosphatase (pyrophosphate phosphohydrolase, EC 3.6.1.1) activity was not constant through all the steps of purification (Moss *et al.* 1967); the ratios for enzymes A, B and C were 13, 20 and 12 respectively.

Moloney, W. C., Dorr, A. D., Dowd, G. & Boschetti, A. E. (1962). *Blood*, **19**, 45.

Morton, R. K. (1955). *Biochem. J.* **61**, 232.

Moss, D. W., Eaton, R. H., Smith, J. K. & Whitby, L. G. (1967). *Biochem. J.* **105**, 32.

Sussman, H. H., Small, P. A., jun. & Cotlove, E. (1968). *J. biol. Chem.* **243**, 160.

The Human Pseudocholinesterase System

By PATRICK J. GAFFNEY, jun. (Medical Research Council Abnormal Haemoglobin Unit, Department of Biochemistry, University of Cambridge)

Multiple forms of human pseudocholinesterase have been demonstrated on starch-gel (La Motta, McComb, Noll, Wetstone & Reinfrank, 1968) and polyacrylamide-gel (Juul, 1968) electrophoresis. It has been shown (Svensmark, 1961) that there are several molecules of *N*-acetylneuraminic acid (NANA) associated with each molecule of pseudocholinesterase, but that these play no part in the enzyme function (Augustinsson & Ekedahl, 1962).

A 460-times enriched preparation of the enzyme was obtained from pooled outdated plasma by ammonium sulphate precipitation and DEAE-cellulose fractionation at pH 4.2. Polyacrylamide-gel electrophoresis at pH 8.9 followed by enzyme detection with butyrylthiocholine iodide showed that some enzyme components of whole serum were lost during this purification process. The consistency of the distribution of the enzyme components of whole serum on polyacrylamide gels under various conditions of gel-pore size and buffering indicated that the heterogeneity may not be completely explained by assuming polymer forms for the enzyme (La Motta *et al.* 1968).

The enriched enzyme preparation, after treatment with neuraminidase (EC 3.2.1.18) for 7 days at 30° was quite homogeneous on polyacrylamide-gel electrophoresis and its mobility was less than that of the major component of the heterogeneous (6–10 components) control. Both the neuraminidase-treated preparation and its control yielded three fractions on elution from Sephadex G-75 with dilute ammonia solution at pH 8.2. The two major protein fractions were poorly resolved and eluted at about the void volume and the third minor fraction was eluted at about twice the void volume. Whereas enzyme activity was associated