startling picture of these variations. The earlier observation that κ light chains contain two disulphide bridges (Milstein, 1964), each including about 60 residues (Hiltschmann & Craig, 1965), encouraged the study of the intrachain bridges in other chains. The results so far support the idea that the 60-residues loop is the basic building block of all immunoglobulin chains.

Studies of the four subtypes of heavy chains ($\gamma 1$, $\gamma 2$, $\gamma 3$ and $\gamma 4$) have shown that three disulphide-bridged peptides are characteristically present in all cases (with occasional differences specific of each type). A fourth, variable, bridge has only been obtained in a few cases. It appears therefore that the heavy chains include a common structure, invariable within each subclass, of three loops: two in the C-terminal half (Fc fragment) and one in the N-terminal half (Fd fragment). This result supports observations on the attachment of the light chains to the heavy chains (Pink & Milstein, 1967a) indicating invariant sections of human Fd fragments. Although evidence that each loop contains about 60 residues is only available in a few cases, the similarity of the bridged peptides suggest that this is so in all of them.

In contrast to this close similarity of intrachain bridges between the different types of γ chains, the number, arrangement and distribution of interchain bridges is remarkably variable. Not less than two [in $\gamma 1$ (Steiner & Porter, 1967; Frangione & Milstein, 1967) and $\gamma 4$ (Pink & Milstein, 1967b)] and up to five [(in $\gamma 3$ (Frangione & Milstein, 1968)] inter-heavy-chain bridges held the two halves of the molecule together. All these isotypic variants are thought to arise from different structural genes coding for the invariant part of the molecule. The C-terminal half of κ chains was found to be identical in two different proteins (Hiltschmann & Craig, 1965) and in all the proteins studied later, with the single exception of residue 191, which was established to be correlated with a genetic (Inv) marker (Milstein, 1966; Baglioni, Alescio Zonta, Cioli & Carbonara, 1966). This and the marker in $\gamma 1$ chains are the only two human allotypic markers that have so far been correlated with sequence variation. They indicate that this part of the molecule is coded by a single gene.

Extensive information concerning the idiotypic variants of κ chains is being collected by several Laboratories. The combined results suggest the presence of a number of basic sequences (not less than three according to Milstein, 1967). Results

on light chains obtained from normal individuals suggest that these basic sequences are not allelic, and in all probability represent an equivalent number of genes involved in the control of the N-terminal half of the molecule. However, in this part of the molecule additional variability (possibly of somatic origin) has been observed. This variability is extended in such a way that so far no two κ chains have been found to be identical.

- Baglioni, C., Alescio Zonta, L., Cioli, D. & Carbonara, A. (1966). *Science*, **152**, 1519.
 Cohen, S. & Milstein, C. (1967). *Advanc. Immunol.* **7**, 1.
 Frangione, B. & Milstein, C. (1967). *Nature, Lond.*, **216**, 939.
 Frangione, B. & Milstein, C. (1968). *J. molec. Biol.* **33**, 893.
 Hiltschmann, N. & Craig, L. C. (1965). *Proc. nat. Acad. Sci., Wash.*, **53**, 1403.
 Milstein, C. (1964). *J. molec. Biol.* **9**, 836.
 Milstein, C. (1966). *Nature, Lond.*, **209**, 370.
 Milstein, C. (1967). *Nature, Lond.*, **260**, 330.
 Pink, J. R. L. & Milstein, C. (1967a). *Nature, Lond.*, **214**, 92.
 Pink, J. R. L. & Milstein, C. (1967b). *Nature, Lond.*, **214**, 941.
 Steiner, L. A. & Porter, R. R. (1967). *Biochemistry*, **6**, 3957.

Haemoglobin

By E. R. HUEHNS. (*Medical Research Council Group on Haemolytic Anaemia, University College Hospital Medical School, London, W.C. 1.*)

Among the large number of human haemoglobin variants so far identified, a few cause disease because the abnormal protein is either unstable or has abnormal oxygen-dissociation properties. The exact location in the three-dimensional structure of the molecule of the amino acid differences causing these specific abnormalities can be determined from the atomic model of haemoglobin built by Perutz and his co-workers (Perutz, 1965; Perutz, Muirhead, Cox & Goaman, 1968). The results indicate that certain regions of the molecule, such as the pockets containing the haem groups and the two contacts between the α - and β -chains, are of particular importance in maintaining the normal function and stability of haemoglobin. Several such amino acid substitutions will be discussed. The findings, in general, are in keeping with the small degree of variation of these residues among normal mammalian haemoglobins, and confirm the importance of these parts of the molecule.

- Perutz, M. F. (1965). *J. molec. Biol.* **13**, 646.
 Perutz, M. F., Muirhead, H., Cox, J. M. & Goaman, L. C. G. (1968). *Nature, Lond.*, **219**, 131.