and to compare this with other avian ovalbumins. Cysteine-containing regions of the ovalbumin polypeptide chain were located by labelling with iodo[^{14}C]acetate after mercaptoethanol reduction, followed by tryptic digestion and high-voltage electrophoresis. It was found that one of the hen ovalbumin cysteine peptides was the C-terminal peptide and that the cysteine residue was the fourth amino acid from the C-terminus (Fothergill & Gilmore, 1968). All the labelled peptides from duck ovalbumin contained lysine or arginine. The involvement of this C-terminal peptide in the disulphide bond of hen ovalbumin was investigated by using a double-labelling technique.

Hen ovalbumin was carboxymethylated with iodo[^{14}C]acetate in 8M-urea (Crestfield, Moore & Stein, 1963) but without the addition of reducing agent. After removal of the unchanged iodo[^{14}C]acetate by gel filtration, the carboxymethylated ovalbumin was treated with mercaptoethanol and iodo[^3H]acetate. The doubly labelled protein, containing approximately 4g.atoms of ^{14}C and 2g.atoms of ^3H /mole, was heat-denatured and digested with trypsin. The tryptic peptides were separated by high-voltage electrophoresis at pH 6.5 on Whatman no. 3 MM paper. The electrophoresis paper was cut transversely at 1cm. intervals and ^{14}C and ^3H were measured by using a liquid-scintillator double-counting technique. ^3H was found in the C-terminal peptide and in the pH 6.5 neutral region, whereas only a small amount of ^{14}C was located in the C-terminal peptide.

We conclude that the cysteine residue close to the carboxyl terminus of hen ovalbumin is involved in the disulphide bond, and that duck ovalbumin probably lacks this cysteine residue. We are investigating by similar techniques other ovalbumins from birds that are more closely related phylogenetically to the hen.

Crestfield, A. M., Moore, S. & Stein, W. H. (1963). *J. biol. Chem.* **238**, 622.

Fothergill, J. E. & Gilmore, L. A. (1968). *Abstr. 5th Meet. Fed. Europ. biochem. Soc., Prague*, p. 36.

Fothergill, J. E. & Perrie, W. T. (1966). *Biochem. J.* **99**, 58P.

Gilmore, L. A. & Fothergill, J. E. (1967). *Biochem. J.* **103**, 39P.

Binding of Glutathione and Cysteine to Cellular Protein by Mixed Disulphide Bonds

By R. C. JACKSON, K. R. HARRAP and C. A. SMITH.
(Chester Beatty Research Institute, Institute of Cancer Research: Royal Cancer Hospital, Fulham Road, London, S.W. 3)

It has been demonstrated by Modig (1968) that protein-bound GSH and cysteine are present in

Ehrlich ascites cells in amounts that form about one-third of the total cellular content of these thiols. The present study investigated the distribution of mixed disulphides in normal tissues, and their mechanism of formation. We attempted also to assess their involvement in the turnover of cellular free thiol.

High-speed-supernatant fractions were obtained from rat spleen, heart, liver, skeletal muscle, kidney and Yoshida ascites sarcoma. The preparations were dialysed, to remove free thiol and disulphide, and analysed for mixed disulphide by a modification of the method of Modig (1968). Extracts were reduced with sodium borohydride in borate buffer, pH 9.0, containing 8M-urea. After deproteinization with sulphosalicylic acid, GSH was estimated by the method of Ball (1966), and total thiol by the method of Ellman, as modified by Harrap (1967). Mixed disulphides were found in all the tissues examined. Concentrations of the protein-GSH mixed disulphide in the different tissues were similar, but amounts of the protein-cysteine disulphide were more variable.

To examine the ability of tissues to reduce mixed disulphide, substrates were prepared from bovine plasma albumin with GSH or cysteine. Homogenates of Yoshida ascites cells, and of kidney, spleen and heart were found to reduce the albumin-GSH disulphide in the presence of NADPH and to a much lesser extent with NADH. During this enzymic reaction, GSH and protein thiol groups were released in equimolar ratio. Little or no such activity was found in erythrocytes, skeletal muscle or liver. None of the tissues examined would reduce the albumin-cysteine disulphide under these conditions.

An attempt was made to test the speculation that these compounds might act as intracellular reserves of thiol compounds by using Yoshida ascites sarcoma cells in which a sixfold increase of cellular GSH content had been induced by chlorambucil treatment (Harrap & Hill, 1968). Under these conditions, far from decreasing as the free thiol concentration increased, both protein-GSH and protein-cysteine mixed disulphides doubled in concentration.

Ball, C. R. (1966). *Biochem. Pharmacol.* **15**, 809.

Harrap, K. R. (1967). *Biochem. Pharmacol.* **16**, 725.

Harrap, K. R. & Hill, B. T. (1968). *Int. J. Cancer* (In the Press).

Modig, H. (1968). *Biochem. Pharmacol.* **17**, 177.