PROCEEDINGS OF THE BIOCHEMICAL SOCIETY

The 488th Meeting of the Society was held at the Institute of Psychiatry, De Crespigny Park, London, S.E. 5, on Thursday and Friday, 21 and 22 November 1968, when the following papers were presented:

COLLOQUIUM ON 'INHERITED ABNORMALITIES AFFECTING THE NERVOUS SYSTEM'

Homocystinuria

D. C. CUSWORTH and C. E. DENT. (*Medical Unit, University College Hospital Medical School, London, W.C. 1*)

Homocystinuria was first detected in two sisters during the screening of mentally retarded children in Northern Ireland for abnormal amino acid excretion (Carson & Neill, 1962). At the same time, homocystine was independently identified in the urine of a retarded child in the U.S.A. (Gerritsen, Vaughn & Waisman, 1962). In addition to mental retardation and motor seizures, the two sisters had dislocated lenses, knock knees, fine friable hair and erythematous blotches on the skin, particularly evident on the cheeks and limb extremities. Further cases with similar features were subsequently found in Northern Ireland and England, and in addition skeletal disorders and thrombo-embolic disease were discovered, several patients having died from pulmonary embolus or renal-artery thrombosis (Carson, Dent, Field & Gaull, 1965). A survey of a large number of patients with non-traumatic dislocated lenses in the U.S.A. showed that about 5% were in fact cases of homocystinuria, and altogether 38 new cases were found. Many of the features of the European cases were found, but a surprising new observation was that 16 of the new cases were of normal intelligence (Schimke, McKusick, Huang & Pollack, 1965). Although many cases have occurred in siblings, parents and offspring of cases have not been affected; both sexes are affected, there is a relatively high proportion of consanguineous marriages and in one family cases have occurred in cousins, all factors which suggest that homocystinuria is inherited as an autosomal recessive character. New cases are being reported with increasing frequency and there are now probably over a hundred in the literature, 6 years after the original observation; next to phenylketonuria, this is the commonest inherited biochemical disorder associated with mental retardation.

The simplest explanation of the homocystine excretion was to postulate a block in the metabolism of methionine between homocysteine and cys-

thionine, but the measured urine homocystine excretions of 50–100mg. per day were very much less than expected if the major part of the 1–2g. dietary methionine intake was being excreted in this form. Homocystine is not found in the urine or plasma of normal subjects, and was not at first detected in the plasma of homocystinuric patients owing to binding to protein during storage, but was later found to be at a concentration of 0.03–0.07 μ mole/ml.; the plasma methionine also was sometimes elevated (Carson *et al.* 1963). The possibility of a methionine-transport defect was considered, but oral and intravenous loads of methionine produced similar plasma methionine concentrations in homocystinuric patients. These tolerance tests showed that whereas the plasma methionine had returned to the pre-load concentration 24 hr. later in normal subjects, the concentration in patients with homocystinuria was still elevated at this time and indeed remained elevated for several days. Homocystine concentrations showed no consistent changes after these methionine loads (Brenton, Cusworth & Gaull, 1965a).

Confirmation of the postulated enzyme defect was elegantly demonstrated *in vitro* by Mudd, Finkelstein, Irreverre & Laster (1964) who measured cystathionine synthetase activity in liver biopsies in a series of patients and found no activity in a case of homocystinuria. Additional evidence was also provided for a recessive mode of inheritance, for the hepatic cystathionine synthetase activity of the parents of this patient was much lower than in normal subjects (Finkelstein, Mudd, Irreverre & Laster, 1964). Further evidence for an enzyme block has been obtained by studies *in vivo* in homocystinuric patients. For example, dietary cystine is essential for the maintenance of positive nitrogen balance in affected subjects (Brenton, Cusworth, Dent & Jones, 1966), and after a tracer dose of [³⁵S]methionine the excretion of labelled inorganic sulphate and the incorporation of labelled cystine into plasma proteins was very much lower than in normal subjects (Brenton & Cusworth, 1966). The urinary excretion of inorganic sulphate after large and repeated doses of methionine is much lower than normal, but there is no difference in sulphate

excretion after cysteine administration (Laster, Mudd, Finkelstein & Irreverre, 1965). A further consequence of the decreased activity of cystathionine synthetase is the absence of cystathionine from the brain, where it is normally present in considerable amount (Brenton, Cusworth & Gaull, 1965b; Gerritsen & Waisman, 1964). This abnormality may have some relation to the mental retardation associated with homocystinuria, but it is not yet known if those patients with normal intelligence have normal amounts of cystathionine in the brain.

There are a number of connective tissue defects, the most constant finding being the dislocation of the lenses, probably due to degenerative changes in the zonular fibres and ciliary body which have been demonstrated by light and electron microscopy (Henkind & Ashton, 1965). Skeletal changes include scoliosis, arachnodactyly, and enlargement and shape changes of vertebral bodies, tibiae and femora (Brenton *et al.* 1966). Structural changes in the elastic fibres of intimal and medial layers of arterial walls are present (Gibson, Carson & Neill, 1964) and these may be partly responsible for the thrombus formation which often occurs. An increase in soluble collagen and a reduction in collagen cross-linking has been demonstrated in the skin of homocystinuric patients (Harris & Sjoerdsma, 1966). We have suggested that homocystine may interfere with protein disulphide cross-linkages (Brenton & Cusworth, 1966) and although this cannot be of importance with collagen, which contains virtually no sulphur amino acid, it may occur with other structural proteins. Aorta from a patient who died from pulmonary emboli showed a marked decrease in glycoprotein and protein-polysaccharide matrix, and it is possible that altered organic sulphate metabolism may be responsible for this (M. J. Smith & S. M. Partridge, quoted by Carson *et al.* 1965).

Treatment by restriction of dietary methionine intake has decreased homocystine and methionine concentrations in body fluids (Komrower, Lambert, Cusworth & Westall, 1966; Perry, 1967). The addition of choline to the diet (Perry, Hansen, Love, Crawford & Tischler, 1968) or the administration of folic acid (Carey, Fennelly & Fitzgerald, 1968) lowers homocystine concentrations, presumably by promoting remethylation to methionine. Large doses of pyridoxine lead to decreased concentrations of methionine and homocystine in some patients (Barber & Spaeth, 1967) but not in others (Schimke, McKusick & Weilbaecher, 1967). We have recently shown that arginine can inhibit the renal tubular reabsorption of homocystine and lead to a large increase in its excretion in the urine, offering yet another therapeutic approach (D. C. Cusworth & A. Gattereau, 1968). The clinical value of these various therapeutic attempts will take many years

to assess as the natural course of the disease is slow and extremely variable. There are many unsolved problems in this intriguing disease. The clinical effects are so variable; in particular some patients are profoundly mentally retarded while others are of normal intelligence. The life-endangering thrombosis may be due only to structural changes in the blood-vessel walls, but there may also be a biochemical cause. The basis of the connective tissue changes is not known. The discrepancy between methionine intake and homocystine excretion and the relative constancy of plasma homocystine concentrations are difficult to explain solely on the basis of an enzymic block between homocystine and cystathionine.

- Barber, G. W. & Spaeth, G. L. (1967). *Lancet*, i, 337.
 Brenton, D. P. & Cusworth, D. C. (1966). *Clin. Sci.* **31**, 197.
 Brenton, D. P., Cusworth, D. C., Dent, C. E. & Jones, E. E. (1966). *Quart. J. Med.* **35**, 325.
 Brenton, D. P., Cusworth, D. C. & Gaull, G. E. (1965a). *J. Pediat.* **67**, 58.
 Brenton, D. P., Cusworth, D. C. & Gaull, G. E. (1965b). *Pediatrics*, N.Y., **35**, 50.
 Carey, M. C., Fennelly, J. J. & Fitzgerald, O. (1968). *Amer. J. Med.* **45**, 26.
 Carson, N. A. J., Cusworth, D. C., Dent, C. E., Field, C. M. B., Neill, D. W. & Westall, R. G. (1963). *Arch. Dis. Childh.* **38**, 425.
 Carson, N. A. J., Dent, C. E., Field, C. M. B. & Gaull, G. E. (1965). *J. Pediat.* **66**, 565.
 Carson, N. A. J. & Neill, D. W. (1962). *Arch. Dis. Childh.* **37**, 505.
 Cusworth, D. C. & Gattereau, A. (1968), *Lancet*, ii, 916.
 Finkelstein, J. D., Mudd, S. H., Irreverre, F. & Laster, L. (1964). *Science*, **146**, 785.
 Gerritsen, T., Vaughn, J. G. & Waisman, H. A. (1962). *Biochem. biophys. Res. Commun.* **9**, 493.
 Gerritsen, T. & Waisman, H. A. (1964). *Science*, **145**, 588.
 Gibson, J. B., Carson, N. A. J. & Neill, D. W. (1964). *J. clin. Path.* **17**, 427.
 Harris, E. D. & Sjoerdsma, A. (1966). *Lancet*, ii, 707.
 Henkind, P. & Ashton, N. (1965). *Trans. ophthal. Soc.* **85**, 21.
 Komrower, G. M., Lambert, A. M., Cusworth, D. C. & Westall, R. G. (1966). *Arch. Dis. Childh.* **41**, 666.
 Laster, L., Mudd, S. H., Finkelstein, J. D. & Irreverre, F. (1965). *J. clin. Invest.* **44**, 1708.
 Mudd, S. H., Finkelstein, J. D., Irreverre, F. & Laster, L. (1964). *Science*, **143**, 1443.
 Perry, T. L. (1967). In *Amino Acid Metabolism and Genetic Variation*, p. 279. Ed. by Nyhan, W. L. New York: McGraw-Hill Book Co.
 Perry, T. L., Hansen, S., Love, D. L., Crawford, L. E. & Tischler, B. (1968). *Lancet*, ii, 474.
 Schimke, R. N., McKusick, V. A., Huang, T. & Pollack, A. D. (1965). *J. Amer. med. Ass.* **193**, 711.
 Schimke, R. N., McKusick, V. A. & Weilbaecher, R. G. (1967). In *Amino Acid Metabolism and Genetic Variation*, p. 297. Ed. by Nyhan, W. L. New York: McGraw-Hill Book Co.