PROCEEDINGS OF THE BIOCHEMICAL SOCIETY

The 489th Meeting of the Society was held at the Woolwich Polytechnic, Wellington Street, London, S.E. 18, on Thursday, 19 December 1968, when the following papers were presented:

COLLOQUIUM ON 'SUBCELLULAR LOCALIZATION AND FUNCTION OF ACID HYDROLASES'

Localization and Function of Acid Hydrolases in Animal Cells

By P. JACQUES. (*Laboratoire de Chimie Physiologique, Université de Louvain*)

Lysosomal and Non-Lysosomal Localization of Acid Hydrolases in Animal Cells

By VIVIANE MAGGI. (*Tissue and Organ Culture Unit, Imperial Cancer Research Fund, London, W.C. 2*)

Several acid phosphatases with different substrate specificities, different responses to inhibitors and different tissue and subcellular localizations have been described in a number of mammals (Neil & Horner, 1964*a,b*; Maggi, 1965; Maggi, Franks & Carbonell, 1966; Nelson, 1966; Franks, Maggi & Carbonell, 1967), protozoa (Holter & Lowy, 1958-60; Allen, Misch & Morrison, 1963), insects (Bowen, 1968; D. J. Beadle and P. B. Gahan, unpublished work) and plants (Gahan & McLean, 1967; Meany, Gahan & Maggi, 1967). In mouse coagulating glands and seminal vesicles one of these acid phosphatases splits β -glycerophosphate and naphthol AS phosphate monoesters, is completely inhibited by sodium fluoride and sodium molybdate at concentrations of between 1 mM and 0.1 μ M, has a pH optimum between 5 and 6 with both substrates and is mainly confined to the secretory border of the epithelium; lysosomes are also localized in this area. A second type of acid phosphatase is also present. This does not split glycerophosphate but only naphthyl phosphate monoesters, is not inhibited by 0.1 M-sodium fluoride or 0.1 M-sodium molybdate, has a pH optimum of between 5.6 and 5.8, and is sited in the basement membrane region (Maggi *et al.* 1966) where lysosomes are rarely found. Although this latter enzyme is present in most tissues in a number of animal species and plants, its localization within the same tissue may vary according to the species. In the mouse kidney it is present in the region of the brush border of the first

portion of the proximal convoluted tubule but not in the rest of the tubule; in the rat kidney it is present along the whole length of the proximal convoluted tubule but with different intracellular localization in different segments of the same tubule. Lysosomes, on the other hand, are present along the whole length of the proximal convoluted tubules in these species: in the brush-border area in the mouse kidney, and predominantly at the cell base in the rat kidney.

The development of a technique for the ultrastructural cytochemical demonstration of acid phosphatase by using naphthol AS/B1 phosphate as a substrate and lead phthalocyanin diazotate as a coupling agent (Maggi, Livingston, Franks & Coombs, 1968) has made it possible to show that whereas one enzyme is localized in the lysosomes and possesses all the properties of the lysosomal β -glycerophosphate phosphatase described by de Duve (1963), the other is present in some parts of the rough endoplasmic reticulum in specific regions of the cells.

As well as the kidney and the prostatic complex of adult rats and mice, other tissues studied include striated muscle, intestine, liver, testis, pancreas and salivary gland of adult and young mice. Tissues from very young mice (1-3 weeks old) have a very low activity of this second, non-lysosomal, acid phosphatase (Maggi Franks & Carbonell, 1967). Cells in culture (V. Maggi unpublished work), or cells from malignant tumours (Maggi & Franks, 1967; V. Maggi & A. P. Wyatt, unpublished work), or cells in which metaplastic changes have been induced by castration and treatment with oestrogens (V. Maggi and A. W. Steggles, unpublished work) do not contain the enzyme, although they show in lysosomal acid phosphatase.

β -Glycerophosphatase, considered to be an exclusively lysosomal enzyme, has also been found in the endoplasmic reticulum of cells of rat kidney adenomas (Seljelid, 1966), the intestine of the desert locust (Bowen, 1968), rat brain (Kalina & Bubis, 1968) and the kidney of mice after short-term starvation (Maggi & Oddy, 1968). It is

possible that β -glycerophosphatase has a predominantly digestive role, whether auto- or heterophagic, and that the naphthol phosphatase described in this paper has a completely different role in cell metabolism.

The distribution of the naphthol phosphatase in both lysosomes and endoplasmic reticulum is similar to that of a number of other acid hydrolases, e.g. esterases (Holt & Hicks, 1966), β -glucuronidase (Fishman, Goldman & De Lellis, 1967) and arylsulphatase (Dodgson, Rose & Spencer, 1957; Dodgson, Spencer & Thomas, 1954, 1955; Viala & Gianetto, 1955; Milsom, Rose & Dodgson, 1968).

- Allen, S. L., Misch, S. M. & Morrison, B. M. (1963). *J. Histochem. Cytochem.* **11**, 706.
- Bowen, I. D. (1968). *J. R. micr. Soc.* **88**, 279.
- de Duve, C. (1963). In *Ciba Found. Symp.: Lysosomes*, p. 1. Ed. by de Reuck, A. V. S. & Cameron, M. P. London: J. and A. Churchill Ltd.
- Dodgson, K. S., Rose, F. A. & Spencer, B. (1957). *Biochem. J.* **66**, 357.
- Dodgson, K. S., Spencer, B. & Thomas, J. (1954). *Biochem. J.* **56**, 177.
- Dodgson, K. S., Spencer, B. & Thomas, J. (1955). *Biochem. J.* **59**, 29.
- Fishman, W. H., Goldman, S. S. & De Lellis, R. (1967). *Nature, Lond.*, **213**, 457.
- Franks, L. M., Maggi, V. & Carbonell, A. W. (1967). *J. Anat., Lond.*, **101**, 777.
- Gahan, P. B. & McLean, J. (1967). *Biochem. J.* **102**, 47P.
- Holt, S. J. & Hicks, R. M. (1966). *J. Cell Biol.* **29**, 361.
- Holter, H. & Lowy, B. A. (1958-1960). *C.R. Lab. Carlsberg*, **31**, 105.
- Kalina, M. & Bubis, J. J. (1968). *Histochemie*, **14**, 103.
- Maggi, V. (1965). *Biochem. J.* **96**, 12P.
- Maggi, V. & Franks, L. M. (1967). *Biochem. J.* **106**, 21P.
- Maggi, V., Franks, L. M. & Carbonell, A. W. (1966). *Histochemie*, **6**, 305.
- Maggi, V., Franks, L. M. & Carbonell, A. W. (1967). *Biochem. J.* **102**, 48P.
- Maggi, V., Livingston, D. C., Franks, L. M. & Coombs, M. M. (1968). *Proc. R. micr. Soc.* **3**, 79.
- Maggi, V. & Oddy, M. F. (1968). *Histochem. J.* **1**, 78.
- Meany, A., Gahan, P. B. & Maggi, V. (1967). *Histochemie*, **11**, 280.
- Milsom, D. W., Rose, F. A. & Dodgson, K. S. (1968). *Biochem. J.* **109**, 40P.
- Neil, M. W. & Horner, M. W. (1964a). *Biochem. J.* **92**, 217.
- Neil, M. W. & Horner, M. W. (1964b). *Biochem. J.* **93**, 220.
- Nelson, B. D. (1966). *Proc. Soc. exp. Biol., N.Y.*, **121**, 998.
- Seljelid, R. (1966). *J. Ultrastruct. Res.* **16**, 569.
- Viala, R. & Gianetto, R. (1955). *Canad. J. Biochem. Physiol.* **33**, 839.

Vacuoles as Lysosomes of Plant Cells

By PH. MATILE. (*Department of General Botany, Swiss Federal Institute of Technology, Zurich, Switzerland*)

The investigation of plant subcellular particles has led to the suggestion that lysosome-like structures as defined by de Duve are absent from plant cells. If plant tissues are homogenized by using conventional techniques the lysosomal enzymes are largely present in the soluble cell fraction. Nevertheless, with appropriate methods it can be shown that these enzymes are localized in cell compartments corresponding to the lysosomes of animal cells.

In yeast cells they are identical with the vacuoles that are destroyed on conventional homogenization. The isolation of intact vacuoles has been achieved by using naked protoplasts instead of whole cells. Under suitable conditions the lysis of osmotically labilized protoplasts is accompanied by the release of vacuoles, which can be separated by centrifugation in the presence of Ficoll. Fractions of isolated yeast vacuoles contain very high specific activities of the lysosomal enzymes. The analysis of various cell fractions suggests that the vacuoles are the only lysosomes of the yeast cell (Matile & Wiemken, 1967).

Evidence favouring the classification of higher plant vacuoles as lysosomes comes from several lines. On cytochemical grounds the localization of acid phosphatase in meristematic vacuoles of wheat embryos (Poux, 1963a), autophagic vacuoles of starving *Euglena* (Brandes & Bertini, 1964) and aleurone vacuoles (Poux, 1963b, 1965) has been established. However, acid phosphatase seems to represent a doubtful marker enzyme for lysosomal structures. It may be completely absent from lysosomes as it is the case in yeast cells. In addition, its exact function in intracellular digestion has not yet been elucidated, and, finally, in higher-plant cells several acid phosphatases differing with regard to intracellular distribution are present (Ph. Matile, unpublished work). Therefore, the evidence based on isolated organelles seems to be more conclusive.

In meristematic vacuoles isolated from root-tip cells of corn seedlings a dozen acid hydrolases including proteolytic enzymes, nucleases, glucosidases, esterases and phosphatase are localized (Matile, 1966, 1968a; Ph. Matile, unpublished work). It is noteworthy that the largest vacuoles that can be extracted from root-tip cells are not sedimentable in the presence of comparatively low concentrations of sucrose. Thus these lysosomes cannot be prepared if sucrose solutions are used as media for homogenization and centrifugation.

According to the view of classical cytologists the