PRETEST AND OBJECTIVES

To receive your
credit certificate
immediately for free—

Go to
www.PSYCHIATRIST.com
and complete this
activity online.

Articles are selected for CME credit designation on the basis of the CME Institute's assessment of the needs of readers of *The Primary Care Companion*, with the purpose of providing readers with a curriculum of CME articles on a variety of topics throughout each volume. There are no prerequisites for participation in this CME activity.

To obtain credit, please study the designated article and complete the Posttest.

Accreditation Statement

The CME Institute of Physicians Postgraduate Press, Inc. is accredited by the Accreditation Council for Continuing Medical Education to provide continuing medical education for physicians.

Credit Designation

The CME Institute of Physicians Postgraduate Press, Inc. designates this educational activity for up to 1 Category 1 credit toward the American Medical Association Physician's Recognition Award. Each participant should claim only those credits that he/she actually spent in the educational activity.

Date of Original Release/Review

This educational activity is eligible for CME credit through August 31, 2007. The latest review of this material was July 2005.

Educational Objective

After studying the article by Ramaswamy et al., you will be able to:

- Screen for and initiate treatment for posttraumatic stress disorder in the primary care setting.

This pretest is designed to facilitate your study of the material.

1. The usual pharmacologic treatment of posttraumatic stress disorder begins with:

- a. A selective serotonin reuptake inhibitor
- b. A mood stabilizer
- c. A benzodiazepine
- d. An antipsychotic

Pretest answer and Posttest on page 188.

Disclosure of Off-Label Usage

The authors have determined that, to the best of their knowledge, citalopram, clonazepam, escitalopram, fluoxetine, lithium, olanzapine, quetiapine, risperidone, temazepam, zolpidem, alprazolam, carbamazepine, lamotrigine, topiramate, trazodone, and zaleplon are not approved by the U.S. Food and Drug Administration for the treatment of posttraumatic stress disorder.

Answers to Vol. 5, No. 2 Posttest 2003

1. c 2. d 3. a 4. d 5. d

Answers to Vol. 5, No. 3 Posttest 2003

1. d 2. b 3. c 4. a 5. c

A Primary Care Perspective of Posttraumatic Stress Disorder for the Department of Veterans Affairs

Sriram Ramaswamy, M.D.; Vishal Madaan, M.D.; Faiz Qadri, M.D.;
Christopher J. Heaney, Psy.D.; Terry C. North, Ph.D.; Prasad R. Padala, M.D.;
Syed P. Sattar, M.D.; and Frederick Petty, M.D., Ph.D.

Posttraumatic stress disorder (PTSD) is a major mental disorder associated with significant morbidity, psychosocial impairment, and disability. The diagnosis of PTSD can be missed in a primary care setting, as patients frequently present with somatic complaints or depression and are often reluctant to discuss their traumatic experiences. As recent studies of veterans returning from the Gulf War and the Iraqi War suggest high rates of PTSD, the U.S. Department of Veterans Affairs (VA) Hospitals are gearing up to face this challenge. It is important to screen these veterans for symptoms of PTSD and make an appropriate referral if required. In this article, we attempt to review PTSD with a special focus on the VA population. In addition to discussing the epidemiology, diagnosis, and treatment options for PTSD, we also suggest screening questions for both combat-related and military sexual trauma-related PTSD.

(*Prim Care Companion J Clin Psychiatry* 2005;7:180-187)

Received Dec. 17, 2004; accepted May 17, 2005. From the Veterans Affairs Medical Center (Drs. Ramaswamy, Heaney, North, Padala, Sattar, and Petty); the Department of Psychiatry, Creighton University (Drs. Ramaswamy, Heaney, North, Sattar, and Petty); the Creighton-Nebraska Psychiatry Residency Program (Drs. Madaan and Qadri); and the University of Nebraska Medical Center (Dr. Padala), Omaha, Nebr.

In the spirit of full disclosure and in compliance with all ACCME Essential Areas and Policies, the faculty for this CME activity were asked to complete a full disclosure statement. The information received is as follows: Dr. Sattar is a consultant for AstraZeneca, Eli Lilly, Bristol-Myers Squibb, Janssen, and Abbott; has received grant/research support from AstraZeneca and Bristol-Myers Squibb; and has received honoraria from and is a member of the speakers/advisory boards for AstraZeneca and Abbott. Dr. Petty is a consultant for Bristol-Myers Squibb, Abbott, and Eli Lilly; has received grant/research support from Eli Lilly, Janssen, Bristol-Myers Squibb, Abbott, Pfizer, and Ortho-McNeil; has received honoraria from Eli Lilly, Bristol-Myers Squibb, Abbott, and Janssen; and is a member of the speakers/advisory boards for Eli Lilly, Bristol-Myers Squibb, Janssen, and Ortho-McNeil. Drs. Ramaswamy, Madaan, Qadri, Heaney, North, and Padala have no significant commercial relationships to disclose relative to this article.

Corresponding author and reprints: Sriram Ramaswamy, M.D., Staff Psychiatrist, Department of Mental Health and Behavioral Sciences, Omaha Veterans Affairs Medical Center, 4101 Woolworth Avenue, Omaha, NE 68105 (e-mail: sramaswamy@creighton.edu).

Posttraumatic stress disorder (PTSD) is a common and often severe mental disorder, with symptoms manifesting after exposure to a traumatic event. Initially in this article, we briefly review the history, definition, and causes of PTSD for the benefit of our primary care colleagues. We also discuss the prevalence of PTSD in the general population as well as specifically among veterans and thereafter provide an overview of the recent data on the prevalence of PTSD among U.S. military personnel in Iraq and Afghanistan. Finally, we discuss the diagnosis and management of PTSD.

HISTORICAL PERSPECTIVE

While the diagnostic term *posttraumatic stress disorder* has been adopted only recently, descriptions suggestive of the syndrome have been found in ancient Greek literature, including Homer's *The Iliad*.¹ "Cardiorespiratory syndrome" (characterized by postwar palpitations, shortness of breath, chest pain, and easy fatigue) was described by Dr. DaCosta in 1872 among veterans of the Battle of Gettysburg in the American Civil War.¹

Possibly the first "molecular" theory of PTSD arose after World War I, when survivors of the trench warfare were found to develop neurologic symptoms, such as tremor, amnesia, stuttering, and ataxia, after combat. It was then believed that explosions drove molecules of air through the skull and caused brain damage termed "shell shock."¹

The first psychological theories of PTSD were developed during and after World War II when the term *combat neurosis* was used to describe these symptoms.¹ In 1957, the "rape trauma syndrome"² was described, which emphasized the common symptomatology of PTSD, whether caused by civilian trauma, such as rape, or by combat trauma. Finally, in 1980, the American Psychiatric Association, in the DSM-III, published the first diagnostic criteria for PTSD and introduced the term *posttraumatic stress disorder* as the official term used to identify this condition.

DEFINITION AND CLASSIFICATION

PTSD is characterized by exposure to a traumatic event followed by reexperiencing of the traumatic event, avoid-

ance of stimuli associated with the trauma, and symptoms of increased arousal for at least 1 month.³ Stress occurs whenever there is change or “a challenge to homeostasis”^{4,5} that forces adaptation to that change. While stress has a neutral emotional valence (adaptive versus maladaptive; good versus bad), “traumatic stress” is qualitatively different and is always potentially harmful. Traumatic stress is a catastrophic, unexpected, uncontrolled event of extreme negative valence, capable of creating fear or horror in anyone. Examples of traumatic stress include combat, child abuse or neglect, domestic violence, rape and sexual assault, life-threatening accidents, torture, criminal assault, natural disasters (fires, floods, tornados, hurricanes), and man-made disasters (terrorism, bombing).

The response to the traumatic stress is the key to determining whether an individual develops PTSD later. The reason some individuals are resilient or resistant to traumatic stress, while others are not, is a focus of intense research. Animal models of stress reaction such as “learned helplessness” are shedding light on the underlying psychophysiology of stress susceptibility.^{6,7}

Traumatic stress is a fairly common life event. Estimates from the National Comorbidity Survey (NCS)⁸ showed that 60.7% of men and 51.2% of women reported exposure to at least 1 lifetime traumatic event. The same study highlighted that 20% of the women and 8% of the men exposed to trauma developed PTSD. Similarly, Breslau et al.⁹ reported that about 10% of men and 17% of women go on to develop severe PTSD after exposure to traumatic stress. However, if moderate and milder cases of PTSD are also included, about one third of individuals will develop PTSD after experiencing traumatic stress.

ETIOLOGY AND VULNERABILITY

PTSD may be unique among psychiatric disorders since it usually has a known etiology, is potentially preventable, and has a predominately environmental cause. While most other major psychiatric disorders have a strong genetic component (50%–80%),^{10,11,12} PTSD has a less frequent genetic association (30%).¹³ The cause of PTSD, obviously, is exposure to traumatic stress. However, as noted above, not everyone develops PTSD after exposure to traumatic stress. Thus, there are vulnerability factors. Among veterans of the Vietnam War, the strongest predictor of PTSD after combat was believed to be the duration of exposure to combat.¹⁴ Another study showed that Vietnam veterans with a history of childhood abuse and those who experienced a greater number of traumatic events prior to joining the military were more likely to have PTSD.¹⁵ Other vulnerability factors include a history of another mental illness, a family history of psychiatric disorders, borderline personality disorder, history of prior PTSD, and female gender. A recent meta-

analysis of predictors of PTSD or of its symptoms showed that peritraumatic psychological processes (and not prior characteristics), including amnesia and hyperarousal, are the strongest predictors of PTSD.¹⁶

Ample data exist that PTSD has a well defined biology. One interesting finding is that persons with PTSD appear to have a smaller hippocampus than persons without PTSD,^{17,18} which implies a dysregulation in the formation and registration of memories. The hippocampus is also highly sensitive to stress hormones, suggesting that high levels of these hormones may have caused brain damage. Recent studies have linked the decreased hippocampal volumes to a history of childhood trauma.^{19,20} Other brain imaging studies have found that amygdaloid activity is increased in persons with PTSD when they are reminded of the trauma.^{21,22} The amygdala is the region of the brain where emotions are regulated and is considered to be the anxiety center of the brain.

Another interesting biological finding in PTSD is low levels of the stress hormone cortisol. Several studies have reported low levels of cortisol in persons with PTSD.^{23,24} Initially, this was interpreted as a depletion effect, thinking that the severity of the stress experience in PTSD had “used up” the cortisol. Another explanation for this finding may be that individuals, who are unable to mount a normal stress response by increasing cortisol, are more vulnerable to developing PTSD after exposure to traumatic stress. This alternate hypothesis is supported by studies demonstrating that low cortisol levels immediately following trauma is a risk factor for developing PTSD at a later date.^{25,26}

HOW COMMON IS PTSD?

While in the past, there has been an unfortunate bias among some researchers about the validity of the diagnosis of PTSD, we now know that not only is PTSD real, but it is also common. The NCS, a large-scale epidemiology study using DSM-III-R criteria, found that the lifetime population prevalence of PTSD in the United States is 8%, with a prevalence of 10% in women and 5% in men.⁸ However, studies using DSM-IV criteria have reported higher lifetime prevalences for PTSD.^{27,28} This disorder is particularly common in primary care settings²⁹ and is associated with significant morbidity, psychosocial impairment, and distress. In these settings, PTSD is a persistent illness,³⁰ and it is unfortunate that substantial proportions of primary care patients go untreated or receive inadequate treatment.³¹

PTSD IN VETERANS

Among veterans of the Vietnam conflict, 14% of males had current PTSD at the time of survey, and 30% had lifetime PTSD.³² In female veterans of Vietnam, 8% had cur-

Table 1. Soldiers and Marines Who Met Screening Criteria for a Mental Disorder After Deployment to Iraq^a

Mental Disorder	Army Group, % (N/N)	Marine Group, % (N/N)
Depression	7.9 (66/840)	7.1 (55/775)
Anxiety	7.9 (66/839)	6.6 (51/776)
Posttraumatic stress disorder	12.9 (114/881)	12.2 (99/811)
Any of the above	17.1 (151/882)	15.6 (127/813)
Alcohol misuse	20.6 (168/815)	29.4 (219/744)

^aAdapted with permission from Hoge et al.³⁸**Table 2. Combat Experiences Reported by Soldiers and Marines After Deployment to Iraq^a**

Traumatic Stress	Army Group, % (N/N)	Marine Group, % (N/N)
Being attacked or ambushed	89 (789/883)	95 (764/805)
Receiving incoming artillery, rocket, or mortar fire	86 (753/872)	92 (740/802)
Being shot at or receiving small arms fire	93 (826/886)	97 (779/805)
Seeing dead bodies or human remains	95 (832/879)	94 (759/805)
Knowing someone seriously injured or killed	86 (751/878)	87 (693/797)
Responsible for death of an enemy combatant	48 (414/871)	65 (511/789)

^aAdapted with permission from Hoge et al.³⁸

rent and 25% had lifetime PTSD.³² These surveys were conducted upon return from Vietnam. After the first Gulf War, 10% of combat veterans developed PTSD.³³ After the conflict in Somalia, 8% of service members developed PTSD.³⁴

The U.S. Department of Veterans Affairs (VA) operates the United States' largest health care delivery system, providing comprehensive care to more than 5 million veterans through a network of 160 hospitals and 850 affiliated clinics.^{35,36} Approximately 600,000 veterans are affected by PTSD, and more than 200,000 have applied for VA service-connected benefits on the basis of PTSD.³⁷ PTSD thus is the most common psychiatric condition for which veterans seek VA service-connected benefits.

Recently, a survey was conducted in 4 U.S. combat infantry units before and after deployment to Iraq and Afghanistan.³⁸ The results of this survey were of great interest. Table 1 lists the percentage of service members deployed to Iraq who reported depression, anxiety, and/or PTSD or alcohol misuse. It is interesting how similar the results were in the Army and Marine study groups, except for perhaps more alcohol misuse among the Marines. Note that about 1 soldier or Marine in 6 endorsed a problem with depression, anxiety, and/or PTSD. These are, indeed, high rates of mental disorder and psychological distress.

Table 2 illustrates the kinds of traumatic stress to which the soldiers and Marines in Iraq were exposed.³⁸ As can be seen, virtually all of these combat infantrymen had

Table 3. Perceived Need for and Use of Mental Health Services Among Soldiers and Marines Who Met Criteria for Depression, Anxiety, or PTSD After Deployment to Iraq^a

Outcome	Army Group, % (N/N)	Marine Group, % (N/N)
Acknowledged a problem	78 (104/133)	86 (91/106)
Interested in receiving help	43 (58/134)	45 (47/105)
Received professional help in past year		
Overall (from any professional)	40 (56/140)	29 (33/113)
From a mental health professional	27 (37/138)	21 (24/112)

^aAdapted with permission from Hoge et al.³⁸**Table 4. Perceived Barriers to Seeking Mental Health Services Among Soldiers and Marines Who Met Criteria for a Mental Disorder^a**

Perceived Barrier	% (N/N)
I would be seen as weak	65 (413/640)
My unit leadership might treat me differently	63 (403/637)
Members of my unit might have less confidence in me	59 (377/642)
There would be difficulty getting time off for treatment	55 (354/643)
It would harm my career	50 (319/640)

^aAdapted with permission from Hoge et al.³⁸

been exposed to severe traumatic stress. From the combined data, we can conclude that about 1 service member in 8 developed PTSD. The survey also included questions regarding the perceived need for and use of mental health services by these soldiers and Marines. Table 3 illustrates the results.

The vast majority of the service members acknowledged a psychiatric problem,³⁸ which is a striking finding and illustrates a major challenge for the VA. However, less than half were interested in receiving help. Only about a third had received professional help of any kind, which could include, for example, counseling with a chaplain. Only about a fourth had received help from a mental health professional, which would include psychologists, psychiatric social workers, psychiatric nurses, and psychiatrists. The most common barriers to seeking mental health services as perceived by soldiers and marines are summarized in Table 4.

In the study, only 25% endorsed "mental health care doesn't work."³⁸ Thus, these service members are generally aware that they have a significant mental health problem, and they generally believe that mental health care is effective, but they do not seek help for fear that it would have a negative impact on their relations with peers and leaders and on their career. Mental health professionals, both in the armed forces and in the VA, clearly have a major challenge in decreasing the stigma associated with mental health care.

PTSD can also result from military sexual trauma. Military sexual trauma refers to both sexual harassment and sexual assault that occurs in military settings. In a survey, at least 23% of women using the VA health care system reported experiencing at least 1 sexual assault during

Table 5. The 3 Symptom Clusters of PTSD^a

Intrusive Category	Avoidant Category	Hyperarousal Category
Recurrent and intrusive distressing memories of the trauma	Avoiding thoughts, feelings, or conversations about the trauma	Difficulty falling or staying asleep
Recurrent and distressing dreams of the trauma (nightmares)	Avoiding activities, places, or people that remind of the trauma	Irritability or outbursts of anger
Acting or feeling as if the traumatic event was recurring (flashbacks)	Difficulty remembering important aspects of the trauma	Difficulty concentrating
Intense psychological distress at exposure to reminders of the trauma	Loss of interest or pleasure in normal activities	Feeling alert or watchful when there is no need to be
Physical distress (heart racing, shaking, etc) when reminded of the trauma	Feeling distant or cut off from others	Exaggerated startle response
	Difficulty experiencing normal feelings such as love and happiness	
	Feeling that the future will be cut short	

^aAdapted with permission from the American Psychiatric Association.³

their military service.³⁹ Another survey of Persian Gulf War military personnel found that rates of sexual assault (7%), physical sexual harassment (33%), and verbal sexual harassment (66%) were higher than those typically found in peacetime military samples.³⁹ There is also evidence that sexual trauma is a powerful contributor to the development of PTSD among female Vietnam veterans.⁴⁰ Sexual trauma associated with military service has some unique characteristics. It most often occurs in settings where the victim lives and works. In most cases, the victim must live and work closely with the perpetrators, which often leads to increased feelings of helplessness, powerlessness, and being at risk of additional victimization. The traumatic experience can cut short the career goals of the victims, and they frequently experience difficulties after being discharged from the military.³⁹

DIAGNOSIS OF PTSD

Considering that PTSD is a common psychiatric, and therefore medical, condition and that it causes great suffering and a high health care cost, it is surprising that the diagnosis of PTSD is often missed, especially in primary care settings.⁴¹

In PTSD, comorbidity is the rule, not the exception. The most common psychiatric comorbidities seen in patients with PTSD are substance use disorders; mood disorders, such as major or minor depressive disorders and bipolar disorder; other anxiety disorders, such as obsessive-compulsive disorder, panic disorder, or generalized anxiety disorder; and personality disorders, such as borderline personality disorder.⁸ Also, general medical conditions, such as anemia, arthritis, asthma, back pain, diabetes, eczema, kidney disease, lung disease, and ulcer are common among patients with PTSD.⁴² Usually it is the comorbid condition that prompts medical attention or requests for treatment, particularly in primary care settings.

Thus, in attending to the presenting problem, the health care provider often does not recognize, diagnose, or treat the underlying PTSD. Often, the trauma survivor may not be aware that the symptoms of PTSD are related to a dis-

ease process, and may have lived in quiet suffering for years. Thus, it is important for all health care professionals, not only in mental health, but also in primary care, internal medicine, and gynecology, to ask about trauma history when examining patients and incorporate some screening questions about PTSD into the medical evaluation. Health care professionals should remember that, depending on the treatment setting, 20% to 50% of patients either have PTSD or are at high risk for developing PTSD.

According to DSM-IV-TR,³ the diagnosis of PTSD requires (1) exposure to traumatic stress; (2) an initial reaction of horror, fear, or helplessness; and (3) symptoms in the 3 symptom clusters (Table 5). To have a diagnosis of PTSD, a person must have at least 1 intrusive symptom, at least 3 avoidant symptoms, and at least 2 arousal symptoms. Also, the symptoms need to last for 1 month or more and cause significant impairment.

SCREENING FOR PTSD

Screening for psychological illness, and especially PTSD, is based on symptom reporting.⁴³ While the U.S. Preventive Services Task Force currently does not recommend routine screening for PTSD, researchers have attempted to find short, validated tools for early detection and assessment of PTSD among veterans, especially in view of the ongoing events in Iraq, Afghanistan, and the Middle East. In a recent study⁴⁴ focusing on PTSD among soldiers returning from the Iraqi War, only about 3% of soldiers showed evidence of traumatic stress. The authors found 4 of the items on the U.S. Department of Defense Form 2796 (DD FORM 2796)⁴⁵ (Table 6) as being an effective screen and stated that the 17-item PTSD Checklist⁴⁶ was no better than the DD Form 2796 in terms of sensitivity and specificity. Interestingly, in screening done 3 to 6 months postdeployment, Hoge et al.³⁸ found rates of PTSD around 15% (as compared to 3% in the study by Bliese et al.⁴⁴), which questions when the most effective timing for screening is.

We suggest that health care professionals can perform a preliminary screen for PTSD by asking 2 questions (follow-up questions are in parentheses).

Table 6. Posttraumatic Stress Disorder Screening Questions From the Department of Defense Form 2796^a

Have you ever had any experience that was so frightening, horrible, or upsetting that, IN THE PAST MONTH, you...		
NO	YES	
☐	☐	Have had any nightmares about it or thought about it when you did not want to?
☐	☐	Tried hard not to think about it or went out of your way to avoid situations that remind you of it?
☐	☐	Were constantly on guard, watchful, or easily startled?
☐	☐	Felt numb or detached from others, activities, or your surroundings?

^aAdapted from U.S. Department of Defense Form 2796.⁴⁵ This material is in the public domain.

Question 1. “What’s the worst thing that ever happened to you?” (“Has there ever been a time when your life was in danger, you were seriously injured or harmed, you were threatened with death or serious injury, or you saw this happen to someone else?”)

If the person has not had traumatic stress exposure, or if the answer to the follow-up to question 1 is “no,” then the person cannot have PTSD, and the examiner may move on. If the person has had traumatic stress exposure, or if the answer to the follow-up to question 1 is “yes,” then the examiner should ask question 2.

Question 2. “Do memories of (the traumatic stress event) still bother you?” (“Do you have nightmares or flashbacks about the event? Do you think about it when you don’t want to? Do you get upset if something reminds you about it?”)

If the patient has a positive response to question 2, they should have additional evaluation for PTSD. This might take the form of a consultation request to a mental health service provider. Alternately, if the examiner wishes to ask screening questions about the other symptom clusters, we suggest the following:

Avoidant symptoms. “Do you avoid situations that might remind you of (the traumatic stress event)? Have your relationships suffered because of (the traumatic stress event)?”

Arousal symptoms. “Have you become more nervous since (the traumatic stress event)? Is it hard for you to relax because of (the traumatic stress event)? Do you have trouble sleeping because of (the traumatic stress event)?”

Screening all veterans for sexual trauma is also important. This should be done with compassion and sensitivity, recognizing the stigma associated with the experience while preserving confidentiality at all times. The National Center for Posttraumatic Stress Disorder has suggested asking the following questions: “While you were in the military did you experience any unwanted sexual attention, like verbal remarks, touching, or pressure for sexual favors?” and “Did anyone ever use force or the threat of force to have sex with you against your will?”⁴⁷

The importance of a sensitive approach by the providers doing the screening cannot be overstated. We are aware of cases in which veterans, male and female, denied any military sexual assault history for years. They

have told us one of the reasons they ultimately acknowledged the past sexual assault was because the screening providers, through their patience, eye contact, voice tone, and nonjudgmental demeanor, appeared to genuinely care for them and desire to refer them to the specialty services they required.

CLINICAL COURSE

The clinical course of PTSD has been studied in both combat veterans and rape trauma survivors. An acute stress disorder may follow a traumatic event. The diagnosis of acute stress disorder includes the criteria for PTSD but adds and emphasizes dissociative symptoms. Current research suggests that acute stress disorder is a predictor of subsequent PTSD.¹ There is some evidence that pharmacologic intervention shortly after trauma exposure may prevent the development of PTSD.^{48,49}

Once PTSD is diagnosed, recovery occurs most frequently in the first year; at 2 years, one half of the cases have remitted; and over the next 5 years, another 20% or so recover. Up to a third of survivors of traumatic stress may go on to develop chronic PTSD.¹ Once PTSD becomes chronic, spontaneous recovery is unusual, and symptoms tend to persist for years or decades. There is some evidence in veterans of the first Gulf War that symptoms can actually worsen over the course of time. Even when symptoms have disappeared, they can return, even years later, particularly if there is a new stress. The new stress need not be anything like the traumatic stress. For example, among veterans of World War II, symptoms of PTSD often returned after retirement. Stressors such as the death of a loved one, a heart attack, or job loss can bring back the symptoms of PTSD. It is important for individuals suffering from PTSD to realize that they are particularly sensitive to stress.

TREATMENT OF PTSD

Ideally, treatment of PTSD should be comprehensive and multidisciplinary. Counseling, education, and psychotherapy are important aspects of this treatment. The VA Medical Centers provide a network of specialized PTSD programs that offer veterans education, evaluation, and treatment conducted by mental health professionals

from a variety of disciplines. The outpatient PTSD programs include 3 basic types of clinics: (1) PTSD Clinical Teams provide one-to-one evaluation, education, and psychotherapy; (2) Substance Use PTSD Teams offer outpatient education, evaluation, and counseling for comorbid PTSD and substance abuse; and (3) Women's Stress Disorder Treatment Teams provide similar services to female veterans. The inpatient PTSD programs provide 24-hour nursing and psychiatric care. Finally, day treatment PTSD units provide one-to-one case management and counseling, group therapy, education, and activities to help clients live successfully with PTSD.⁵⁰

As mentioned, psychological treatment of PTSD is important and may be the treatment of first choice. Counseling and psychotherapy should be done by an experienced and skilled therapist because of the risk of "retraumatization." Often symptoms may worsen for a while during the therapy process as a person remembers the traumatic event.

Exposure therapy aims to disrupt the link between a trauma-related cue and the intense anxiety and avoidance that are typical for PTSD. The technique often stimulates the patient to experience a traumatic memory to modify the response to that memory. Repeated exposure helps patients learn to master their fear. In the hands of experienced therapists, it is a safe and effective treatment for PTSD⁵¹ and has the most empirical support.⁵²

Treatment issues in the case of military sexual trauma should address immediate health concerns, validate feelings, support existing adaptive coping strategies, promote development of new coping skills, and explore affective and cognitive reactions like fear, self-blame, anger, and disillusionment.

The pharmacologic management of PTSD usually begins with a selective serotonin reuptake inhibitor (SSRI). These medications include sertraline, paroxetine, fluoxetine, citalopram, and escitalopram. Of these, sertraline and paroxetine are approved by the U.S. Food and Drug Administration for treatment of PTSD. This class of medications is safe, has few side effects, and has no addictive potential.

If there is a partial response to the SSRIs, with some symptoms persisting, other psychotropic medications may be added. Anticonvulsants and mood-stabilizing agents like carbamazepine, lamotrigine, topiramate, and lithium can be useful for mood swings, irritability, and impulsivity and violent behavior.⁵³ Benzodiazepines such as alprazolam, clonazepam, and temazepam may be useful for arousal symptoms and to promote sleep^{54,55}; however, given the risk of promoting drug dependence and worsening PTSD by virtue of their dissociative and disinhibiting properties, benzodiazepines should be avoided in PTSD. If a hypnotic is necessary, the use of nonbenzodiazepines (e.g., zolpidem or zaleplon), antihistamines, or trazodone is recommended.^{54,55} Adjunctive use of atypical

antipsychotic medications, such as risperidone,⁵⁶ olanzapine,⁵⁷ and quetiapine,⁵⁸ can be helpful in combat-related PTSD. It is not unusual for a patient being treated for PTSD to be taking several psychotropic medications. The treating practitioner should be an experienced psychopharmacologist. The goal of medication management of PTSD is symptom reduction and stabilization. Even if all the symptoms do not entirely disappear, a patient may derive great benefit from getting a good night's sleep and being less anxious and irritable. Acute PTSD responds better than chronic PTSD. The earlier treatment begins, the better the outcome.

Education is another important aspect of PTSD treatment. Persons with PTSD are often unaware of the cause of their symptoms. Just being able to understand that PTSD is a biological disease, often associated with irritability, insomnia, and so forth, can be helpful to the patient in understanding the symptomatology and associated behavior. Group therapy is helpful for many patients. Often, persons with PTSD feel isolated and alone and doubt that anyone else in the world can understand what they have gone through and how they feel. Group therapy helps provide peer support and can expedite the healing process. Obviously, as in the treatment of other mental illnesses and behavioral disorders, psychotherapy may be combined with medication management.

The treating clinician should project a positive attitude, since many persons with PTSD are demoralized and may doubt they will ever improve. The family should be involved in the treatment plan. Treatment should emphasize hope, spirituality, growth, and healing. In this regard, a chaplain can be very helpful. Many persons with PTSD are suffering guilt and asking questions such as "why did this happen to me?" A priest, minister, or spiritual counselor can be helpful in providing comprehensive care for these survivors of trauma.

SUMMARY

PTSD is a serious and chronic mental illness that can cause severe psychosocial and functional impairment in the life of trauma survivors. It is characterized by intrusive, avoidant, and arousal symptoms after exposure to traumatic stress. PTSD affects more than 600,000 U.S. veterans and is the most common psychiatric condition for which veterans seek VA disability benefits. PTSD among veterans gains enormous significance especially in view of the ongoing events in Iraq and Afghanistan. Researchers are attempting to devise short, well-validated screening tools for early detection of this chronic illness. Treatment of PTSD usually involves both psychotherapy and medication. PTSD responds to treatment, sometimes with dramatic improvement, not only in the symptom profile, but also in the individual's overall quality of life.

Drug names: alprazolam (Xanax, Niravam, and others), carbamazepine (Tegretol, Carbatrol, and others), citalopram (Celexa and others), clonazepam (Klonopin and others), escitalopram (Lexapro), fluoxetine (Prozac and others), lamotrigine (Lamictal), lithium (Lithobid, Eskalith, and others), olanzapine (Zyprexa), paroxetine (Paxil, Pexeva, and others), quetiapine (Seroquel), risperidone (Risperdal), sertraline (Zoloft), temazepam (Restoril and others), topiramate (Topamax), trazodone (Desyrel and others), zaleplon (Sonata), zolpidem (Ambien).

REFERENCES

- Ornstein RD, Pitman RK. Trauma and post-traumatic stress disorder. In: Stern TA, Herman JB, eds. *Psychiatry Update & Board Preparation*. 2nd ed. New York, NY: McGraw-Hill; 2004:127–136
- Burgess AW, Holmstrom LL. Rape trauma syndrome. *Am J Psychiatry* 1974;131:981–986
- American Psychiatric Association. *Diagnostic and Statistical Manual of Mental Disorders, Fourth Edition, Text Revision*. Washington, DC: American Psychiatric Association; 2000
- Selye H. Stress and psychiatry. *Am J Psychiatry* 1956;113:423–427
- Selye H. What is stress? *Metabolism* 1956;5:525–530
- Petty F, Kramer GL, Wu J, et al. Posttraumatic stress and depression: a neurochemical anatomy of the learned helplessness animal model. *Ann N Y Acad Sci* 1997;821:529–532
- Petty F, Chae Y, Kramer G, et al. Learned helplessness sensitizes hippocampal norepinephrine to mild restress. *Biol Psychiatry* 1994;35:903–908
- Kessler RC, Sonnega A, Bromet E, et al. Posttraumatic stress disorder in the National Comorbidity Survey. *Arch Gen Psychiatry* 1995;52:1048–1060
- Breslau N, Davis GC, Andreski P, et al. Traumatic events and posttraumatic stress disorder in an urban population of young adults. *Arch Gen Psychiatry* 1991;48:216–222
- Henderson DC, Kunkel L, Goff DC. Psychosis and schizophrenia. In: Stern TA, Herman JB, eds. *Psychiatry Update & Board Preparation*. 2nd ed. New York, NY: McGraw-Hill; 2004:100
- Matthews J, Papakostas G. Mood disorders: depression. In: Stern TA, Herman JB, eds. *Psychiatry Update & Board Preparation*. 2nd ed. New York, NY: McGraw-Hill; 2004:105
- Perlis TH, Ghaemi SN. Bipolar disorder. In: Stern TA, Herman JB, eds. *Psychiatry Update & Board Preparation*. 2nd ed. New York, NY: McGraw-Hill; 2004:115
- True WR, Rice J, Eisen SA, et al. A twin study of genetic and environmental contributions to liability for posttraumatic stress symptoms. *Arch Gen Psychiatry* 1993;50:257–264
- Buydens-Branchey L, Noumair D, Branchey M. Duration and intensity of combat exposure and posttraumatic stress disorder in Vietnam veterans. *J Nerv Ment Dis* 1990;178:582–587
- Bremner JD, Southwick SM, Johnson DR, et al. Childhood physical abuse and combat-related posttraumatic stress disorder in Vietnam veterans. *Am J Psychiatry* 1993;150:235–239
- Ozer EJ, Best SR, Lipsey TL, et al. Predictors of posttraumatic stress disorder and symptoms in adults: a meta-analysis. *Psychol Bull* 2003;129:52–73
- Hedges DW, Allen S, Tate DF, et al. Reduced hippocampal volume in alcohol and substance naive Vietnam combat veterans with posttraumatic stress disorder. *Cogn Behav Neurol* 2003;16:219–224
- Villareal G, Hamilton DA, Petropoulos H, et al. Reduced hippocampal volume and total white matter volume in posttraumatic stress disorder. *Biol Psychiatry* 2002;52:119–125
- Bremner JD, Vythilingam M, Vermetten E, et al. MRI and PET study of deficits in hippocampal structure and function in women with childhood sexual abuse and posttraumatic stress disorder. *Am J Psychiatry* 2003;160:924–932
- Bremner JD, Randall P, Vermetten E, et al. Magnetic resonance imaging-based measurement of hippocampal volume in posttraumatic stress disorder related to childhood physical and sexual abuse: a preliminary report. *Biol Psychiatry* 1997;41:23–32
- Shin LM, Orr SP, Carson MA, et al. Regional cerebral blood flow in the amygdala and medial prefrontal cortex during traumatic imagery in male and female Vietnam veterans with PTSD. *Arch Gen Psychiatry* 2004;61:168–176
- Nutt DJ, Malizia AL. Structural and functional brain changes in post-traumatic stress disorder. *J Clin Psychiatry* 2004;65(suppl 1):11–17
- Yehuda R, Kahana B, Binder-Brynes K, et al. Low urinary cortisol excretion in Holocaust survivors with posttraumatic stress disorder. *Am J Psychiatry* 1995;152:982–986
- Yehuda R, Southwick SM, Nussbaum G, et al. Low urinary cortisol excretion in patients with posttraumatic stress disorder. *J Nerv Ment Dis* 1990;178:366–369
- Delahanty DL, Raimonde AJ, Spoonster E. Initial posttraumatic urinary cortisol levels predict subsequent PTSD symptoms in motor vehicle accident victims. *Biol Psychiatry* 2000;48:940–947
- Yehuda R, McFarlane AC, Shalev AY. Predicting the development of posttraumatic stress disorder from the acute response to a traumatic event. *Biol Psychiatry* 1998;44:1305–1313
- Breslau N, Kessler RC, Chilcoat HD, et al. Trauma and posttraumatic stress disorder in the community: the 1996 Detroit area survey of trauma. *Arch Gen Psychiatry* 1998;55:626–632
- Breslau N, Kessler RC. The stressor criterion in DSM-IV posttraumatic stress disorder: an empirical investigation. *Biol Psychiatry* 2001;50:699–704
- Bruce SE, Weisberg RB, Dolan RT, et al. Trauma and posttraumatic stress disorder in primary care patients. *Prim Care Companion J Clin Psychiatry* 2001;3:211–217
- Zlotnick C, Rodriguez BF, Weisberg RB, et al. Chronicity in posttraumatic stress disorder and predictors of the course of posttraumatic stress disorder among primary care patients. *J Nerv Ment Dis* 2004;192:153–159
- Rodriguez BF, Weisberg RB, Pagano ME, et al. Mental health treatment received by primary care patients with posttraumatic stress disorder. *J Clin Psychiatry* 2003;64:1230–1236
- Schlenger WE, Kulka RA, Fairbank JA, et al. The prevalence of post-traumatic stress disorder in the Vietnam generation: a multimethod, multisource assessment of psychiatric disorder. *J Trauma Stress* 1992;5:333–363
- Kang HK, Natelson BH, Mahan CM, et al. Post-traumatic stress disorder and chronic fatigue syndrome-like illness among Gulf War veterans: a population-based survey of 30,000 veterans. *Am J Epidemiol* 2003;157:141–148
- Litz BT, Orsillo SM, Friedman M, et al. Posttraumatic stress disorder associated with peacekeeping duty in Somalia for US military personnel. *Am J Psychiatry* 1997;154:178–184
- Wilson N, Kizer K. The VA health care system: an unrecognized national safety net. *Health Aff (Millwood)* 1997;16:200–204
- Rosenheck R, Massari. Wartime military service and utilization of VA health care services. *Mil Med* 1993;158:223–228
- Murdoch M, Hodges J, Cowper D, et al. Regional variation and other correlates of Department of Veterans Affairs Disability Awards for patients with posttraumatic stress disorder. *Med Care* 2005;43:112–121
- Hoge CW, Castro CA, Messer SC, et al. Combat duty in Iraq and Afghanistan, mental health problems, and barriers to care. *N Engl J Med* 2004;351:13–22
- Street A, Stafford J. Military sexual trauma: issues in caring for veterans. Washington, DC: National Center for PTSD. Available at: http://www.ncptsd.org/war/military_sexual_trauma.html. Accessed Oct 16, 2004
- Fontana A, Schwartz LS, Rosenheck R. Posttraumatic stress disorder among female Vietnam veterans: a causal model of etiology. *Am J Public Health* 1997;87:169–175
- Magruder KM, Frueh BC, Knapp RG, et al. Prevalence of posttraumatic stress disorder in Veterans Affairs primary care clinics. *Gen Hosp Psychiatry* 2005;27:169–179
- Weisberg RB, Bruce SE, Machan JT, et al. Nonpsychiatric illness among primary care patients with trauma histories and posttraumatic stress disorder. *Psychiatr Serv* 2002;53:848–854
- Rona RJ, Hyams KC, Wessely S. Screening for psychological illness in military personnel. *JAMA* 2005;293:1257–1260
- Bliese P, Wright K, Adler A, et al. Screening for traumatic stress among re-deploying soldiers. *Traumatic Stress Validation Report*. US Army Medical Research Unit-Europe. 2004. Available at: <http://www.usamrue.hq.usareur.army.mil/Traumatic%20Stress%20Screen%20Validation%20Report1.pdf>. Accessed March 31, 2005

45. Army Medical Surveillance Activity Documents. US Department of Defense Form 2796. Available at: http://amsa.army.mil/documents/DoD_PDFs/EPDA-042003.pdf. Accessed June 30, 2005
46. US Department of Defense Deployment Health Clinical Center. Standard Health Assessment Tools: Post Traumatic Stress Disorder (PTSD) Checklist (PCL); November 1994. Available at: <http://www.pdhealth.mil/guidelines/appendix4.asp>. Accessed June 24, 2005
47. Street A, Stafford J. Military sexual trauma. In: Schnurr PP, Cozza SJ, eds. *Iraq War Clinician Guide*. 2nd ed. Washington, DC: Department of Veterans Affairs National Center for PTSD; 2004: 68–69. Available at: http://www.ncptsd.va.gov/war/iraq_clinician_guide_v2/iraq_clinician_guide_ch_9.pdf. Accessed June 30, 2005
48. Vaiva G, Ducrocq F, Jezequel K, et al. Immediate treatment with propranolol decreases posttraumatic stress disorder two months after trauma. *Biol Psychiatry* 2003;54:947–949
49. Pitman RK, Sanders KM, Zusman RM, et al. Pilot study of secondary prevention of posttraumatic stress disorder with propranolol. *Biol Psychiatry* 2002;51:189–192
50. Department of Veterans Affairs. Specialized PTSD treatment programs in the US Department of Veterans Affairs: a national center for PTSD fact sheet. Available at: http://www.ncptsd.va.gov/facts/veterans/fs_treatment_programs.html. Accessed March 31, 2005
51. Rothbaum BO, Schwartz AC. Exposure therapy for posttraumatic stress disorder. *Am J Psychother* 2002;56:59–75
52. Foa EB. Psychosocial treatment of posttraumatic stress disorder. *J Clin Psychiatry* 2000;61(suppl 5):43–48
53. Schoenfeld FB, Marmar CR, Neylan TC. Current concepts in pharmacotherapy for posttraumatic stress disorder. *Psychiatr Serv* 2004;55:519–531
54. Asnis GM, Kohn SR, Henderson M, et al. SSRIs versus non-SSRIs in post-traumatic stress disorder: an update with recommendations. *Drugs* 2004;64:383–404
55. Davis LL, English BA, Ambrose SM, et al. Pharmacotherapy for posttraumatic stress disorder: a comprehensive review. *Expert Opin Pharmacother* 2001;2:1583–1595
56. Bartzokis G, Lu PH, Turner J, et al. Adjunctive risperidone in the treatment of chronic combat-related posttraumatic stress disorder. *Biol Psychiatry* 2005;57:474–479
57. Stein MB, Kline NA, Matloff JL. Adjunctive olanzapine for SSRI-resistant combat-related PTSD: a double-blind, placebo-controlled study. *Am J Psychiatry* 2002;159:1777–1779
58. Hamner MB, Deitsch SE, Brodrick PS, et al. Quetiapine treatment in patients with posttraumatic stress disorder: an open trial of adjunctive therapy. *J Clin Psychopharmacol* 2003;23:15–20

For the CME Posttest for this article, see pages 188–189.

For more CME activities, visit

CMEInstitute.MedFair.com

INSTRUCTIONS AND POSTTEST

Participants may receive up to 1 Category 1 credit toward the American Medical Association Physician's Recognition Award by reading the CME article and correctly answering at least 70% of the questions in the Posttest that follows:

Go to www.psychiatrist.com/cmehome to take this Posttest online and earn credit immediately.

Or

1. Read each question carefully and circle the answer on the Registration Form.
2. Type or print the registration information in the spaces provided and complete the evaluation.
3. Send the Registration Form to the address or fax number listed on the Registration Form.

All replies and results are confidential. Answer sheets, once graded, will not be returned. Unanswered questions will be considered incorrect and so scored. The CME Institute of Physicians Postgraduate Press, Inc. will keep only a record of participation, which indicates the completion of the activity and the designated number of Category 1 credits that have been awarded. Correct answers to the Posttest will be made available to the participants of this activity upon request after the submission deadline.

Accreditation Statement

The CME Institute of Physicians Postgraduate Press, Inc. is accredited by the Accreditation Council for Continuing Medical Education to provide continuing medical education for physicians.

Answer to Pretest: 1. a

1. The life stressor that usually is *not* associated with development of posttraumatic stress disorder (PTSD) is:
 - a. Sexual trauma or rape
 - b. Domestic violence or criminal assault
 - c. A terrorist bombing in which many were injured
 - d. The expected death of a loved one after chronic illness
2. Risk factors that may increase vulnerability to developing PTSD after traumatic stress exposure do *not* include:
 - a. Prior history of another psychiatric illness
 - b. Higher socioeconomic status
 - c. History of childhood abuse
 - d. History of prior PTSD
3. Which of the following is *not* a known biological finding in patients with PTSD?
 - a. Smaller hippocampus compared to patients without PTSD
 - b. Increased amygdaloid activity when patients are reminded of the traumatic event
 - c. Increased ventricular size
 - d. Low cortisol levels
4. In PTSD, comorbidity is:
 - a. Common but usually limited to another anxiety disorder
 - b. Rare
 - c. Dependent on the gender of the trauma survivor
 - d. Common and may include a wide range of other psychiatric or medical diagnoses
5. The authors indicate that primary care providers who screen patients for PTSD:
 - a. Should rely solely on patient responses to written questions and not engage in follow-up questions, to avoid upsetting patients
 - b. Can improve patient disclosure of past sexual traumas by virtue of a genuine, empathic, and nonjudgmental demeanor
 - c. Should make sure only male providers screen male patients for a history of sexual assault
 - d. Should be aware that veterans often are interested in receiving professional help because of increased public awareness of PTSD

REGISTRATION FORM

Circle the one correct answer for each question.

1. a b c d 4. a b c d
2. a b c d 5. a b c d
3. a b c d

Print or type

Name _____

Last 4 digits of Social Security Number _____
(Required to issue CME credit)

Birth Date (mm, dd, yy) _____
(Required to issue CME credit)

Degree _____ Specialty _____

Affiliation _____

Address _____

City, State, Zip _____

Phone () _____

Fax () _____

E-mail _____

☐ Hospital ☐ Private Practice ☐ Resident ☐ Intern

Deadline for submission

For a credit certificate to be issued, please complete this Registration Form no later than August 31, 2007. Online submissions will receive credit certificates immediately. Faxed or mailed submissions will receive credit certificates within 6 to 8 weeks.

Keep a copy for your files

Retain a copy of your answers and compare them with the correct answers, which will be published after the submission deadline.

Payment

If you complete the test online, no payment is necessary. A \$10 payment must accompany this form. You may pay by check, money order, or credit card (Visa or MasterCard). Make check or money order payable to Physicians Postgraduate Press, Inc. If paying by credit card, please provide the information below.

Check one: ☐ Visa ☐ MasterCard

Card number _____

Expiration date _____

Your signature _____

Please evaluate the effectiveness of this CME activity by answering the following questions.

1. Was the educational content relevant to the stated educational objective? ☐ Yes ☐ No
2. Did this activity provide information that is useful in your clinical practice? ☐ Yes ☐ No
3. Was the format of this activity appropriate for the content being presented? ☐ Yes ☐ No
4. Did the method of presentation hold your interest and make the material easy to understand?
☐ Yes ☐ No
5. Achievement of educational objective:
A. Enabled me to screen for and initiate treatment for posttraumatic stress disorder in the primary care setting. ☐ Yes ☐ No
6. Did this CME activity provide a balanced, scientifically rigorous presentation of therapeutic options related to the topic, without commercial bias? ☐ Yes ☐ No
7. Does the information you received from this CME activity confirm the way you presently manage your patients? ☐ Yes ☐ No
8. Does the information you received from this CME activity change the way you will manage your patients in the future? ☐ Yes ☐ No
9. If you answered yes, what change(s) do you intend to make in your practice?

10. Please offer comments and/or suggested topics for future CME activities.

11. How much time did you spend completing this CME activity?

12. What is your preferred format for CME activities? Circle one.
 - a. Print media (eg, journals, supplements, and newsletters)
 - b. Internet text
 - c. Internet multimedia
 - d. Audio CD
 - e. Live meeting

TEAR OUT AND SEND THIS PAGE, ALONG WITH YOUR PAYMENT, TO:

CME INSTITUTE • PHYSICIANS POSTGRADUATE PRESS, INC. • P.O. Box 752870 • MEMPHIS, TN 38175-2870

IF YOU ARE PAYING BY CREDIT CARD, YOU MAY FAX THIS PAGE TO: CME INSTITUTE AT 901-751-3444

QUESTIONS? CALL 1-800-489-1001 EXT. 8
WWW.PSYCHIATRIST.COM