4. This non-plasma antibody is unlikely to be mainly in the lymphocytes or lymphoid tissue.

5. Injection of antigen into immune, but not into normal, rabbits causes a profound leucopenia, which is due to a fall in the numbers of both lymphocytes and granulocytes. This phenomenon is not modified by the prior injection of cortisone in amounts sufficient to cause a lymphocytopenia.

We are most grateful to Professor A. Wormald for his constant advice and encouragement. We are indebted to the Medical Research Council and the Central Research Fund of the University of London for grants (to Professor A. Wormald) which have partly covered the costs of these investigations. We are also thankful to Miss A. Routledge for technical help throughout these investigations.

REFERENCES

- Abell, R. G. & Schenck, H. P. (1938). *J. Immunol.* **34**, 195.
- Banks, T. E., Tupper, R. L., Watts, R. W. E. & Wormald, A. (1955). *Biochem. J.* **59**, 149.
- Benson, J. A., Kim, K. S. & Bollman, J. L. (1955). *Amer. J. Physiol.* **182**, 217.
- Bierman, H. R., Kelly, K. H., Petrakis, N. L., Cordes, F., Foster, M. & Lose, E. A. (1951). *Blood*, **6**, 926.
- Cohen, S., Holloway, R. C., Matthews, C. & McFarlane, A. S. (1956). *Biochem. J.* **62**, 143.
- Dixon, F. J. & Talmage, D. W. (1951). *Proc. Soc. exp. Biol., N.Y.*, **78**, 123.
- Dougherty, T. F., Chase, J. H. & White, A. (1945). *Proc. Soc. exp. Biol., N.Y.*, **58**, 135.
- Dragstedt, C. A., Ramirez de Arellano, M., Lawton, A. H. & Youmans, G. P. (1940). *J. Immunol.* **39**, 537.
- Fischel, E. E., LeMay, M. & Kabat, E. A. (1949). *J. Immunol.* **61**, 89.
- Francis, G. E., Hawkins, J. D. & Wormald, A. (1955). *Biochem. J.* **59**, vii.
- Francis, G. E., Hawkins, J. D. & Wormald, A. (1957). *Biochem. J.* **65**, 560.
- Francis, G. E., Mulligan, W. & Wormald, A. (1954). *Isotopic Tracers*. London: Athlone Press.
- Gitlin, D. & Janeway, C. A. (1953). *Science*, **118**, 301.
- Harris, S. & Henle, W. (1948). *J. Immunol.* **59**, 9.
- Hawkins, J. D. (1957). *Biochem. J.* **67**, 537.
- Heidelberger, M. & Kendal, F. E. (1935). *J. exp. Med.* **62**, 697.
- Knox, W. C. & Endicott, F. C. (1950). *J. Immunol.* **65**, 523.
- Kolmer, J. A. & Boerner, F. (1945). *Approved Laboratory Technique*, 4th ed. New York: D. Appleton-Century-Crafts Inc.
- Lecomte, J. & Hugues, J. (1956). *Int. Arch. Allergy, N.Y.*, **8**, 72.
- Proom, H. (1943). *J. Path. Bact.* **55**, 419.
- Sanders, A. G., Florey, H. W. & Barnes, J. M. (1940). *Brit. J. exp. Path.* **21**, 254.
- Stavitsky, A. B. (1952). *J. Immunol.* **69**, 63.
- Stavitsky, A. B., Stavitsky, R. & Ecker, E. E. (1949). *J. Immunol.* **63**, 389.
- Sterling, K. & Gray, S. J. (1950). *J. clin. Invest.* **29**, 1614.
- van der Scheer, J., Bohnel, E., Clarke, F. H. & Wyckoff, R. W. G. (1942). *J. Immunol.* **44**, 165.

The Effects of 8-Azaguanine on the Biosynthesis of Ribonucleic Acid in *Bacillus cereus*

By H. G. MANDEL* AND R. MARKHAM

Agricultural Research Council, Virus Research Unit, Moltano Institute, University of Cambridge

(Received 15 November 1957)

The incorporation of the guanine analogue 8-azaguanine into the micro-organism *Bacillus cereus* has already been described (Matthews & Smith, 1956; Mandel, 1957; Mandel, Sugarman & Apter, 1957*b*). Almost all the combined 8-azaguanine was found in the ribonucleic acid (RNA) fraction of the cells (Mandel *et al.* 1957*b*) and much of it was released from this fraction by pancreatic ribonuclease in the form of 8-azaguanosine 2':3'-phosphate (Matthews & Smith, 1956). When [¹⁴C]8-azaguanine was used, a small amount of the analogue was found in the deoxyribonucleic acid (DNA) fraction, and the 'acid-soluble' fraction of the bacteria was also found to be radioactive

(Mandel *et al.* 1957*b*). The addition of a purine or a purine compound to a bacterial culture in which growth had been inhibited by the presence of 8-azaguanine had the effect of restoring normal growth. Simultaneously, some of the 8-azaguanine was lost from the RNA fraction and was released into the medium (Mandel, 1957).

In the present investigation the effects of 8-azaguanine on synthesis of ribonucleic acid in *B. cereus* were examined. Since the analogue was incorporated into the RNA it seemed possible that a number of alternative effects might be expected. One possibility was that the total amount of RNA per cell might increase in order to compensate to some extent for the presence of the unnatural base. On the other hand, the normal amount of RNA could be present with some of its purines replaced

* Present address: Department of Pharmacology, The George Washington University School of Medicine, Washington, D.C., U.S.A.

by the analogue. It also seemed likely that a study of the distribution of the analogue in the polynucleotide chains might indicate the way in which the synthesis of such chains proceeded.

Studies were also carried out to determine the presence of intermediary metabolites of 8-azaguanine because the effects of the analogue might be due to the blocking of some coenzyme systems in addition to its incorporation into the RNA. As far as was possible the behaviour of the normal base guanine was studied under similar conditions and the mechanism of the reversal of growth inhibition by this substance was investigated. Preliminary accounts of some of this work have been published (Mandel & Markham, 1957*a, b*).

MATERIALS AND METHODS

The micro-organism used in the present investigations, *Bacillus cereus* strain 569 *H*, has been used in previous work (Matthews & Smith, 1956; Mandel, 1957; Mandel *et al.* 1957*b*). This is a constitutive penicillinase-producing mutant isolated from the inducible penicillinase-producing wild strain by Kogut, Pollock & Tridgell (1956). The medium was that of Matthews & Smith (1956) and contained casein hydrolysate (Vitamin-free Casamino Acids, Difco Laboratories Inc., Detroit, Mich., U.S.A.). For experiments with ^{32}P , a low-phosphate medium was prepared in which 2 g. of 2-amino-2-hydroxymethylpropane-1:3-diol/l. and 1.5 g. of K_2SO_4 /l. replaced 3.35 g. of the usual 3.5 g. of KH_2PO_4 /l.

[2- ^{14}C]5-Azaguanine was furnished by the Southern Research Institute, Birmingham, Ala., U.S.A.; [2- ^{14}C]uracil was synthesized as reported previously (Mandel & Brown, 1952); [8- ^{14}C]guanine and $^{32}\text{PO}_4^{3-}$ ions (carrier-free) were purchased from the Radiochemical Centre, Amersham, Bucks. [^{14}C]Formate was obtained from Isotopes Specialties Co., Burbank, California, U.S.A.

Bacterial growth and inhibition. Bacteria were grown in a shaking device at 37° so that aeration was optimum. A heavy inoculum of cells was used (5×10^8 cells/100 ml. of medium). The bacteria grew exponentially with a characteristic mean generation time of 55 min. Growth was measured turbidimetrically in a Unicam SP. 500 spectrophotometer at 540 m μ , such a procedure being a satisfactory index of the concentration of bacterial suspensions (Mandel, 1957).

Growth was recorded as the difference between the optical density reading (ΔE_{540}) after a given time and that at the introduction of the isotope.

8-Azaguanine was dissolved in 0.5% Na_2CO_3 soln. The analogue was added to the micro-organisms to a final concentration of 20 μg ./ml. of medium, when the bacterial concentration reached 10^8 cells/ml. ($E_{540} = 0.2$). The experiments were usually terminated at twice this bacterial concentration. Within 30 min. after the addition of the analogue the generation time increased approximately fivefold.

Fractionation of micro-organisms. Bacteria were harvested by centrifuging, followed by washing in 0.15 M-NaCl or 0.15 M-sodium phosphate buffer, pH 7. The residue was extracted with 5% (w/v) trichloroacetic acid at room

temperature, the extract being shaken several times with ether to remove the acid, followed by evaporation and chromatography on Whatman no. 3 MM paper in the 70% (v/v) propan-2-ol-aq. NH_3 soln. system of Markham & Smith (1952*a*). The residue after the extraction with trichloroacetic acid was taken up in 70% (w/v) ethanol at 60°, followed by a wash in ethanol-ether (1:1, v/v). The washed residue was then hydrolysed by N-KOH at room temperature overnight to liberate the ribomononucleotides from the ribonucleic acids.

Electrophoretic separation of mononucleotides. The alkaline digest was neutralized with N-HClO₄ to remove excess of alkali and K^+ ions and centrifuged at 0°. The supernatant solution was applied to paper strips moistened with 0.05 M-ammonium formate buffer, pH 3.1, and the separation was carried out at that pH for 7 hr. at 20 v/cm. (Markham & Smith, 1952*b*). At this pH the four pairs of normal nucleotides and 8-azaguanilyc acid may be easily separated from each other. Guanylic acid and 8-azaguanilyc acid were clearly split into the 2'- and 3'-isomers, which move at slightly different mobilities in this system. The order of increasing mobilities was cytidylic, adenylic, guanylic, 8-azaguanilyc and uridylic acid. The nucleotides were detected by their ultraviolet-light absorption, and were eluted in 0.1 N-HCl, with an appropriate blank treated similarly, and measured spectrophotometrically. Radioactivity was measured in a liquid counter (20th Century Electronics, London, type M2H) for ^{32}P , or after plating on metal dishes and counting with an end-window tube.

Uptake of isotopic precursors. Two bacterial cultures of identical bacterial concentration received 20 μg . of the labelled compound/ml., and then 20 μg . of 8-azaguanine/ml. was added to one culture. Precursors studied included $^{32}\text{PO}_4^{3-}$ ions, sodium [^{14}C]formate and [^{14}C]uracil. Samples of both cultures were taken at frequent intervals, and bacterial growth was measured turbidimetrically. The samples were mixed immediately afterwards with an equal volume of 10% (w/v) trichloroacetic acid, and the mixture was filtered with membrane filters (Schleicher and Schuell Co., Keene, New Hampshire, U.S.A.) as described by Britten, Roberts & French (1955). The filters were then counted for radioactivity. This procedure has been used previously to measure accurately the effects of an agent inhibiting growth on bacterial biosynthetic pathways (Bolton & Mandel, 1957).

Assay for ribose. The orcinol method for ribose, as described by Miltzer (1946) and modified by Markham (1955), was used.

Metabolite analysis. For an analysis of the metabolites of 8-azaguanine occurring in the bacterial medium after inhibition of bacterial growth by the analogue, the medium was centrifuged and evaporated almost to dryness. The residue was then subjected to chromatography on Whatman no. 3 MM paper in the 70% (v/v) propan-2-ol-aq. NH_3 soln. system of Markham & Smith (1952*a*).

Ribonuclease treatment. Crystalline ox-pancreas ribonuclease (E. Light and Co., Colnbrook, Bucks) was dissolved in water and heated to 100° to inactivate other enzymes. Solutions to be digested were adjusted to about pH 7 by the addition of a dilute aq. NH_3 soln. with bromothymol blue as an internal indicator. The bacterial sample, after the usual extractions, was ground with fine glass powder to rupture cell walls; the mixture was incubated with approx. 0.3 mg. of pancreatic ribonuclease/ml. for

3 days at 37° in the presence of CHCl_3 to prevent bacterial growth. The mixture was centrifuged, and the soluble portion concentrated by evaporation and applied on Whatman no. 3MM paper for chromatography in the propan-2-ol-aq. NH_3 soln. solvent system for 3 days.

RESULTS

Nucleic acid content and composition

To determine whether the quantity of polynucleotide per cell was affected by 8-azaguanine, the uptake of $^{32}\text{PO}_4^{3-}$ ions by a bacterial culture partially inhibited by the drug, and by a control culture, was measured in relation to growth. It was found that for the same increase in cell mass, approximately twice as much of the isotope was incorporated by the culture grown in the presence of 8-azaguanine. In a similar experiment in which $[2\text{-}^{14}\text{C}]\text{uracil}$ served as the only labelled supplement to the two proliferating cultures, the presence of 8-azaguanine caused 100% increase in incorporation of the isotope (Fig. 1). The uptake of labelled formate into the nucleic acid was also increased by the analogue when comparisons were made after similar increases in cell mass. Bacteria grown in the presence of 8-azaguanine also contained more orcinol-reactive material than did a control culture. These results are consistent with the conclusion that in the presence of 8-azaguanine, the inhibition of growth of *B. cereus* was associated with incorporation of the analogue into nucleic acids (Matthews & Smith, 1956; Mandel *et al.* 1957*b*) as well as with an increase in the content of the latter.

The relative compositions of the ribopolynucleotides isolated from bacteria grown in the presence and in the absence of 8-azaguanine were compared by measurement of the isolated mononucleotides.

The possibility of contamination by traces of the analogue base from the medium, where it occurs in excess, was thus ruled out. Since it is impossible to separate those micro-organisms in the bacterial

culture formed before the addition of the inhibitor from those grown later, the sensitivity of measurement was increased by adding ^{32}P to the culture simultaneously with the inhibitor. By using a radioactive tracer in this manner, the biosynthetic processes occurring only after the addition of 8-azaguanine could be examined, since in the growing bacterial cell no turnover of nucleic acids has been found (Roberts, Abelson, Cowie, Bolton & Britten, 1955). Comparisons were made with a control culture which received the same amount of labelled phosphate added at the same bacterial concentration. Both cultures were harvested after approximately equal increases in cell mass, and the nucleotides separated electrophoretically as described above. The various nucleotides were identified as shown in Fig. 2. Guanylic acid fluoresced under acidic conditions only (Smith & Markham, 1950) whereas 8-azaguanilyc acid fluoresced in acidic and ammoniacal atmospheres (Matthews, 1954). All the nucleotides were radioactive because of the presence of ^{32}P .

The relative composition of nucleotides from the two bacterial cultures is shown in Table 1. It was demonstrated that, of the normal nucleotides, only the relative content of guanylic acid was affected. When only labelled nucleotides were considered, as in the ^{32}P assay column, it appears that slightly more 8-azaguanilyc acid was synthesized than guanylic acid, and that the summation of these two nucleotides produced polynucleotides resembling the composition of normal nucleic acid.

Since other phosphorus-containing metabolites may be associated with the bacterial nucleic acid fraction, as has been reported for similar alkali digests obtained from rat liver (Davidson & Smellie, 1952), the above results were calculated also in terms of molar activities. These values have been converted into relative molar activity so that the average activity of the nucleotides of the control culture would equal 100. The addition of ^{32}P to a control bacterial culture growing exponentially in a medium containing some phosphate labels all nucleotides in accordance with their rate of formation, so that the molar activities would be expected to be identical. It will be noted that this procedure does not involve any assumptions about the pathway of synthesis of the nucleotides. Since the activities were equal, as shown in Table 2, extraneous ^{32}P compounds apparently did not interfere with the present measurements.

On the other hand, if a change in the relative rates of formation of the nucleotides should occur as, for example, an effect produced by the simultaneous introduction of the analogue with the ^{32}P , the molar activities of the nucleotides would differ from 100. Thus if the synthesis of a nucleotide were accelerated in relation to growth, its relative molar

Fig. 1. Uptake of $[^{14}\text{C}]\text{uracil}$ (counts/min.) by cultures of *Bacillus cereus* in relation to growth measured turbidimetrically at 540 μ . ○, Control culture; ●, culture treated with 8-azaguanine.

Fig. 2. Electrophoretic separation of mononucleotides from normal and 8-azaguanine-containing nucleic acid. (a) Control; (b) from nucleic acid containing 8-azaguanine. Compounds from top (positive end) down are uridylic (8-azaguanidylic), guanylic, adenylic and cytidylic acid. Lane 1, ultraviolet-light absorption due to nucleotides; lane 2, acid fluorescence of guanylic and 8-azaguanidylic acid; lane 3, ammoniacal fluorescence of 8-azaguanidylic acid only; lane 4, radioautograph from ^{32}P -labelled nucleotides.

Table 1. *Nucleotide composition of nucleic acids from cultures of Bacillus cereus grown in the presence and the absence of 8-azaguanine*

Analogue and $^{32}\text{PO}_4^{3-}$ ions were added simultaneously to growing bacterial cultures.

Nucleotide isolated	Polynucleotide composition (moles/100 moles of nucleotides)			
	Extinction assay		^{32}P assay	
	Inhibited	Control	Inhibited	Control
Cytidylic acid	21.8	23.2	20.8	22.2
Uridylic acid	26.5	24.1	26.4	25.2
Adenylic acid	23.0	23.5	23.0	24.7
Guanylic acid	20.6	29.2	13.5	27.9
8-Azaguanidylic acid	8.1	0.0	16.2	0.0

activity would be increased. Since the tracer was added to the two cultures when both had identical bacterial concentrations, the corresponding nucleo-

tides isolated after further growth from the control and inhibited cultures would be diluted by identical quantities of unlabelled material.

Thus the relationship between the increased synthesis (n_i'/n_c') and the increased relative molar activity would be:

$$\frac{m_i}{m_c} = \frac{R_i}{n_i' + n_i} \bigg/ \frac{R_c}{n_c' + n_c},$$

where m_i represents molar activity of [^{32}P]nucleotides from inhibited culture, m_c molar activity of [^{32}P]nucleotides from control culture, R_i radioactivity of [^{32}P]nucleotides from inhibited culture, R_c radioactivity of [^{32}P]nucleotides from control culture, n_i' moles of [^{32}P]nucleotides from inhibited culture synthesized after addition of tracer, n_c' moles of [^{32}P]nucleotides from control culture synthesized after the addition of tracer, and n_c ($=n_i$) moles of [^{32}P]nucleotides from either culture synthesized before addition of tracer. Since

$$\frac{R_i}{n_i'} = \frac{R_c}{n_c'},$$

$$\frac{m_i}{m_c} = \frac{(n_c' + n_c) n_i'}{(n_i' + n_c) n_c'}.$$

In these experiments n_c is proportional to the turbidimetric reading, measured at 540 m μ , of the bacterial cultures at the time of addition of ^{32}P , and n_c' corresponds to the difference between the readings at harvest and at the addition of ^{32}P .

Since the control and inhibited cultures did not proliferate to exactly the same extent, the molar activities of the inhibited cultures were adjusted by use of the formula

$$\frac{m_{i_2}}{m_{i_1}} = \frac{n_{i_2}(n_{i_1} + n_c)}{n_{i_1}(n_{i_2} + n_c)},$$

where the subscripts 1 and 2 refer to the bacterial concentrations at various harvests.

The results of these calculations are shown in Table 2.

The activity of 8-azaguanylic acid is higher than that of the other nucleotides because only the former compound is undiluted by unlabelled ribonucleotide existing before the introduction of the ^{32}P . Again, the guanylic acid isolated from the inhibited micro-organisms differs in molar activity from the other natural nucleotides. The increase in nucleic acid content is also confirmed. The similarity between the adjusted activity of guanylic acid of the nucleic acid containing 8-azaguanylic acid and the activities of the nucleotides from control nucleic acid signifies that synthesis of guanylic acid was not impaired by the action of the inhibitor. Instead, the biosynthesis of cytidylic acid, uridylic acid and adenylic acid was approximately doubled in these experiments, and 8-azaguanylic acid replaced guanylic acid in that nucleic acid formed in excess of the normal quantity.

When [^{14}C]8-azaguanine was used to inhibit the bacterial culture and the usual procedure for the isolation of nucleotides was followed, the 8-azaguanylic acid was radioactive. No activity could be found in the position where 8-azadenylic acid would be expected, if such conversion into this compound had taken place.

Treatment with ribonuclease

A ground bacterial preparation which contained 1 part of [^{14}C]8-azaguanine to 3 parts of nucleic acid guanine was treated with ribonuclease and chromatographed. Additional fractionation was obtained by paper electrophoresis at pH 3.5 in ammonium formate buffer, perpendicular to the chromatographic separation. The normal constituents of such a digest can readily be recognized by this procedure (Markham & Smith, 1952*b*), and those containing 8-azaguanine were located by fluorescence and measured by their radioactivity. The latter metabolites were eluted and subjected to further electrophoresis at pH 1.9 in glycine buffer (3.75 g. of glycine/l. of 0.05N-HCl) and 7.4 in phosphate buffer (KH_2PO_4 , Na_2HPO_4 ; 3:7, w/w; 0.05M). The separation at pH 1.9 gave information about the composition of the non-8-azaguanine part of the fractions, whereas the treatment at pH 7.4 revealed particularly the terminal groupings of the fractions, such as the nucleosides, the nucleoside 2':3'-phosphates, or the nucleoside 3'-phosphates. The results of this fractionation scheme are shown in Fig. 3 and Table 3, where the mobilities relative to 8-azaguanylic acid are shown, and the identity of the various fractions is suggested. Although final identification of the many fragments would require a more rigorous analysis with larger amounts of material, the chromatographic and electrophoretic behaviour of the various compounds matched quite well the values calculated for those metabolites of 8-azaguanine which might be expected in such a ribonuclease digest. The largest portion of the analogue was recovered in the form of 8-azaguanosine 2':3'-phosphate, as well as dinucleotides terminated by the latter.

Table 2. *Relative molar radioactivities of nucleotides isolated from nucleic acids of cultures of Bacillus cereus grown in the presence and the absence of the inhibitor 8-azaguanine*

Analogue and $^{32}\text{PO}_4^{3-}$ ions were added simultaneously to growing bacterial cultures. Figures in parentheses are adjusted to allow for unequal proliferation of inhibited and control cultures.

Nucleotide isolated	Treated		Control	
	Expt. 1	Expt. 2	Expt. 1	Expt. 2
Cytidylic acid	1.12 (1.44)	1.29 (1.48)	0.97	1.01
Uridylic acid	1.18 (1.51)	1.26 (1.45)	1.05	1.02
Adenylic acid	1.18 (1.51)	1.23 (1.42)	1.05	1.01
Guanylic acid	0.77 (0.99)	0.94 (1.08)	0.97	0.95
8-Azaguanylic acid	2.37	1.94	—	—
Average	— (1.49)*	— (1.45)*	1.00	1.00
Increase in nucleic acid content over control (%)	109	93	0	0

* For inhibited cultures, averages include only cytidylic acid, uridylic acid, and adenylic acid.

8-Azaguanine metabolites in the acid-soluble fraction

The relatively small amount of 8-azaguanine metabolites in the acid-soluble fraction has complicated the search for the minor constituents of these samples. Chromatography regularly revealed two major metabolites which have been identified as 8-azaguanosine 5'-phosphate and 8-azaguanosine. These compounds were usually present in the molar ratio 1:4.

8-Azaguanosine 5'-phosphate. This nucleotide had the same R_F in propan-2-ol-aq. NH_3 soln. as 8-azaguanosine 3'-phosphate, obtained by alkaline hydrolysis of polynucleotides. The low R_F value was similar to that of guanylic acid. When subjected to paper electrophoresis at pH 9.2 in sodium borate buffer (19 g./l.; 0.05M), the metabolite showed a slightly higher mobility than that of the 3' derivative. Upon rechromatography in propan-2-ol-aq. NH_3 soln., however, the 3' derivative retained its original R_F , whereas 8-azaguanosine 5'-phosphate, which had formed a borate complex, had a much lower R_F value in this solvent. Whereas 8-azaguanosine 5'-phosphate has a *cis*-glycol structure the 3' derivative does not, and therefore it does not form such a complex.

For additional confirmation, snake-venom phosphodiesterase was prepared by the method of Butler (1955). Treatment of bacterial nucleic acids containing 8-azaguanine with this enzyme by incubation at 37° for 24 hr. liberated 8-azaguanosine 5'-phosphate, which was identical in behaviour with the compound isolated from the acid-soluble fraction.

Fig. 3. Separation pattern of compounds derived from ribonuclease treatment of nucleic acids containing 8-azaguanine. Mixture is placed at the top left, followed by chromatography in propan-2-ol-aq. NH_3 soln. in a vertical direction and electrophoresis at pH 3.5, negatively charged compounds moving towards the right. Black spots are due to 8-azaguanine metabolites, with numbers corresponding to those in Table 3. Other spots (lettered as in Table 3) are due to normal fragments isolated from the digest, as indicated.

Table 3. Characterization of 8-azaguanine metabolites found and expected after ribonuclease treatment of 8-azaguanine containing nucleic acids from *Bacillus cereus*

Chromatography zones of oligonucleotides after Markham & Smith (1952b). A = Adenylic; C = cytidylic; G = guanylic; U = uridylic; Z = 8-azaguanilyc. Mobilities are given relative to 8-azaguanilyc acid (=100). Negative values refer to mobilities in the opposite direction from 8-azaguanilyc acid.

Compound isolated	Relative (%)	Chromatographic position	Suggested identity	Mobility					
				pH 3.5		pH 1.9		pH 7.4	
				Obs.	Calc.	Obs.	Calc.	Obs.	Calc.
1	2	1	AZU	115	(125)	—	(100)	100	(105)
2	2	1	AZZ*	115	(125)	—	(90)	85	(80)
3	8	1	ZU	155	(145)	135	(155)	105	(115)
4	2	1	ZZ*	155	(145)	—	(145)	75	(85)
5	10	2	Z	100	(100)	100	(100)	100	(100)
6	9	2	GZ*	125	(130)	135	(120)	85	(85)
7	4	4	ZC*	70	(85)	75	(60)	90	(85)
8	5	4	AZ*	100	(105)	75	(60)	70	(80)
9	58	6	Z*	100	(100)	100	(100)	70	(65)
			ZC	—	(85)	—	(60)	—	(110)
			AAZ*	—	(100)	—	(35)	—	(80)
			ZU*	—	(145)	—	(155)	—	(90)
			Z (side)	—	(0)	—	(-10)	—	(15)
			AZ (side)	—	(35)	—	(-25)	—	(45)
			GZ (side)	—	(55)	—	(35)	—	(45)

* Nucleoside 2':3'-phosphate.

8-Azaguanosine. This compound had the characteristically high R_f of other nucleosides, and was identified by comparing its mobility at various pH values with the calculated mobilities. It lacked mobility at pH 3.5, and moved more slowly than 8-azaguanine at pH 7.4 (phosphate buffer) and pH 9.2 (glycine buffer). In sodium borate buffer at pH 9.2, however, it moved faster than the free base. These results were anticipated for a nucleoside, since the characteristic *cis*-glycol structure enabled it to form a charged borate complex, resulting in greater mobility in such a system. After acid hydrolysis of the metabolite, 8-azaguanine was recovered.

To establish further the presence of the nucleoside, the metabolite was oxidized with sodium metaperiodate, followed by incubation at 37° at pH 10 overnight, the method of Whitfeld (1954) for *cis*-glycols being used. The products were purified by chromatography in propan-2-ol-aq. NH_3 soln. followed by paper electrophoresis at pH 9.2. 8-Azaguanine was liberated during the oxidation, confirming the identity of the metabolite as the nucleoside.

For a comparison with the authentic nucleoside, 8-azaguanic acid prepared by alkali hydrolysis of bacterial nucleic acids was caused to react with prostatic phosphomonoesterase at 37° for 30 min. The resulting nucleoside corresponded in its properties with the isolated metabolite.

Other compounds. 8-Azaguanine was also present in this fraction. Occasionally 8-azaxanthosine was found in trace amounts. This compound had a slightly greater R_f in propan-2-ol-aq. NH_3 soln., and had slightly greater mobility in sodium borate buffer at pH 9.2 than did 8-azaguanosine. The presence of 8-azaxanthosine in ethanol extracts of *B. cereus* has already been reported (Matthews & Smith, 1956). After periodate oxidation (Whitfeld, 1954) 8-azaxanthine was isolated.

No other metabolites could be detected.

8-Azaguanine metabolites in the bacterial medium

Chromatographic analysis of the bacterial medium after inhibition of bacterial growth by 8-azaguanine regularly revealed two major metabolites, in addition to 8-azaguanine which was present in excess. These were 8-azaguanosine and another metabolite which has not yet been identified.

8-Azaguanosine. This compound was identical with that from the acid-soluble fraction. Approximately 5% of the 8-azaguanine of the medium had been converted into the nucleoside.

Unidentified metabolite. The second metabolite, which was derived from another 5% of the analogue of the medium, had a very small R_f in propan-2-ol-aq. NH_3 soln., thus resembling a

nucleotide. The metabolite, however, evidently did not contain phosphate, since it did not become radioactive when a culture growing in the presence of ^{32}P was inhibited with unlabelled 8-azaguanine.

The metabolite apparently has a relatively strong basic group, since at pH 1.9 it moved toward the cathode more rapidly than did 8-azaguanine. At pH 9.2 the metabolite had a greater mobility than 8-azaguanine though less than that of 8-azaguanic acid, thus suggesting a relatively strong acidic group. No difference could be detected in mobility in glycine or sodium borate buffers at pH 9.2, indicating the lack of a sugar group containing *cis*-glycol. After treatment with N-HCl at 100° for 1 hr. the compound retained its original properties. The metabolite apparently contained the C-2 atom since it became labelled when $[2\text{-}^{14}\text{C}]\text{-8-azaguanine}$ was used. It therefore seemed likely that the purine skeleton remained intact, and that a conjugate was formed with a part having a strong acid and a part with a basic charge. The identity of this substance is being studied to determine whether a conjugate of 8-azaguanine is formed with an amino acid. Normal purines have been known to form such complexes (Carter & Cohen, 1956).

Other compounds. A trace of 8-azaxanthosine was found in the medium also.

8-Azaguanosine ejected by bacteria into a medium supplemented with purine compounds

The introduction of a purine compound, such as guanosine, into a culture already inhibited with 8-azaguanine results in the recovery of the growth rate, together with the ejection of some of the 8-azaguanine from the RNA fraction of the cells, as already described (Mandel, 1957). In the present experiments, bacteria inhibited with labelled 8-azaguanine were transferred to a new medium containing either guanosine or guanylic acid. After 1 hr., during which the culture was allowed to resume its normal growth, the medium was analysed for radioactivity. 8-Azaguanosine was isolated and shown to be identical with the metabolite characterized above.

It was observed in these investigations that after the addition of guanylic acid to the bacterial culture, a large amount of this nucleotide was converted into the nucleoside which was present in the medium, indicating that extensive dephosphorylation had taken place. It is very likely therefore that the 8-azaguanosine isolated after the reversal of growth was derived from an 8-azaguanosine phosphate by dephosphorylation.

No other metabolites of 8-azaguanine, nor other nucleotides or nucleosides, could be detected in the medium.

*Comparisons with [¹⁴C]guanine metabolism
in Bacillus cereus*

Several experiments were carried out by procedures identical with those described, but in which labelled guanine was used rather than 8-azaguanine, to serve as comparison.

Nucleic acid fraction. Isolation of the mononucleotides after paper electrophoresis at pH 3.1 revealed that some of the labelled guanine was converted into adenine. The molar activity of adenylic acid was found to be 45% of that of guanylic acid.

Acid-soluble fraction. Guanosine and guanosine 5'-phosphate were isolated, as well as several other compounds of higher R_F , in propan-2-ol-aq. NH_3 soln. Guanosine 5'-pyrophosphate could not be detected.

Bacterial medium. Traces of two metabolites were found, probably guanine and guanosine, but no compound could be isolated which corresponded in R_F in the propan-2-ol-aq. NH_3 soln. system to the unidentified metabolite present in the medium of a bacterial culture inhibited by 8-azaguanine.

DISCUSSION

The replacement of guanylic acid by 8-azaguanilyc acid in a portion of the ribonucleic acids of a bacterial culture of *B. cereus* growing in the presence of 8-azaguanine is not surprising in view of the close structural analogy of the latter to guanine. In experiments with tobacco mosaic virus and 8-azaguanine, where the extent of incorporation of the analogue was extremely small, the 8-azaguanine in the nucleic acid fraction was found to be present at the expense of guanine (Matthews, 1953). There is no information about whether a change in the nucleic acid content of the virus occurred, but such a change would not be expected to be appreciable because of the slight extent of incorporation of the analogue.

It has been reported previously (Mandel, 1957) that the relative extent of incorporation of the analogue decreased when it was added at increasing bacterial concentrations. Under conditions such that a large incorporation of the analogue took place, it was shown that a culture inhibited by 8-azaguanine incorporated 60% as much of the analogue into nucleic acids as that amount of exogenous guanine taken up by a control culture during the same increase in cell mass (Mandel, 1957). In the normal micro-organism growing in the presence of [¹⁴C]guanine, however, a certain amount of this purine was converted into the adenine of polynucleotides. Calculations based on the relative molar activities of guanine and adenine, as well as their relative abundances in the nucleic acids, showed that only about 70% of the guanine

taken up by a culture of *B. cereus* was used for synthesis of guanylic acid. Since synthesis of guanylic acid was not affected in these experiments, it appears that guanylic acid and 8-azaguanilyc acid were synthesized at approximately the same rate.

The formation and incorporation of 8-aza-adenine from 8-azaguanine could not be observed in any of the experiments, even though the reverse process, i.e. the formation of 8-azaguanine from 8-aza-adenine and the subsequent incorporation of 8-azaguanine, has been described for tobacco mosaic virus (Matthews & Smith, 1955). In *B. cereus*, however, 8-aza-adenine is not converted into the guanine analogue (Smith & Matthews, 1957), nor could conversion of 8-azaguanine into 8-aza-adenine be demonstrated in the mouse (Mandel, Carló & Smith, 1954).

The great preponderance of cyclic 8-azaguanosine 2':3'-phosphate in the ribonuclease digest is in agreement with the findings of Matthews & Smith (1956). The isolation in the present experiments of what appear to be dinucleotides ending in cyclic 8-azaguanilyc acid strengthens their hypothesis that incorporation of 8-azaguanine is greatest in the terminal portions of polynucleotide chains. The concentration of recovered cyclic 8-azaguanilyc acid implies that polynucleotides built during inhibition with 8-azaguanine are relatively short.

The ejection of 8-azaguanine from the bacterial cell under conditions of reversal of inhibition of growth suggests that during the restoration of growth the 8-azaguanine incorporated primarily at the terminal ends of polynucleotides was removed, since no other nucleotide fragments were evident. Since the 8-azaguanine was localized principally at these end groups, such a degradation of part of the nucleic acid could take place without any additional turnover. The possibility remains, however, that normal nucleotides from degraded nucleic acids might be re-utilized preferentially and thus would not be found in the medium.

Recent experiments on the incorporation of thiouracil into tobacco mosaic virus (Mandel, Markham & Matthews, 1957a) have revealed a similar distribution of that analogue in polynucleotides. It was concluded that thiouracil occupied several of the positions normally occupied by uracil, with a tendency for the analogue to be concentrated at the ends of polynucleotide chains, particularly more at one end than the other.

It is interesting that 8-azaguanosine 5'-phosphate was isolated from the acid-soluble fraction of the inhibited micro-organisms, in view of the probability that this compound acts as a precursor for formation of nucleic acid, perhaps by an enzyme similar to that isolated by Grunberg-Manago & Ochoa (1955) from *Azotobacter*, which uses nucleoside 5'-pyrophosphates as substrates. Although the

last-named derivative of 8-azaguanine could not be isolated, it is possible that this compound was present in too small an amount to be detectable, or else that it was immediately incorporated into the polynucleotide. Under the experimental conditions, guanosine 5'-pyrophosphate could not be detected after bacteria had been grown in the presence of labelled guanine. The presence of guanosine pyrophosphate in the acid-soluble portion of rat tissues has been reported (Schmitz, Hurlbert & Potter, 1954).

8-Azaguanosine 5'-phosphate could well form the pyrophosphate or some other intermediate which is directly responsible for the inhibition of growth produced by 8-azaguanine, perhaps by antagonizing guanosine pyrophosphate. This last-named compound acts, among other ways, as a cofactor for the incorporation of amino acids into proteins in a rat-liver enzyme system (Keller & Zamecnik, 1956). The 8-azaguanine metabolite could then be incorporated into the nucleic acids. The lack of a direct relationship between the extent of inhibition of growth caused by the analogue and the total amount of 8-azaguanine incorporated into the nucleic acid fraction (Mandel, 1957) might be explained by the existence of such an intermediate.

The isolation of both 8-azaguanosine 5'-phosphate in the acid-soluble fraction and the presence of (cyclic) 8-azaguanosine 2':3'-phosphate in the ribonuclease digest resulting from incorporation of 8-azaguanine suggests the existence of more than one route of synthesis of nucleic acid and incorporation of the analogue.

In addition to the pathway involving the polymerization of the 5'-phosphates, as postulated by Grunberg-Manago & Ochoa (1955), the cyclic nucleotides might also serve as intermediates in synthesis of nucleic acid, perhaps with the utilization of a reversible enzyme similar to ribonuclease (Heppel, Whitfeld & Markham, 1955).

SUMMARY

1. During the partial inhibition of growth of *Bacillus cereus* produced by 8-azaguanine, the newly formed ribonucleic acids (RNA) contain 8-azaguanine as well as the normal base guanine.

2. The relative base composition of the nucleic acids remains unchanged, with the analogue substituting for some of the guanine.

3. The ribonucleic acid guanylic acid content of the inhibited bacteria is similar to that of the control micro-organisms, but the amount of adenylic, cytidylic and uridylic acid increases in proportion to the quantity of the analogue incorporated.

4. It is deduced that 8-azaguanine replaces

guanylic acid in the excess portion of the bacterial nucleic acids only.

5. A major portion of the incorporated 8-azaguanine is present at the ends of polynucleotide chains in the form of cyclic 8-azaguanosine 2':3'-phosphate.

6. Additions of guanine or guanosine to a culture inhibited by 8-azaguanine apparently leads to the loss of 8-azaguanine from the polynucleotides.

7. 8-Azaguanosine 5'-phosphate has been isolated from the acid-soluble fraction of the cells, and its possible function in synthesis of nucleic acid is discussed.

One of the authors (H. G. M.) is indebted to the Commonwealth Fund for an Advanced Fellowship in the Health Field. A part of this research was aided by research grant CY-2978 from the National Cancer Institute, U.S. Public Health Service.

REFERENCES

- Bolton, E. T. & Mandel, H. G. (1957). *J. biol. Chem.* **227**, 833.
 Britten, R. J., Roberts, R. B. & French, E. F. (1955). *Proc. nat. Acad. Sci., Wash.*, **41**, 863.
 Butler, G. C. (1955). *Methods in Enzymology*, vol. 2, p. 561. Ed. by Colowick, S. P. & Kaplan, N. O. New York: Academic Press Inc.
 Carter, C. E. & Cohen, L. H. (1956). *J. biol. Chem.* **222**, 17.
 Davidson, J. N. & Smellie, R. M. S. (1952). *Biochem. J.* **52**, 599.
 Grunberg-Manago, M. & Ochoa, S. (1955). *J. Amer. chem. Soc.* **77**, 3165.
 Heppel, L. A., Whitfeld, P. R. & Markham, R. (1955). *Biochem. J.* **60**, 8.
 Keller, E. B. & Zamecnik, P. C. (1956). *J. biol. Chem.* **221**, 45.
 Kogut, M., Pollock, M. R. & Tridgell, E. J. (1956). *Biochem. J.* **62**, 391.
 Mandel, H. G. (1957). *J. biol. Chem.* **225**, 137.
 Mandel, H. G. & Brown, C. L. (1952). *J. Amer. chem. Soc.* **74**, 2439.
 Mandel, H. G., Carló, P. E. & Smith, P. K. (1954). *J. biol. Chem.* **206**, 181.
 Mandel, H. G. & Markham, R. (1957a). *Biochem. J.* **65**, 25p.
 Mandel, H. G. & Markham, R. (1957b). *Trans. Faraday Soc.* **53**, 260.
 Mandel, H. G., Markham, R. & Matthews, R. E. F. (1957a). *Biochim. biophys. Acta*, **24**, 205.
 Mandel, H. G., Sugarman, G. I. & Apter, R. A. (1957b). *J. biol. Chem.* **225**, 151.
 Markham, R. (1955). In *Modern Methods of Plant Analysis*. Ed. by Paech, K. & Tracey, M. V. Berlin: Julius Springer.
 Markham, R. & Smith, J. D. (1952a). *Biochem. J.* **52**, 552.
 Markham, R. & Smith, J. D. (1952b). *Biochem. J.* **52**, 558.
 Matthews, R. E. F. (1953). *Nature, Lond.*, **171**, 1065.

- Matthews, R. E. F. (1954). *J. gen. Microbiol.* **10**, 521.
Matthews, R. E. F. & Smith, J. D. (1955). *Advanc. Virus Res.* **3**, 49.
Matthews, R. E. F. & Smith, J. D. (1956). *Nature, Lond.*, **177**, 271.
Militzer, W. E. (1946). *Arch. Biochem.* **9**, 85.
Roberts, R. B., Abelson, P. H., Cowie, D. B., Bolton, E. T. & Britten, R. J. (1955). *Studies of Biosynthesis in Escherichia coli*, publication no. 607. Washington: Carnegie Institution.
Schmitz, H., Hurlbert, R. B. & Potter, V. R. (1954). *J. biol. Chem.* **209**, 41.
Smith, J. D. & Markham, R. (1950). *Biochem. J.* **46**, 509.
Smith, J. D. & Matthews, R. E. F. (1957). *Biochem. J.* **66**, 323.
Whitfield, P. R. (1954). *Biochem. J.* **58**, 390.

Effect of Nitrophenols and Halogenophenols on the Enzymic Activity of Rat-liver Mitochondria

By V. H. PARKER

Toxicology Research Unit, Medical Research Council Laboratories, Woodmansterne Road, Carshalton, Surrey

(Received 27 November 1957)

Many studies on oxidative phosphorylation involve the use of 2:4-dinitrophenol (DNP) as an inhibitor of that process. Investigations of this kind and of other manifestations of DNP activity are numerous (for a review, see Brody, 1955). In some respects, other nitrophenols and also halogenophenols have similar biological properties to DNP (Clowes, Keltch, Strittmatter & Walters, 1950; De Deken, 1955). In particular, pentachlorophenol has been the subject of a study in parallel with DNP (Weinbach, 1954, 1956), and *p*-nitrophenol has been used by Terner (1954).

In spite of the extensive use of DNP in the study of plant- and animal-tissue metabolism, little is known of the way in which it produces its effects. There is general acceptance of the hypothesis that DNP accelerates the breakdown of some 'high-energy' compound formed during the synthesis of adenosine triphosphate coupled with the oxidation of substrates of the tricarboxylic acid cycle (Lardy, 1956; Slater & Lewis, 1954). The more restricted theory that DNP causes the breakdown of some phosphorylated intermediate was proposed by Hunter (1951), who also suggested the possibility that DNP itself may be phosphorylated with the formation of dinitrophenyl phosphate. The latter hypothesis has been rejected mainly on the grounds of the stability of dinitrophenyl phosphate (Lardy & Wellman, 1953). A similar argument has been advanced by Weinbach (1956) and Bringham, Brinch-Johnsen & Walaas (1956).

It is the object of the present work to record some observations on the similarities and differences of a series of nitro- and halogeno-phenols with respect to their action on the enzymic activity of rat-liver mitochondria *in vitro* and also to record the relative stabilities of some substituted phenyl

phosphates in mitochondrial preparations. For the purpose of this paper, characteristic effects of DNP on liver mitochondria have been compared with effects of the other phenols. There are three well-established biological actions of DNP, namely, the inhibition of oxidative phosphorylation (Loomis & Lipmann, 1948), which is usually demonstrated in the presence of hexokinase and glucose, the stimulation of oxidation of various substrates (Potter & Recknagel, 1951), which is revealed in the absence of hexokinase and glucose, and finally, the stimulation of adenosine triphosphatase activity (Myers & Slater, 1957).

EXPERIMENTAL

Methods

Preparation of mitochondria. The isolation of rat-liver mitochondria and the source and treatment of the reagents used in the enzymic procedures were identical with those described by Aldridge (1957) except in two points of detail. The homogenizer was a Nelco (Measuring and Scientific Equipment, London) used at 5000 rev./min. for 1½ min. and the cell debris and nuclei were removed by centrifuging at 650 g instead of 800 g. By this method mitochondria were produced which satisfied the criterion of normality as defined by Aldridge (1957) and earlier by Potter & Recknagel (1951). That is, in the presence of an appropriate substrate and cofactors these mitochondria are capable of a large increase of oxidative activity if presented with either DNP or hexokinase and glucose, or other substances which increase the rate of utilization of adenosine triphosphate.

Oxidative phosphorylation. The incubations were made in Warburg flasks. Each flask contained 0.5 ml. of mitochondrial suspension equivalent to 250 mg. wet wt. of liver, 0.00115 M-sodium adenosine triphosphate, 0.00115 M-adenosine monophosphate, 0.125 M-KCl, 0.014 M-MgCl₂·6H₂O, 0.001 M-disodium ethylenediaminetetra-acetic acid, 0.0154 M-potassium phosphate, 0.06 M-glucose, 0.0166 M-glycylgly-