ation with KMnO_4 had m.p. 175–176° after recrystallization from ethanol (microrosettes of needles) (Found: C, 48.3; H, 2.7; N, 1.85; Br, 29.1. $\text{C}_{22}\text{H}_{20}\text{O}_3\text{NBr}$ requires C, 47.9; H, 2.5; N, 1.75; Br, 29.9%). The *p*-bromophenacyl ester of authentic berberonic acid had the same m.p., 175–176°, and there was no depression on mixing.

Action of methylamine on sclerotiorin: *N*-methylsclerotiorazin. Sclerotiorin (0.25 g.) was treated with aqueous methylamine (33%, w/v; 5 ml.). The substance immediately became dark red but complete solution was not attained. The mixture was well stirred and all lumps were broken up. After 15 min., water (25 ml.) was added, the mixture was made just acid to Congo red and the solid (0.22 g.) was collected. Crystallization from aq. methanol (50%, v/v) furnished an *N*-methylsclerotiorazin as purple-black needles, m.p. 220–222° (Found: C, 65.3; H, 6.3; N, 3.8; Cl, 8.8. $\text{C}_{22}\text{H}_{26}\text{O}_4\text{NCl}$ requires C, 65.4; H, 6.5; N, 3.5; Cl, 8.8%).

Reduction of N-methylsclerotiorazin. *N*-Methylsclerotiorazin (0.93 g.) in acetic acid (40 ml.) was well stirred during the addition of Zn dust (1 g.). The colour of the solution changed from deep red to orange. The reaction was considered complete when no further lightening occurred (about 20 min.). The solution was filtered to remove residual Zn and poured into water. The gelatinous ppt. produced gradually crystallized in sheaves of needles. The product (0.74 g., m.p. 239°) formed golden-yellow needles of *deacetyldihydro-N*-methylsclerotiorazin, m.p. 239–241°, on recrystallization from methanol (Found: C, 65.9, 66.0; H, 7.4, 7.3; N, 3.5; Cl, 9.85, 9.9. $\text{C}_{20}\text{H}_{26}\text{O}_3\text{NCl}$ requires C, 66.0; H, 7.2; N, 3.85; Cl, 9.7%).

N-Methylsclerotiorazin (2 g.) was dissolved in methanol and reduced in presence of Zn dust (2 g.) and H_2SO_4 (50%, v/v; 2 ml.). The product was poured into water and the filtrate was distilled in the usual manner for collection of volatile acids. Titration of the distillate revealed that 46.8 ml. of 0.1*N*-volatile acid had been produced (94% of 1 mol.prop.). Conversion into the *p*-bromophenacyl ester afforded a product of m.p. 84–85°, not depressed on admixture with *p*-bromophenacyl acetate.

SUMMARY

1. The product of the reaction of sclerotiorin with ammonia, $\text{C}_{21}\text{H}_{24}\text{O}_4\text{NCl}$, named sclerotiorazin, is readily reduced by zinc and acetic acid, yielding sclerotazin ($\text{C}_{19}\text{H}_{22}\text{O}_3\text{NCl}$) and acetic acid.

2. By mild alkaline hydrolysis of sclerotazin the chlorine atom is eliminated and replaced by a hydroxyl group, affording hydroxydechlorosclerotazin, $\text{C}_{19}\text{H}_{23}\text{O}_3\text{N}$.

3. Sclerotazin forms a diacetate, from which the unsaturated side chain is removed by ozonolysis and replaced by a carboxyl group, affording the diacetate of sclerazinic acid $\text{C}_{11}\text{H}_8\text{O}_4\text{NCl}$.

4. Oxidation of sclerazinic acid with alkaline potassium permanganate solution gives berberonic acid. From these degradations a tentative partial molecular structure for sclerotazin is deduced.

5. Sclerotiorin reacting with methylamine produces *N*-methylsclerotiorazin ($\text{C}_{22}\text{H}_{26}\text{O}_4\text{NCl}$), which is reduced by zinc and acetic acid to deacetyldihydro-*N*-methylsclerotiorazin, $\text{C}_{20}\text{H}_{26}\text{O}_3\text{NCl}$.

We wish to thank Imperial Chemical Industries Ltd., Akers Research Laboratories, The Frythe, Welwyn, Herts, for carrying out some of the elementary analyses. The Uvispek Spectrophotometer used in this work was purchased by means of a grant from the Central Research Fund of London University.

REFERENCES

- Birkinshaw, J. H. (1952). *Biochem. J.* **52**, 283.
 Birkinshaw, J. H. (1956). *G. Microbiol.* **2**, 116.
 Curtin, T. P. & Reilly, J. (1940). *Biochem. J.* **34**, 1419.
 Powell, A. D. G., Robertson, A. & Whalley, W. B. (1956). *Spec. Publ. Chem. Soc.* **5**, 27.
 Raistrick, H. & Stössl, A. (1958). *Biochem. J.* **68**, 647.
 Watanabe, H. (1952). *J. pharm. Soc. Japan*, **72**, 807.

Comparative Detoxication

5. CONJUGATION OF AROMATIC ACIDS IN REPTILES: FORMATION OF ORNITHURIC ACID, HIPPURIC ACID AND GLUCURONIDES*

By J. N. SMITH

Department of Biochemistry, St Mary's Hospital Medical School, London, W. 2

(Received 28 January 1958)

The detoxication of aromatic acids in the animal body may involve condensation with a variety of compounds in different species and, as a rule, more than one type of conjugation is found in each species. Glycine conjugation occurs in most mammals (Williams, 1947), in frogs (Bunge &

Schmiedeberg, 1876; Komori, Sendju, Sagara & Takamatsu, 1926), in some birds (Wu Chang & Johnson, 1957) and in insects (Smith, 1955). Ester glucuronides can also be formed in mammals (Williams, 1947) and in the chick (Wu Chang & Johnson, 1957; B. C. Baldwin, D. Robinson & R. T. Williams, unpublished work). Glutamine conjugations, which have been thought peculiar to

* Part 4: Smith (1955).

man and the chimpanzee, have now also been reported in rats, rabbits and chickens (Hiratani, 1957). The conjugation of xanthuronic acid with serine has been reported in rats (Rothstein & Greenberg, 1957). The ornithine conjugation has hitherto been found only in birds, where ornithuric acids result from the detoxication of benzoic acid (Jaffe, 1877).

It has sometimes been suggested that reptiles might also form ornithuric acids since other aspects of their nitrogen metabolism are similar to those of birds (Baldwin, 1948; Sahyun, 1948). The limited data available, however, appear to indicate the operation of the more usual glycine conjugation in reptiles. Thus Komori *et al.* (1926) isolated hippuric acid from the urine of an unidentified chelonian after feeding sodium benzoate. In an early report Schiff (1859) also stated that hippuric acid crystals had been recognized in the urine of *Testudo tabulata*. Khalil (1947, 1948*a, b*, 1951) has quoted analyses for hippuric acid, measured by extraction with ether and nitrogen analysis, in the normal urine of the snakes *Zamenis diadema* and *Eryx thebaicus*, the turtle *Chelone mydas* and the lizard *Chalcides ocellatus*.

A number of aromatic acids have now been fed to six species of reptile and it is shown below that, while in some cases a small amount of glycine conjugation occurs, the major amino acid conjugate excreted is an ornithuric acid.

MATERIALS AND METHODS

Reference compounds. L(+) - Ornithine was prepared from L(+) - arginine by enzymic hydrolysis (Hunter, 1939) and treated, in *N*-Na₂CO₃ solution, with benzoyl chloride to give L(+) - ornithuric acid (*N*²N⁵-dibenzoylornithine), m.p. 183°, $[\alpha]_D^{25}$ 9.7°. DL-Ornithuric acid (m.p. 185°) was similarly prepared from DL-ornithine.

L(+) - Ornithine was shaken with a solution of *p*-nitrobenzoyl chloride in toluene and a slight excess of *N*-Na₂CO₃ at 0° till the purple colour faded. Acidification with conc. HCl gave L(+) - *N*²N⁵-bis-(*p*-nitrobenzoyl)ornithine (*pp'*-dinitro-ornithuric acid), which after crystallization from ethanol had m.p. 100°, solidifying and remelting at 150–160°, $[\alpha]_D^{25}$ 6.0° in 0.1*N*-NaOH (*c*, 1) (Found: C, 52.5; H, 4.8; N, 11.9. C₁₉H₁₈O₈N₄·Et·OH requires C, 52.9; H, 5.1; N, 11.8%). It had $\lambda_{\max}$ 270 m μ and $\epsilon_{\max}$ 24 000 in 0.1*N*-NaOH or 0.1*N*-HCl.

DL-*N*²N⁵-Bis-(*p*-nitrobenzoyl)ornithine, prepared similarly from DL-ornithine and crystallized from ethyl acetate and dilute acetic acid, had m.p. 193° (Found: C, 52.7; H, 4.2; N, 12.8. C₁₉H₁₈O₈N₄ requires C, 53.0; H, 4.2; N, 13.0%).

L(+) - *N*²N⁵-Bis-(*p*-nitrobenzoyl)ornithine (1.1 g.) in 100 ml. of H₂O was heated on a steam bath for 0.5 hr. with FeSO₄·7H₂O (7.2 g.) and K₂CO₃ (4.1 g.). The crystals (0.8 g.) which formed on cooling the filtrate were recrystallized from water and the L(+) - *N*²N⁵-bis-(*p*-aminobenzoyl)ornithine (*pp'*-diamino-ornithuric acid) had m.p. 182°, $[\alpha]_D^{25}$ 7° in 0.1*N*-NaOH (*c*, 1) (Found: C, 60.4; H, 5.9; N, 15.1. C₁₉H₂₂O₄N₄·0.5H₂O requires C, 60.1; H, 6.1; N, 14.8%).

DL-*N*²N⁵-Bis-(*p*-aminobenzoyl)ornithine, m.p. 191°, was similarly prepared from the inactive nitro compound (Found: C, 60.4; H, 5.8; N, 14.8. C₁₉H₂₂O₄N₄·0.5H₂O requires C, 60.1; H, 6.1; N, 14.8%). It had $\lambda_{\max}$ 273, $\epsilon_{\max}$ 32 000 in 0.1*N*-NaOH and $\lambda_{\max}$ 222, $\epsilon_{\max}$ 22 000 in 0.1*N*-HCl.

DL-*N*⁵-Benzyloxycarbonylornithine (Harris & Work, 1950) was dissolved in a slight excess of *N*-Na₂CO₃ at 0° and shaken with a solution in toluene of an equivalent amount of *p*-nitrobenzoyl chloride until the purple colour faded, the solution being kept at pH 10–11 with additions of 2*N*-NaOH. The precipitate which formed on acidification with conc. HCl was crystallized from methanol, and then ethanol, to give DL-*N*⁵-benzyloxycarbonyl-*N*²-(*p*-nitrobenzoyl)ornithine, m.p. 150–152° (Found: N, 9.8%; equiv. 412. C₃₀H₄₁O₇N₃ requires N, 10.1%; equiv. 415). This compound (2 g.) was reduced with H₂ and Adams Pd catalyst (0.2 g.) in 1:1 (v/v) aqueous methanolic solution (200 ml.). Concentration gave a crystalline precipitate (0.9 g.), which was recrystallized from H₂O. The DL-*N*²-(*p*-aminobenzoyl)ornithine had m.p. 190° (decomp.) (Found: C, 53.1; H, 6.9; N, 15.7. C₁₈H₁₇O₃N₃·H₂O requires C, 53.5; H, 7.1; N, 15.6%).

DL-*N*⁵-(*p*-Nitrobenzoyl)ornithine, m.p. 220° (Found: C, 51.4; H, 5.2; N, 14.9. C₁₈H₁₅O₅N₃ requires C, 51.2; H, 5.4; N, 14.9%), and DL-*N*⁵-(*p*-aminobenzoyl)ornithine, m.p. 215° (Found: C, 53.4; H, 7.0; N, 15.0. C₁₈H₁₇O₃N₃·H₂O requires C, 53.5; H, 7.1; N, 15.6%), were prepared as described by McGilvery & Cohen (1950) for the preparation of the corresponding optically-active compounds.

DL-Ornithine dissolved in *N*-Na₂CO₃ was shaken with salicylic azide, the solution being maintained at pH 10–11 by adding 2*N*-NaOH. After 0.5 hr. the solution was acidified with conc. HCl, and the precipitate removed, washed with CHCl₃ and crystallized from a large volume of water, in which it was sparingly soluble. The DL-*N*²N⁵-disalicyloyl-ornithine (*oo'*-dihydroxyornithuric acid) had m.p. 173° after drying over P₂O₅ (Found: C, 61.2; H, 5.2; N, 7.9. C₁₉H₂₀O₈N₄ requires C, 61.3; H, 5.4; N, 7.5%).

p-Aminobenzoylglucuronide was prepared biosynthetically (cf. Bray, Lake, Neale & Thorpe, 1948). *p*-Aminobenzoic acid (6 g.) was fed to three rabbits and the glucuronide gum obtained by the lead acetate procedure (cf. Kamil, Smith & Williams, 1951). This gum contained much free *p*-aminobenzoic acid, which crystallized on standing. The lead acetate separation was therefore repeated on the glucuronide gum and a second clear glucuronide gum (300 mg.) obtained free from *p*-aminobenzoic acid. This was treated with 100 mg. of *p*-anisidine in 2 ml. of water, when a crystalline precipitate of the *p*-anisidine salt of *p*-aminobenzoyl glucosiduronic acid was obtained (150 mg.). After crystallization from water it had m.p. 184°, $[\alpha]_D^{25}$ –5° in water (*c*, 1) (Found: C, 50.7; H, 5.8; N, 5.7. C₂₀H₂₄O₉N₂·2H₂O requires C, 50.8; H, 6.0; N, 5.9%).

Another preparation of the glucuronide gum, after free *p*-aminobenzoic acid has been filtered off, was methylated and acetylated (cf. Kamil *et al.* 1951) and the product crystallized repeatedly from ethanol to give a small yield of methyl (*p*-acetamidobenzoyl triacetylglucosid)uronate, m.p. 205°, $[\alpha]_D^{25}$ –28.5° in CHCl₃ (*c*, 1) (Found: N, 3.2. C₂₂H₂₅O₁₂N requires N, 2.8%).

Salicylic acid, m.p. 168°, and *p*-nitrohippuric acid, m.p. 135°, were prepared by standard methods. *p*-Aminohippuric acid, m.p. 199°, was purchased.

Animals. Tortoises (*Testudo graeca*) which weighed about 700–900 g. were fed on cabbage and boiled potatoes and were kept over a funnel on wire gauze to collect the milky urine. A 100 w electric lamp suspended at one corner of the enclosure kept the temp. at 30–35°. Water tortoises (*Emys orbicularis*) which weighed about 350 g. were fed on raw fish and kept in an aquarium tank at 25° with a brick island at one end. For metabolic experiments they were transferred to a 12 in. diameter glass-covered dish containing 500 ml. of water which was removed daily, evaporated and examined for metabolites.

Grass snakes (*Natrix natrix*) which weighed about 50–100 g. were fed occasionally on fresh, chopped raw codfish and water and kept in a glass tank at 25° on a layer of gravel with some rough bricks. In metabolic experiments they were placed in covered 5 l. beakers from which any excreta were washed out daily.

Slow worms (*Anguis fragilis*) weighed about 25–50 g. and were treated in the same way as the snakes except that they were fed on small white garden slugs.

Green lizards (*Lacerta viridis*) weighed about 50 g. and were kept in a dry, open, glass tank containing a hollow brick and kept at 25–30° by a suspended electric-light bulb. They were fed on mealworms and water. In metabolic experiments they were kept individually in 5 l. beakers from which any excreta were washed out daily.

Young alligators (*Caiman* sp.) weighed 100–150 g. and were kept in a glass-covered glass tank in about 3 cm. of

water which was kept at 30° by an electric lamp suspended over the surface. A hollow brick was provided near the lamp as an island and cave. They were fed on earthworms. In qualitative metabolic experiments urine was withdrawn from the cloaca by blood pipette. In quantitative experiments, a glass funnel having a 15 cm. diameter wire-gauze grid was set up inside the aquarium tank so that the crocodiles could be kept in a water-saturated atmosphere. Water (2–3 ml.) was given by mouth daily, and with this arrangement urine could be collected for periods up to 7 days.

Administration of compounds. These were given by subcutaneous injection as their sodium salts or as the free acids orally in gelatin capsules.

Paper ionophoresis. Compounds were applied as 2 mm. diameter spots to 8 cm. × 50 cm. strips of Whatman no. 1 paper moistened with electrolyte. These strips were clamped between 45 cm. sheets of plate glass and the ends dipped into reservoirs of electrolyte containing carbon electrodes. A potential difference of 500 v was applied across 45 cm. of paper. When 0.02 N-NaOH, 0.02 N-HCl or Britton & Robinson (1931) buffer (British Drug Houses Ltd.) was used, the current was about 1 mA/cm. width of paper. For identifications the 'unknowns' were run side by side with reference compounds for periods of 1–2 hr. No significance could be attached to the absolute directions of migration since no allowance was made for electro-osmotic effects. Migrations are therefore quoted in Table 2 relative

Table 1. R_f values of some acids and their conjugated derivatives

Whatman no. 1 or no. 4 paper was used; the chromatogram was run till the front had moved about 12–15 in. *A*, Butanol-acetic acid-H₂O (4:1:5); *B*, propanol-NH₃ (sp.gr. 0.88) (7:3); *C*, butan-2-one saturated with water; *D*, butan-2-one-2N-NH₃ (1:1); *E*, butan-2-one-2N-acetic acid-water (200:1:100); *F*, benzene-acetic acid-water (1:1:2); *G*, benzene-butanol-NH₃ (sp.gr. 0.88) (2:5:2). All proportions are by volume.

	Solvent						
	<i>A</i>	<i>B</i>	<i>C</i>	<i>D</i>	<i>E</i>	<i>F</i>	<i>G</i>
Benzoic acid	0.90	0.74	0.95	0.13	—	0.95	0.30
Hippuric acid	0.85	0.69	0.35	0.08	—	0.15	0.20
Ornithuric acid	0.95	0.93	0.50	0.38	—	0.30	0.60
Salicylic acid	0.90	0.81	0.90	0.42	—	0.90	0.40
Salicyluric acid	0.80	0.44	0.25	0.01	—	0.50	0.06
Disalicyloylornithine	0.95	0.59	0.80	0.01	—	0.23	0.19
<i>p</i> -Nitrobenzoic acid	0.95	0.75	0.50	0.37	—	0.80	0.30
<i>p</i> -Nitrohippuric acid	0.85	0.68	0.40	0.16	—	0.15	0.20
Bis- <i>p</i> -nitrobenzoylornithine	0.97	0.86	0.20	0.35	—	0.02	0.45
<i>p</i> -Aminobenzoic acid	0.90	0.32	0.85	—	0.90	0.80	0.05
<i>p</i> -Aminohippuric acid	0.67	0.34	0.10	—	0.26	0.05	0.05
<i>N</i> ⁵ -(<i>p</i> -aminobenzoyl)ornithine	0.30	0.39	0.02	—	0.02	0.05	0.15
<i>N</i> ² -(<i>p</i> -aminobenzoyl)ornithine	0.33	0.25	0.01	—	0.01	0.05	0.05
<i>N</i> ² <i>N</i> ⁵ -Bis-(<i>p</i> -aminobenzoyl)ornithine	0.73	0.47	0.10	—	0.45	0.05	0.05
<i>p</i> -Aminobenzoylglucuronide	0.43	0.31	0.05	—	0.05	0.05	0.05

Table 2. Ionophoretic separation of some *p*-aminobenzoic acid derivatives

The figures represent migration in centimetres, relative to that of *p*-aminobenzoic acid, to anode (+) or cathode (–) in 1 hr. Whatman no. 1 paper, 12 v/cm., was used.

	0.02N-HCl	pH 3*	pH 4*	pH 5*	0.02N-NaOH
<i>p</i> -Aminobenzoic acid	0	0	0	0	0
<i>p</i> -Aminohippuric acid	0	+0.1	+2.2	+0.1	–1.0
<i>p</i> -Aminobenzoylglucuronide	+3.0	+1.6	+2.2	–0.5	–1.5
Bis- <i>p</i> -aminobenzoylornithine	–0.8	0	–0.6	–1.0	–2.5
<i>N</i> ² - <i>p</i> -Aminobenzoylornithine	–2.9	–0.9	+1.0	–3.5	–4.0
<i>N</i> ⁵ - <i>p</i> -Aminobenzoylornithine	–2.0	–0.7	+0.4	–3.0	–2.5

* Britton & Robinson (1931) buffer mixture.

to that of *p*-aminobenzoic acid to show the degree of separations achieved.

Paper chromatography. This was carried out by downward displacement as in earlier papers (e.g. Smith, Smithies & Williams, 1953) and approximate R_F values are quoted in Table 1 to indicate degrees of separation achieved. Identifications were made by running reference compound and unknown on the same paper in at least four of the solvents of Table 1.

Large-scale partition chromatography. When this was carried out on sheets of Whatman no. 3 MM or no. 1 paper the solutions were applied as bands 2 cm. wide across the width of the paper. Partition columns were also used. In the latter, diatomaceous earth (Hiflo Supercel, Johns Manville Co. Ltd.) was rubbed with approximately its own weight of the lower layer of a partition system to give a dry, crumbly powder. This was suspended in the top layer of the solvent mixture and the slurry poured into a glass tube plugged loosely at the lower end with cotton wool. After being allowed to settle, the column was washed with one column-length of upper layer and the lower end stoppered when the solvent reached the level of the filling. Mixtures for separation were dissolved in a little lower solvent layer and stirred with enough Hiflo Supercel to give a dry powder. This mixture was placed on top of the column and development with upper solvent layer started. Volumes of eluate were collected periodically and tested on paper chromatograms to see the progress of the separation.

Detection of compounds on paper. Colour reactions and fluorescence phenomena described previously (Friedler & Smith, 1954) were used. In general the ornithine conjugates behaved in these tests in the same way as the glycine derivatives. Hippuric acids were also detected by the orange or red colours given by the specific reaction of Gaffney, Schreier, Di Ferrante & Altman (1954) and aromatic amino compounds were revealed as yellow spots when sprayed with a mixture of equal volumes of 1% (w/v) ethanolic *p*-dimethylaminobenzaldehyde and 0.5N-HCl. Both N^2 - and N^5 -*p*-aminobenzoylornithines were detected as purple spots by the ninhydrin reagent.

Quantitative measurements. Lizards were kept in covered 5 l. beakers at 30–35° and the excreta washed out periodically into 25 ml. graduated flasks with aq. 0.1N-NH₃.

Alligators were kept as described above, the urine being diluted to 25 ml. Small volumes (usually 0.1–0.2 ml.) were separated on Whatman no. 3 MM paper in solvent *E* and

the zones corresponding to different metabolites eluted chromatographically with water and assayed by the Bratton & Marshall (1939) method (cf. Friedler & Smith, 1954). Calibration curves were constructed for each metabolite, using the Unicam SP 600 spectrophotometer, and recoveries of known amounts of material, after chromatography and elution, ranged from 90 to 100%.

RESULTS

Chromatographic experiments

The acids were administered to the reptiles at a dose level of 0.2 mg./g. and excreta collected as described above. Spots of liquid urine (tortoises, alligators) were chromatographed directly. Solid excreta (snakes, lizards) were stirred to a cream with N-NH₃ and the liquid was filtered before chromatography. Results of chromatographic identifications are summarized in Table 3. In some experiments, where trace amounts could not be identified with certainty, large-scale paper chromatograms were run, appropriate zones were eluted and the concentrated eluates examined on further chromatograms.

The elimination of excretory products was slow and irregular. Diazotizable amines were detectable in snake excreta 6 weeks after dosing with *p*-aminobenzoic acid and in lizards after 3 weeks. Metabolites other than hippuric acids or ornithuric acids were present in the excreta after each aromatic acid had been fed but these were further investigated only in experiments with *p*-aminobenzoic acid (see below).

Isolation of hippuric and ornithuric acid from tortoise urine

A tortoise (*Testudo graeca*) weighing 900 g. was given 50 mg. of benzoic acid by mouth and kept on a funnel for 24 hr. The urine (10 ml.) reduced Benedict solution and gave a positive naphtha-resorcinol reaction. It was evaporated to dryness *in vacuo* and extracted with a little aq. 2N-NH₃.

Table 3. Paper-chromatographic identification of conjugates of aromatic acids in reptile urines

Acid fed	Metabolite detected	Detected in urine of					
		Snake	Green lizard	Slow worm	Alligator	Tortoise	Water tortoise
Benzoic acid	Hippuric acid	Trace	—	—	.	+	+
	Ornithuric acid	+	+	+	.	+	+
Salicylic acid	Salicyluric	.	.	.	.	+	.
	Disalicyloylornithine	.	.	.	.	+	.
<i>p</i> -Nitrobenzoic acid	<i>p</i> -Nitrohippuric acid	.	Trace	.	.	.	.
	Bis- <i>p</i> -nitrobenzoylornithine	.	+	.	.	.	.
	<i>p</i> -Aminobenzoic acid	.	+	.	.	.	+
	Bis- <i>p</i> -aminobenzoylornithine	.	+	.	.	.	+
<i>p</i> -Aminobenzoic acid	<i>p</i> -Aminohippuric acid	Trace	Trace	.	+	+	.
	Bis- <i>p</i> -aminobenzoylornithine	+	+	.	+	+	.
	N^5 - <i>p</i> -Aminobenzoylornithine	+	+	.	+	+	.
	<i>p</i> -Aminobenzoylglucuronide	+	+	.	+	+	.

Table 4. *Excretion of p-aminobenzoic acid and conjugates in Lacerta viridis and Caiman sp.*

	Percentage of dose excreted as <i>p</i> -amino-								
	-ornithuric acid			-hippuric acid			-benzoic acid		
	<i>a</i>	<i>b</i>	<i>c</i>	<i>a</i>	<i>b</i>	<i>c</i>	<i>a</i>	<i>b</i>	<i>c</i>
<i>Lacerta viridis</i>									
Days 1 and 2	9.2	3.1	9.8	.	.	.	3.4	1.0	2.6
Days 3 and 4	11.9	14.3	3.1	.	.	.	3.9	2.2	1.7
Days 5 and 6*	1.4	9.8	2.2	.	.	.	1.1	1.3	0.6
Totals	22.5	27.2	15.1	.	.	.	8.4	4.5	4.9
<i>Caiman sp.</i>									
Days 1-6*	11.8	17.1	.	3.5	9.3	.	4.4	5.5	.

* Excretion still incomplete.

After insoluble material had been filtered off the solution was applied as a strip 2 cm. $\times$ 30 cm. on Whatman no. 1 paper and separated in solvent *G*. Bands which quenched the paper fluorescence, corresponding to the R_f values of hippuric acid and ornithuric acid, were cut out and eluted chromatographically with water. Acidification of the ornithuric acid eluate gave a precipitate which was crystallized from water (13 mg.). This had m.p. 184°, not depressed by mixture with authentic L-ornithuric acid, and $[\alpha]_D^{25}$ 9.3 in 0.1N-NaOH (*c*, 1.5) (Found: C, 67.2; H, 5.9; N, 8.0. $C_{19}H_{20}O_4N_2$ requires C, 67.0; H, 5.9; N, 8.2%). In another experiment in which 200 mg. of sodium benzoate was injected subcutaneously, 18 mg. of L-ornithuric acid was isolated from the first day's urine.

The hippuric acid zone on a large-scale paper chromatogram in solvent *G* was eluted with water and rechromatographed in solvent *F*. The eluate of the zone giving the colour reactions of Gaffney *et al.* (1954) gave a very small precipitate on acidification, which after crystallization from water had m.p. 188° (0.2 mg.), not depressed by mixing with hippuric acid.

In a similar experiment with the water tortoise (*Emys*) 100 mg. of benzoic acid was given orally and the first day's excretion examined. L-Ornithuric acid (1 mg.), m.p. and mixed m.p. 183°, and benzoic acid (3 mg.), m.p. and mixed m.p. 122°, were isolated.

Conjugation of *p*-aminobenzoic acid

Quantitative experiments. The acid was given orally to lizards and alligators at a dose level of 0.5 mg./g. The results are summarized in Table 4. *p*-Aminohippuric acid was present only in traces in lizard excreta and was not estimated. No evidence of any significant amount of acetylation was found in any of the reptiles in any qualitative experiment, and acid hydrolysis of urines in the quantitative

experiments gave no increase of diazotizable amine.

The excreta of the reptiles in these quantitative experiments was usually entirely urine but in some experiments with alligators this was contaminated with faeces. In these experiments the diazotizable amino compound present was almost entirely *p*-aminobenzoic acid. Qualitative experiments showed that alligator faeces, which appeared to consist mainly of the earth from the earthworm diet, enzymically hydrolysed bis-*p*-aminobenzoyl-ornithine to *p*-aminobenzoic acid; a small amount of *N*⁵-*p*-aminobenzoylornithine was also produced. Prolonged exposure of the urine to the faeces resulted in the disappearance of any diazotizable amine. Experiments with obvious faecal contamination were therefore rejected.

Identification of ornithine conjugates. Six green lizards were each given 25 mg. of *p*-aminobenzoic acid by mouth and the excreta were collected for 6 days. The excreta were roughly ground and extracted with 20 ml. of boiling water for 30 sec. The filtrate was evaporated to dryness *in vacuo* and the residue separated on a partition column 2 cm. diam. $\times$ 50 cm. with solvent system *E*. Eluates containing material behaving like *p*-aminobenzoic acid and bis-*p*-aminobenzoylornithine on test paper chromatograms were collected.

The eluates containing bis-*p*-aminobenzoyl-ornithine were evaporated to small bulk and further purified by large-scale chromatography in solvent system *A*. Insufficient material was obtained for preparing solid derivatives but the product behaved identically with bis-*p*-aminobenzoylornithine in each of the solvent systems of Table 1 and at each of the pH values of Table 2 in ionophoresis experiments. Hydrolysis with N-HCl at 100° for 4 hr. gave a mixture containing *p*-aminobenzoic acid, ornithine and *N*²- and *N*⁵-*p*-aminobenzoylornithines, which were detected chromatographically.

Material remaining on the original partition column was eluted with 500 ml. of water. After evaporation, the residue gave an intense naphtharesorcinol reaction and reduced Benedict solution. It was separated by large-scale chromatography on paper in solvent *E* into two fractions *X* and *Y*. The slower-moving one (*Y*) was eluted and was identified as *N*⁵-*p*-aminobenzoylornithine by paper chromatography and ionophoretic experiments. No *N*²-*p*-aminobenzoylornithine was found.

In other experiments, bis-*p*-aminobenzoylornithine and small amounts of *N*⁵-*p*-aminobenzoylornithine were separated and identified in the excreta of *Natrix*, *Testudo* and *Caiman* by the methods described above.

Identification of *p*-aminobenzoylglucuronide. The faster-moving band (*X*) on the large-scale paper chromatogram in solvent *E* was eluted and behaved identically with *p*-aminobenzoylglucuronide when chromatographed in the solvent systems of Table 1. It also migrated at the same speed as the glucuronide in the ionophoretic systems of Table 2.

The eluate, which gave an intense naphtharesorcinol reaction, was hydrolysed with locust-crop fluid (which contains a β -glucuronidase) or by boiling for 1 min. with *N*-NaOH, to give *p*-aminobenzoic acid and glucuronic acid, which were identified on paper chromatograms. Quantitative naphtharesorcinol estimations of glucuronic acid (see Mead, Smith & Williams, 1958) gave values equivalent to a total of 14 mg. of *p*-aminobenzoylglucuronide in this eluate. The same value was also obtained by the Bratton & Marshall (1939) diazo estimation by using a calibration curve prepared from authentic *p*-aminobenzoylglucuronide. The eluate had an absorption maximum at 270 m μ in 0.1*N*-NaOH, with an intensity corresponding to the presence of 15.4 mg. of *p*-aminobenzoylglucuronide, if this was assumed to be the only absorbing material present.

In other similar experiments *p*-aminobenzoylglucuronide was identified in the excreta of *Natrix*, *Testudo* and *Caiman*. Identification was made, as above, by chromatography, ionophoresis and hydrolysis to *p*-aminobenzoic acid and glucuronic acid.

***p*-Aminohippuric acid.** Only traces of *p*-aminohippuric acid were detected in the excreta of the lizards and snakes used. *Testudo* and *Caiman* excreted a greater proportion of this conjugate and excreta from these species were separated in large-scale paper chromatograms in solvent *E*. The band corresponding to *p*-aminohippuric acid was eluted and shown to contain material behaving identically with the reference compound in all the solvent systems used for chromatography and at each of the pH values used for ionophoresis. Hydrolysis of

the eluate with *N*-HCl for 2 hr. at 100° gave *p*-aminobenzoic acid and glycine, which were identified on paper chromatograms (cf. Friedler & Smith, 1954).

DISCUSSION

Except for the data on reptiles quoted above, evidence for the existence of an ornithine conjugation is confined to birds. Information about birds is almost entirely concerned with the hen, which is well known to form ornithuric acids from ingested aromatic acids. Experiments with enzymes suggest that ducks and pigeons may also be capable of this conjugation (Schachter, Manis & Taggart, 1955; Efimochkina, 1951). Small amounts of a glycine conjugate may occur in the hen (Suga, 1919) and *N*²- and *N*⁵-substituted ornithines have been detected by Wu Chang & Johnson (1957) after nicotinamide had been fed to chicks. Similarly, small amounts of *N*⁵-substituted ornithine and a glycine conjugate have been found in the reptile experiments.

All orders of terrestrial vertebrates which have been investigated, including reptiles and birds, possess to some degree the ability to conjugate aromatic acids with glycine and glucuronic acid. It is only in reptiles and birds where the additional reaction with ornithine is possible, and it is of interest to consider the evolutionary relationship of these two groups to the rest of the vertebrates (Fig. 1). Present-day birds and reptiles are thought to have evolved from a different group of primitive reptiles from the reptilian ancestors of the modern mammals (see Young, 1950). Since amphibia, which branch off from the evolutionary tree at an earlier stage, use only glycine and glucuronic acid conjugations it seems possible that

Fig. 1. Evolutionary relationships.

the ability to use ornithine in detoxication arose about 200 million years ago, when the cotylosaurian reptiles were becoming diversified into the ancestors of the later reptile orders.

There appears to be a relation between the presence of the ornithine conjugation and the excretion of waste nitrogen as uric acid. It may be significant that in the developing hen's egg the ornithine conjugation appears at the same time as the change to full uricotelic metabolism (Takahashi, 1928). Hens, the grass snake and the two species of lizard studied are all uricotelic animals (see Prosser, 1950, for summary) and it is in these species that the ornithine detoxication predominates over the formation of hippuric acids. The alligators and both aquatic (*Emys*) and terrestrial (*Testudo*) tortoises are partly uricotelic and partly ureotelic. In these species the proportion of hippuric acids formed is much higher (e.g. Table 4) though the ornithuric acid is still the major metabolite.

Florkin (1949) has suggested that the mixed nitrogen metabolism in ancient orders like tortoises and crocodiles is a reflexion of their primitive nature, and he distinguishes them from the more recently evolved lizards and snakes. It is possible that the retention of a glycine-conjugating mechanism alongside the main ornithine detoxication is a similar relic of their amphibian ancestry.

The coincidence of uricotelism with an ornithine conjugation need not necessarily indicate any direct metabolic connexion between the two processes. The advantage of uric acid to organisms reproducing via a cleidoic egg has frequently been pointed out (e.g. Baldwin, 1948), but this in its turn may have introduced another biochemical problem. In the synthesis of uric acid by bird liver, four of the carbon atoms arise directly or indirectly from glycine, and the excretion of the daily intake of nitrogen must create a greater need for glycine in uricotelic than in ureotelic animals. It seems reasonable to believe that in the early stages of evolution of the reptilian ancestors of the birds the glycine-sparing action of an ornithine conjugation might have been an evolutionary asset.

Although ornithine conjugation has been found, so far, only in association with a uricotelic nitrogen metabolism, the converse is not true. Insects are primarily uricotelic but those species for which results are available form hippuric acids from aromatic acids (Smith, 1955a). The mechanism of the synthesis of uric acid in insects is not known; but it is of interest that, of the few insects whose requirements are known, two (*Aedes* and *Blattella*) require glycine as an essential amino acid (Levinson, 1955).

SUMMARY

1. Ornithuric acids have been detected as the major amino acid conjugate in the excreta of six species of reptile after feeding aromatic acids.

2. L(+)-Ornithuric acid has been isolated from excreta of *Testudo graeca* and *Emys orbicularis* dosed with benzoic acid.

3. After *p*-aminobenzoic acid was fed to *Natrix natrix*, *Lacerta viridis*, *Testudo graeca* and *Caiman* sp., small amounts of *p*-aminobenzoyleglucuronide were detected in the excreta.

4. After feeding *p*-aminobenzoic acid, traces of *p*-aminohippuric acid were found in the excreta of *Natrix* and *Lacerta* and large amounts in the excreta of *Caiman* and *Testudo*.

I am grateful to Dr A. d'A. Bellairs of St Mary's Hospital Medical School for advice on keeping reptiles, and to Professor R. T. Williams for his interest in this work.

REFERENCES

- Baldwin, E. (1948). *Comparative Biochemistry*, 3rd ed., p. 59. Cambridge: University Press.
- Bratton, A. C. & Marshall, E. K. (1939). *J. biol. Chem.* **128**, 537.
- Bray, H. G., Lake, H. J., Neale, F. C. & Thorpe, W. V. (1948). *Biochem. J.* **42**, 434.
- Britton, H. T. S. & Robinson, R. A. (1931). *J. chem. Soc.* p. 1456.
- Bunge, G. & Schmiedeberg, O. (1876). *Arch. exp. Path. Pharmac.* **6**, 233.
- Efimochkina, E. F. (1951). *C.R. Acad. Sci. U.R.S.S.* **80**, 793.
- Florkin, M. (1949). *Biochemical Evolution*, p. 42. New York: Academic Press.
- Friedler, L. & Smith, J. N. (1954). *Biochem. J.* **57**, 396.
- Gaffney, G. W., Schreier, K., Di Ferrante, N. & Altman, K. I. (1954). *J. biol. Chem.* **206**, 695.
- Harris, J. I. & Work, T. S. (1950). *Biochem. J.* **46**, 582.
- Hiratani, T. (1957). *Osaka Daigaku Igaku Zasshi*, **9**, 317. Quoted in *Chem. Abstr.* **51**, 14138.
- Hunter, A. (1939). *Biochem. J.* **33**, 27.
- Jaffe, M. (1877). *Ber. dtsch. chem. Ges.* **10**, 1925.
- Kamil, I. A., Smith, J. N. & Williams, R. T. (1951). *Biochem. J.* **50**, 235.
- Khalil, F. (1947). *J. biol. Chem.* **171**, 611.
- Khalil, F. (1948a). *J. biol. Chem.* **172**, 101.
- Khalil, F. (1948b). *J. biol. Chem.* **172**, 105.
- Khalil, F. (1951). *J. biol. Chem.* **189**, 443.
- Komori, Y., Sendju, Y., Sagara, J. & Takamatsu, M. (1926). *J. Biochem., Tokyo*, **6**, 21.
- Levinson, Z. H. (1955). *Riv. Parassit.* **16**, 1.
- McGilvery, R. W. & Cohen, P. P. (1950). *J. biol. Chem.* **183**, 179.
- Mead, T., Smith, J. N. & Williams, R. T. (1958). *Biochem. J.* **68**, 61.
- Prosser, C. L. (1950). *Comparative Animal Physiology*. London: W. B. Saunders.
- Rothstein, M. & Greenberg, D. M. (1957). *Arch. Biochem. Biophys.* **68**, 206.

- Sahyun, M. (1948). *Outline of Amino Acids and Proteins*, 2nd ed., p. 168. New York: Reinhold Publishing Corporation.
- Schachter, D., Manis, J. G. & Taggart, J. V. (1955). *Amer. J. Physiol.* **182**, 537.
- Schiff, H. (1859). *Liebigs Ann.* **111**, 368.
- Smith, J. N. (1955a). *Biol. Rev.* **30**, 455.
- Smith, J. N. (1955b). *Biochem. J.* **60**, 436.
- Smith, J. N., Smithies, R. H. & Williams, R. T. (1953). *Biochem. J.* **54**, 225.
- Suga, T. (1919). *Kyoto med. J.* **15**, 225. Quoted in *Chem. Abstr.* (1921). **15**, 881.
- Takahashi, M. (1928). *Hoppe-Seyl. Z.* **178**, 294.
- Williams, R. T. (1947). *Detoxication Mechanisms*, p. 95. London: Chapman and Hall.
- Wu Chang, M. L. & Johnson, B. C. (1957). *J. biol. Chem.* **226**, 799.
- Young, J. Z. (1950). *The Life of Vertebrates*, p. 368. Oxford: Clarendon Press.

Countercurrent Distribution: Quantitative Analysis of Results, Especially where Separation is Incomplete

By A. A. LEVI

Research Department, Imperial Chemical Industries Pharmaceuticals Division, Hexagon House, Blackley, Manchester 9

(Received 26 September 1957)

A method has been described (Goodall, 1954) for calculating the amounts of each component in a mixture of two substances when separation in, say, a Craig machine, is incomplete. Pairs of tubes containing reasonable amounts of both substances are selected, and the amounts of the components found by solving sets of simultaneous equations. The possibility of extending the method to more than two components was also pointed out.

By an adaptation of standard statistical techniques it is now shown to be possible to calculate not only the composition of a mixture, but also the values of the partition ratios of the components if these are unknown, even when quite a small degree of separation has been achieved by countercurrent distribution. All the relevant tubes are used in one operation.

Apart from its obvious value in enabling information to be obtained in general countercurrent-distribution experiments, the method might occasionally have useful analytical applications. Thus for a pair of acids difficult to separate or to determine in the presence of each other, a system might be found for which a partial separation could be obtained in a relatively small number of countercurrent transfers. From a knowledge of the distribution ratio of each pure substance, and a measurement of the total acidity in each tube, the amount of each acid can be found. Examples are described below.

Bacher (1951) has described a simpler method from a different viewpoint for calculating partition ratios, by using all the relevant tubes in a zone, and has extended the method to the analysis of mixtures. The latter analysis, however, requires more than one measurement (e.g. optical density

at different wavelengths) on each tube. Bland, Hillis & Williams (1952) also discuss this method of finding partition ratios. These procedures, although less elaborate, are of less general application than those proposed here.

THEORETICAL

As is well known, when A units of a substance of partition ratio K_A have been submitted to n transfers in a countercurrent apparatus of the Craig type the amount of substance, Y_r , in tube r is given by the expression $Y_r = Aa_r$, where a_r is the appropriate term of the binomial distribution, which can be calculated (Williamson & Craig, 1947) or found from suitable tables (Eisenhart, 1952) when K_A and n are known. There will be a series of such equations, one for each tube. If Y_r is measured experimentally and A is unknown then a_r can be treated as an independent variable and Y_r as a dependent variable in the series of equations. The best estimate of A , when all the tubes containing measurable amounts of substance are used, is then found by the method of least squares. If A were known the true value of Y_r would be Aa_r , and the difference between this value and the experimental value of Y_r is $Y_r - Aa_r$. The principle of least squares states that the best estimate of A from the experimental data is the value which makes the sum of the squares of these differences, $\Sigma(Y_r - Aa_r)^2$, a minimum. By the usual process of differentiating and equating to zero we then get (removing the subscript r)

$$A = \Sigma Y_r a_r / \Sigma a_r^2. \quad (1)$$

One can also calculate a sum of squares, R , representing the deviations of the experimental