Optimum deamination of threonine proceeds over the range pH 9.0–10.0. This includes the optimum pH of 9.3 for the *Neurospora* enzyme (Yanofsky & Reissig, 1953). Sayre & Greenberg (1956) reported that the optimum pH for the sheep-liver enzyme was 7.4.

L-Serine appears to be deaminated in LC1 by the same enzyme which acts upon L-threonine, a finding in agreement with the work by Umbarger & Brown (1957) on the *E. coli* enzymes and Yanofsky & Reissig (1953) with *Neurospora*. In contrast, Sayre & Greenberg (1956) demonstrated that two easily separated enzymes were present in sheep liver: one specific for L-serine and the other for L-threonine.

The K_m values for L-threonine and L-serine were 3.0 and 6.8 mM respectively. These figures are similar to those found by Yanofsky & Reissig (1953) for the *Neurospora* enzyme, and by Wood & Gunsalus (1949) for the deamination of L-threonine by *E. coli* extracts. The sheep-liver enzyme has a lower affinity for its substrate ($K_m = 29$ mM; Sayre & Greenberg, 1956).

SUMMARY

1. The L-threonine deaminase of the rumen micro-organism LC1 was purified over 100-fold from extracts of vacuum-dried cells by successive fractionation with protamine sulphate and ammonium sulphate, gradient elution with ammonium sulphate and fractionation on a calcium phosphate gel-cellulose column.

2. The enzyme has an optimum pH in the range 9.0–10.0. K_m values of 6.8 mM and 3.0 mM were obtained for L-serine and L-threonine respectively. D-Threonine appeared to inhibit the enzyme.

3. The ratio of L-serine deaminase activity to L-threonine deaminase activity varied little at different stages of the purification, suggesting that one enzyme attacks both substrates.

4. The purified enzyme, after freeze-drying in the presence of ammonium sulphate or glutathione, required pyridoxal phosphate. Glutathione also appeared to be necessary for maximum activity.

I wish to express my thanks to Dr S. R. Elsdon for his continued interest and advice and to Dr J. L. Peel for permission to publish his method for large-scale culture of LC. This work was supported in part by the Rockefeller Foundation. I also wish to acknowledge the receipt of a Research Studentship from the Agricultural Research Council.

REFERENCES

- Conway, E. J. & O'Malley, E. (1942). *Biochem. J.* **38**, 655.
 Dixon, M. (1953). *Biochem. J.* **54**, 457.
 Elsdon, S. R. & Lewis, D. (1953). *Biochem. J.* **55**, 183.
 Elsdon, S. R., Volcani, B. E., Gilchrist, F. M. C. & Lewis, D. (1956). *J. Bact.* **72**, 681.
 Friedemann, T. E. & Haugen, G. E. (1943). *J. biol. Chem.* **147**, 415.
 Keilin, D. & Hartree, E. F. (1936). *Proc. Roy. Soc. B*, **119**, 114.
 Lewis, D. & Elsdon, S. R. (1955). *Biochem. J.* **60**, 683.
 Lineweaver, H. & Burk, D. (1934). *J. Amer. chem. Soc.* **56**, 658.
 McIntire, W. M., Palmer, G. & Marshall, H. L. (1945). *Biochem. J.* **48**, 171.
 Metzler, D. E. & Snell, E. E. (1952). *J. biol. Chem.* **198**, 363.
 Roberts, E. (1952). *J. biol. Chem.* **198**, 495.
 Robertson, W. B. (1942). *Science*, **96**, 93.
 Sayre, F. W. & Greenberg, D. M. (1956). *J. biol. Chem.* **220**, 787.
 Schwimmer, S. (1953). *Nature, Lond.*, **171**, 443.
 Shinn, L. A. & Nicolet, B. H. (1941). *J. biol. Chem.* **138**, 91.
 Stadtman, E. R. & Barker, H. A. (1949). *J. biol. Chem.* **180**, 1085.
 Stadtman, E. R., Novelli, G. D. & Lipmann, F. (1951). *J. biol. Chem.* **191**, 365.
 Swingle, S. M. & Tiselius, A. (1951). *Biochem. J.* **48**, 171.
 Umbarger, H. E. & Brown, B. (1956). *J. Bact.* **71**, 443.
 Umbarger, H. E. & Brown, B. (1957). *J. Bact.* **73**, 105.
 Umbreit, W. W., Burris, R. H. & Stauffer, J. F. (1949). In *Manometric Techniques and Tissue Metabolism*, p. 173. Minneapolis: Burgess Publishing Co.
 Walker, D. J. (1957). *Biochem. J.* **67**, 4P.
 Wood, W. A. & Gunsalus, I. C. (1949). *J. biol. Chem.* **181**, 171.
 Yanofsky, C. & Reissig, J. L. (1953). *J. biol. Chem.* **202**, 567.

Condensed Tannins

1. A STUDY OF COMPLEX LEUCO-ANTHOCYANINS PRESENT IN CONDENSED TANNINS

By D. G. ROUX AND S. R. EVELYN

Leather Industries Research Institute, Rhodes University, Grahamstown, South Africa

(Received 16 December 1957)

Condensed tannins are widely distributed in Nature, and constitute a heterogeneous group possessing the common characteristic of forming insoluble 'phlobaphenes' on treatment with hot dilute mineral acids in aqueous medium. This rather general characteristic is shared by the

catechins, which occur in high proportion in association with tannins in some commercial tanning extracts, e.g. cube gambier from the leaves of *Uncaria gambir* and cutch from the heartwood of *Acacia catechu*. Freudenberg & Maitland (1934) and Freudenberg & Weinges (1955)

considered the highly complex condensed tannins to be polymers of catechins, a view also supported by Hathway & Seakins (1957). So far it has not been possible, however, to obtain direct structural evidence of the $C_6-C_3-C_6$ nature of the hypothetical basic units present in naturally occurring condensed tannins, although it is readily conceded that the evidence provided specifically for gambier and cutch extracts appears to be conclusive.

The remaining noteworthy concept is the leuco-anthocyanin hypothesis of Bate-Smith & Swain (1953), who regarded leuco-anthocyanins as well as catechins as possible prototypes of condensed tannins, from a study of the absorption spectra and phlobaphene reaction of these substances. Plant extracts which are classed as 'condensed tannins' and widely used for commercial tannage were shown by Pigman, Anderson, Fischer, Buchanan & Browning (1953) to be rich in leuco-anthocyanins. The extracts examined include those from black-wattle bark (*Acacia mollissima*), quebracho heartwood (*Schinopsis balansae* and *S. quebracho-colorado*), mangrove bark (mainly *Rhizophora* spp.), western hemlock wood (*Tsuga heterophylla*) and the bark of the black spruce (*Picea mariana*). The scarlet colorations developed from black-wattle and quebracho extracts were shown to exhibit lower absorption maxima (532 and 525 $m\mu$ respectively) than those obtained from other extracts (540–545 $m\mu$). Hillis (1956*a, b*) found that 'condensed tannins' suitable for tanning purposes, and rich in leuco-anthocyanins, are also located in the heartwoods of various *Eucalyptus* species and in the bark of *Pinus radiata*.

King & Bottomley (1954) and King & Clark-Lewis (1955) linked the structure of the monomeric leuco-anthocyanin, melacacidin, with that of the condensed tannins. They, however, overlooked the fact that monomeric flavan-3:4-diols are in-

capable of tanning collagen, and provided no evidence of the presence of complex leuco-anthocyanins in those condensed tannins used for commercial tannage.

The present work shows that both wattle and quebracho extracts contain appreciable proportions of monomeric and polymeric leuco-anthocyanins of the same $C_6-C_3-C_6$ structural pattern, and with varying affinity for cellulose and collagen in aqueous medium. The flavan-3:4-diol structure thus appears to form the basis for at least one type of condensed tannin, i.e. that based on leuco-anthocyanins. Anthocyanidins resulting from wattle and quebracho extracts belong to a new group compared with the naturally occurring delphinidin, cyanidin and pelargonidin series studied by Bate-Smith (1954). This has necessitated a more detailed study of chromatographic identification methods for anthocyanidins and for simple and complex leuco-anthocyanins (Roux, 1957*a, b*), as well as a detailed study of the molecular-weight distribution of wattle and quebracho tannins.

EXPERIMENTAL

Synthesis and origin of anthocyanidins. Robinetinidin chloride or 3:7:3':4':5'-pentahydroxyflavylium chloride (I, $R=H$), fisetinidin chloride or 3:7:3':4'-tetrahydroxyflavylium chloride (II, $R=H$) and 3:7:4'-trihydroxyflavylium chloride (III, $R=H$) were synthesized as previously described (Roux, 1957*a*).

Other anthocyanidins were obtained from natural sources. 3:7:8:3':4'-Pentahydroxyflavylium chloride was obtained by generation from melacacidin (King & Bottomley, 1954), separated from the heartwood extractives of *Acacia melanoxylon* by using the 'preparative' method. Water containing 2% of acetic acid was used as irrigant for thick cellulose sheets (Whatman no. 3 or 3MM) and the

melacacidin was eluted with 70% ethanol after chromatography. Contaminants were removed by repeating the process with butanol-acetic acid-water (4:1:5, upper phase) as irrigant. The anthocyanidin was generated from melacacidin by the method of Pigman *et al.* (1953).

Anthocyanidins containing the phloroglucinol nucleus were generated from naturally occurring anthocyanins present in flowers by hydrolysis with 3N-HCl on the water bath. The aqueous solutions were extracted with amyl alcohol to furnish a concentrate of the anthocyanidin (Bate-Smith, 1954). Delphinidin (I, R=OH) was obtained from *Delphinium* and *Salvia* spp.; cyanidin (II, R=OH) from the rose *Daily Mail* (scented), and pelargonidin (III, R=OH) from various *Pelargonium* spp.

Chromatographic identification of anthocyanidins. Chromatograms were run on Whatman no. 1 paper in two solvent systems: (a) the 'Forestal Solvent', namely water-acetic acid-conc. HCl (10:30:3, v/v) and (b) 90% (w/v) formic acid-3N-HCl (1:1, v/v) at 21°. After noting the R_F values the absorption maxima of the anthocyanidins were determined directly on the paper by the method of Bradfield & Flood (1953). Finally, after allowing the escape of excess of HCl from the paper by airing for 4-7 days at room temperature (20°), the anthocyanidin spots were treated with a few drops of 5% aluminium sulphate solution. The paper was allowed to dry in air and the absorption maximum of the blue colorations developed were determined in the spectrophotometer as before. Visual comparison and distinction between the various shades of blue could be made by comparison with reference compounds.

The above procedures are complicated by the presence of 'background' polyphenols, particularly where anthocyanidins are generated from condensed tannin mixtures. In such instances the pigment generated was applied in an intense streak on the origin of the paper chromatogram. After development, preferably in 90% formic acid-3N-HCl mixture, which tends to keep the highly condensed tannins at the origin, the scarlet band was cut out as a strip. The strip was cut and developed on the wick principle (Reith, 1957) with methanol. Tannins and other phenolic compounds tend to move forward while the anthocyanidins lag behind. The strip is removed once the methanol front nears the top of the tube and the paper is air-dried. The absorption spectrum of the anthocyanidin is investigated as above with the spectrophotometer. The trailing portion of the scarlet coloration was found to give the most satisfactory measurements of absorption maxima where other contaminants are present.

Generation of anthocyanidins from leuco-anthocyanins. The method of Pigman *et al.* (1953) was used and the scarlet colorations developed were examined qualitatively by means of paper chromatography, or used for estimating the yield of anthocyanidin quantitatively. Where the rate of generation of anthocyanidins from leuco-anthocyanins or the stability of the anthocyanidins were studied a number of similar tubes were used, and these were removed from the boiling-water bath at regular intervals and cooled quickly.

Origin and synthesis of monomeric leuco-anthocyanins or flavan-3:4-diols. Melacacidin, 7:8:3':4'-tetrahydroxyflavan-3:4-diol (King & Bottomley, 1954) was obtained from the methanolic extract of the heartwood of *Acacia melanoxylon* by the preparative method with chromatographic sheets, as previously outlined.

7:3':4':5'-Tetrahydroxyflavan-3:4-diol, leucorobinetidin, and 7:3':4'-trihydroxyflavan-3:4-diol, leucofisetinidin, were obtained by the catalytic hydrogenation of dihydro-robinetin and dihydrofisetin (fustin) respectively with platinum oxide as catalyst by the method of Freudenberg & Roux (1954).

5:7:3':4'-Tetrahydroxyflavan-3:4-diol, leucocyanidin, was obtained by the reduction of dihydroquercetin (taxifolin) with sodium borohydride by the method of Swain (1954).

Identification of leuco-anthocyanins on paper chromatograms. Toluene-*p*-sulphonic acid (3-5%, w/v) in absolute ethanol is evenly sprayed on two-way paper chromatograms. After heating the paper at 80-90° in an air oven, red spots developed in those areas in which leuco-anthocyanins are present. During the heating period (5-10 min.) the development of the pink area must be watched, as prolonged heating causes the cellulose to darken, obscuring the colours which have developed. The sheets may be rolled into a tube and heated in an oven, provided that the chromatographic paper rests on thick absorbent paper and remains out of contact with the hot metal frame of the oven. Where this procedure is adopted the paper must be removed at 5 min. intervals for inspection. The paper is quite stable after development of the spots and may be stored for record purposes.

Fractionation of black-wattle tannins for study of leuco-anthocyanins. Black-wattle tannins, when examined by two-way paper chromatography, may be arbitrarily subdivided into a more mobile fraction which travels mainly as discrete spots A-F, and a low R_F trail in which no clear differentiation of constituents is apparent (Fig. 1).

Constituents of the mobile higher R_F fraction were separated on a micro-scale by paper chromatography on thick cellulose sheets (Whatman no. 3 or 3MM, 18½ × 22½ in) by the 'preparative' method, as outlined by Nordström & Swain (1953) and others. Fresh wattle bark or commercial extract were exhaustively extracted with methanol in order to eliminate the gums naturally present. The methanolic solution was streaked on to the sheets and the chromatogram developed in butan-1-ol-acetic acid-water (6:1:2). By using traces of the brilliantly fluorescing flavanols naturally present (e.g. robinetin, R_F 0.51, and

Fig. 1. Two-dimensional chromatogram of black-wattle tannins, indicating relative positions of the main mobile 'constituents' and the low R_F trail.

Table 1. *Preliminary fractionation of black-wattle tannins*

Fraction	Extraction period	Polyphenolic tannins* (%)	Percentage of total polyphenols	Mol. wt.
C_1	24 hr.	82.6	49.5	1290
C_2	72 hr.	81.9	13.3	1305
C_3	24 hr.	45.5	26.2	1507
C_4	8 days	39.8	5.3	1556
C_5	14 days	38.7	3.2	1540

* Estimation by the ultraviolet method (Roux, 1951).

fisetin, R_f 0.72) as reference compounds, bands corresponding to constituents *D* and *E*, *B* and *C*, *A*, A_1 , A_2 and *F*, and *G* and *H* were cut. The various groups of constituents were eluted with ethanol-water mixtures as described by Nordström & Swain (1953).

The constituents were separated by re-running the bands on paper sheets with 2% acetic acid as developing medium. The constituents were eluted as before and concentrated. Overlap of constituents occurred to a large extent, but by re-running the bands and by cutting and discarding strips between bands, good separation was achieved and the 'constituents', *A*, A_1 , *B*, *C*, *D*, *E*, *G* and *H* were isolated in a form sufficiently 'pure' for their migration as discrete spots on two-way paper chromatograms when the above solvents were used. These constituents were, in fact, still not completely pure, as shown by further chromatography with water-saturated phenol as irrigant, and it must be assumed that overlap of constituents occurs even after using two solvent systems. The fractions or 'constituents' obtained by these methods were, however, sufficiently 'pure' for a study of their leuco-anthocyanin reactions.

Estimation of molecular weight of wattle and quebracho tannins

Materials. Owing to the complexity of the mixture wattle tannins were fractionated first by successive solvent extraction of the fresh mature bark, initially with ethyl acetate and finally with methanol (Table 1). Fractions C_1 and C_2 were obtained by successive extractions of macerated fresh bark with ethyl acetate, followed by three successive extractions with methanol, giving fractions C_3 , C_4 and C_5 . Fractions C_1 and C_2 are similar in tannin:non-tannin ratio and in molecular weight (1300), as were also fractions C_3 , C_4 and C_5 (1500–1550). Fractions C_1 and C_3 were selected as representative fractions for detailed examination.

Quebracho extract ('ordinary' or 'unsulphited') from commercial sources (71.6% tannin content) was used without preliminary separation.

Chromatographic fractionation of tannins. The polyphenolic tannins were fractionated by paper chromatography on thick sheets, e.g. Whatman no. 3 or 3MM by the methods previously described (Nordström & Swain, 1953), with 2% acetic acid as irrigant. The presence of acetic acid limits atmospheric oxidation during paper chromatography. After air-drying, strips were cut from the sheet corresponding to R_f values 0, 0.1, 0.2, 0.3, 0.4 and 0.5 + 0.6, so that the centre of each band corresponds to these R_f values. The bands (approx. 2 in. wide) were eluted with ethanol-water mixtures as previously described, and the eluent from each band concentrated under vacuum.

Sugars present in the extract migrate just behind the

solvent front in the chromatographic separation and are therefore eliminated.

Preparation of methoxyl derivatives of tannins. The fractions were methylated with excess of diazomethane (8 g. of nitrosomethylurea to 0.5 g. of tannins) in methanolic solution maintained at -10° for 48 hr. Diazomethane reacts selectively with polyphenolic tannins and permits separation of any residual carbohydrate material during recovery of the product by pouring into water (Evelyn, 1954a). Substituted tannins were vacuum-dried at 110° over P_2O_5 .

Analysis for methoxyl content. Methoxyl values of the fractions were determined by duplicate semi-micro Zeisel analysis, the bromine-acetate absorbent of Vieboch & Brecher (1930) being used.

Estimations of molecular weights. Molecular weights of the methylated tannins were determined in benzene with the Ray (1952) ebulliometer, as modified by Evelyn (1954b) and incorporating improvements detailed by Smith (1956). A further modification of the same apparatus (Evelyn, van Berge & Cooper, 1958) was used to determine molecular weights in chloroform or acetone for those methylated tannin fractions which are only partly soluble in benzene.

The apparatus constant was determined daily and the average value used for calculation of the molecular weights. Average molecular weights were obtained from the mean of readings at four concentrations, as the readings were independent of concentration.

Molecular-weight values of tannins. Results are given as the molecular weights of the unsubstituted polyphenolic tannins, and are calculated from the determined molecular weights of the methylated derivatives and their methoxyl values.

RESULTS

Identification of anthocyanidins

The R_f values of the anthocyanidins in the different solvent systems have been tabulated (Roux, 1957a). 3:7:4'-Trihydroxyflavylium chloride was erroneously represented as having R_f 0 in 90% formic acid-3N-HCl (1:1) mixture, owing to the presence of impurities resulting from synthesis. Subsequent work has shown the correct value to be R_f 0.6, and the anthocyanidin exhibits yellowish fluorescence under ultraviolet light. The absorption maxima of the anthocyanidins and their aluminium complexes (Roux, 1957b) provide additional parameters for their identification. The absorption

maxima of the aluminium complexes tend to be broad, and visual comparison with reference compounds provides useful confirmation of identity.

Differentiation between monomeric and polymeric leuco-anthocyanins

The flavan-3:4-diols were examined by two-way paper chromatography with butanol-acetic acid-water (4:1:5, upper phase) for direction 1 and 2% acetic acid for direction 2. The compounds were located on the chromatograms with the toluene-*p*-sulphonic acid reagent, and R_F values were noted (Table 2). King & Bottomley's (1954) melacacidin was found to consist of two compounds running to different R_F values in each solvent system. The shade of red developed with the reagent on heating varied for the different leuco-anthocyanins. Leuco-fisetinidin (7:3':4'-trihydroxyflavan-3:4-diol) developed a purple shade initially which became light pink, whereas the melacacidins developed a deep red. The others formed red colorations of intermediate shade.

These synthetic and naturally occurring flavan-3:4-diols have a higher R_F value in 2% acetic acid or in water than catechins and 2:3-dihydroflavonols (Table 2), and are also more highly mobile than the polymeric tannins. Two-dimensional chromatograms run under the above conditions may therefore be used to locate monomeric flavan-3:4-diols (R_F 0.47–0.55 in 2% acetic acid) and also polymeric leuco-anthocyanins (R_F 0–0.45 in 2% acetic acid) with the aid of the toluene-*p*-sulphonic acid reagent.

Of 144 'heartwoods' examined from trees generally classified as 'hard-woods' (mainly those indigenous to South Africa, but also including

exotic types), 44 (33%) gave strongly positive leuco-anthocyanin reactions, and 17 (12%) showed chromatographic evidence of the presence of monomeric forms (Table 3). The monomeric leuco-anthocyanins are usually accompanied by polymeric leuco-anthocyanins of lower R_F in 2% acetic acid. These were shown up prominently by the toluene-*p*-sulphonic acid reagent. In most species examined only polymeric leuco-anthocyanins are present, as judged by their low R_F on paper chromatograms and the formation of anthocyanidins under the conditions of Pigman *et al.* (1953).

Table 2. R_F values of flavan-3:4-diols, catechins and 2:3-dihydroflavonols

	R_F in developing solvent	
	Butan-1-ol-acetic acid-water	2% Acetic acid
Flavan-3:4-diols		
7:3':4'-Trihydroxy-	0.76	0.49
7:3':4':5'-Tetrahydroxy-	0.57	0.55
7:8:3':4'-Tetrahydroxy-	0.42	0.47
	0.50	0.54
5:7:3':4'-Tetrahydroxy-	0.52	0.53
Catechins		
(+)-Gallocatechin	0.57	0.33
(+)-Catechin	0.76	0.39
(±)- <i>epi</i> -3:7:3':4'-Tetrahydroxyflavan	0.92	0.42
		0.47
2:3-Dihydroflavonols		
Dihydrofisetin (fustin)	0.90	0.35
Dihydromyricetin	0.74	0.36
Dihydroquercetin (taxifolin)	0.90	0.29
Dihydromyricetin (ampeloptin)	0.70	0.35

Table 3. *Leuco-anthocyanins in the 'heartwoods' of dicotyledonous plants*

All heartwoods listed give a strong reaction for leuco-anthocyanins; those marked * also contain monomeric leuco-anthocyanins.

Anacardiaceae	<i>Harpephyllum caffrum</i> , <i>Rhus legati</i> *, <i>Schinopsis quebracho-colorado</i> * (syn. <i>S. lorentzii</i>), <i>S. balansae</i> *
Celastraceae	<i>Gymnosporia acuminata</i> , <i>Pleurostylia capensis</i> , <i>Pterocelastrus tricuspidatus</i> *
Cornaceae	<i>Curtisia faginea</i>
Cunoniaceae	<i>Cunonia capensis</i>
Ebenaceae	<i>Diospyros mespiliformis</i> , <i>Euclea lanceolata</i> *
Euphorbiaceae	<i>Phyllanthus discoideus</i>
Flacourtiaceae	<i>Kiggelaria africana</i> , <i>Scolopia munduti</i>
Leguminosae	
Caesalpinioidae:	<i>Baikiaea minor</i> , <i>B. plurijuga</i> , <i>Brachystegia apendulata</i> *, <i>B. spiciiformis</i> , <i>Burkea africana</i> , <i>Copaifera coleosperma</i> *, <i>C. tessmannii</i> *, <i>Erythrophloeum africana</i> , <i>E. guineense</i> , <i>Isobertlinia tomentosa</i> , <i>Schotia latifolia</i> *
Mimosoideae:	<i>Acacia dealbata</i> *, <i>A. decurrens</i> *, <i>A. melanoxylon</i> *, <i>A. mollissima</i> *, <i>A. nigrescens</i> *, <i>A. pycnantha</i> *
Papilionatae:	<i>Berlinia niambiensis</i>
Myristicaceae	<i>Staudtia gabonensis</i>
Myrtaceae	<i>Eucalyptus marginata</i> , <i>Syzygium cordatum</i>
Ochnaceae	<i>Ochna arborea</i> , <i>Ochna</i> sp.*
Rhamnaceae	<i>Rhamnus zeyheri</i> *
Rosaceae	<i>Pygeum africanum</i>
Rhizophoraceae	<i>Weinmannia africana</i>
Sapotaceae	<i>Mimusops obovata</i> , <i>Manilkara cuneifolia</i>
Ulmaceae	<i>Trema guineensis</i>

Generation of anthocyanidins from wattle and quebracho extracts

Quebracho tannins. Quebracho extract yields fisetinidin (mainly) and also a second anthocyanidin (traces) when heated with HCl in propan-2-ol solution under pressure (Roux, 1957*a*). On paper chromatograms the whole of the lower R_f area containing the tannins gives the leuco-anthocyanin reaction with toluene-*p*-sulphonic acid spray reagent, but the presence of other constituents, probably catechins, is indicated as brown spots (Fig. 2). This behaviour indicates the presence of leuco-anthocyanins throughout the quebracho-tannin fractions. On the paper chromatogram the leuco-anthocyanins overlap each other so as to form a continuous band. This streaking without separation into discrete spots may result from the complexity of the mixture and also from the structural complexity of the individual tannins. Brown constituents which remain on the starting-line in direction 2 do not react readily with the toluene-*p*-sulphonic acid spray, but nevertheless yield fisetinidin on treatment with HCl in propan-2-ol solution under pressure.

No pink coloration developed in that position (Fig. 2) where the monomeric leuco-fisetinidin, 7:3':4'-trihydroxyflavan-3:4'-diol, should be located. The absence of Freudenberg & Maitland's (1934) 'quebracho catechin', or (±)-*epi*-7:3':4'-trihydroxyflavan-3-ol, from commercial quebracho extract has also been established (Roux, 1957*b*).

Black-wattle tannins. Black-wattle extracts forms three pigments on treatment with HCl under the conditions of Pigman *et al.* (1953). Two of these have been identified as the anthocyanidins robinetinidin (major proportion) and fisetinidin, but the identity of the third pigment is unknown (Roux,

1957*b*). Robinetinidin is the only anthocyanidin generated from constituents *C* and *E*, whereas both robinetinidin and fisetinidin are generated from A_1 . Constituents *B* and *D* give mainly the unidentified orange-yellow coloration. The anthocyanidins and the unknown pigment are also generated in about the same proportions as from the whole extract, from that fraction of wattle tannins which remains immobile on the starting-line in direction 2 (Roux, 1957*b*).

Yields of anthocyanidins. The amount of anthocyanidin generated from simple or complex leuco-anthocyanins increases rapidly over the first hour, but apparently the rate of reaction declines rapidly after this period. Robinetinidin and fisetinidin also undergo partial transformation into the corresponding flavonols, resulting in a decrease in the absorption density at the absorption maximum of each coloration, and other variables (Pigman *et al.* 1953) also affect the yield of anthocyanidins from leuco-compounds.

To obtain an indication of the extent to which leuco-anthocyanins occur in wattle and quebracho extracts, a comparative study of the yields of anthocyanidins from monomeric and complex leuco-anthocyanins of similar type was made. Thus most of the above variables are eliminated.

Quebracho extract yields mainly fisetinidin ($\lambda_{\max} = 525 \text{ m}\mu$), but for black-wattle extract three pigments are generated, ranging from 470 to 532 $\text{m}\mu$ in absorption maxima. Measurement of the percentage conversion for quebracho (at 525 $\text{m}\mu$) and for wattle (at 532 $\text{m}\mu$), as represented in Table 4, is limited by these factors, but nevertheless furnishes approximate values. Polyphenols and extracts were treated with HCl for 2 hr. under the conditions of Pigman *et al.* The percentage conversion is calculated from the absorption-density values of synthetic robinetinidin ($\text{C}_{15}\text{H}_{11}\text{O}_6\text{Cl}$, absorption density = 0.810 at 532 $\text{m}\mu$ and 3.5 mg./500 ml.), and fisetinidin ($\text{C}_{15}\text{H}_{11}\text{O}_6\text{Cl}, \text{H}_2\text{O}$, absorption density = 0.595 at 525 $\text{m}\mu$ and 2.3 mg./500 ml.). Correction is made during calculation of yields for the water content of fisetinidin.

Range of molecular weights in condensed tannins

Estimation of molecular weights of tannin fractions. The molecular weights of the extracts and of their fractions of different R_f values, as well as their weight distribution and percentage recovery, are given in Table 5. Wattle extracts C_1 and C_3 were obtained from the same bark (Table 1). No attempts were made to elute the strips exhaustively, as oxidation effects are likely to result from prolonged elution. The recovery of simple polyphenols, e.g. catechins, from paper chromatograms is, for example, never quantitative, in spite of their

Fig. 2. Two-dimensional chromatogram of quebracho tannins, indicating relative positions of *epi*-7:3':4'-trihydroxyflavan-3-ol ('quebracho catechin') and 7:3':4'-trihydroxyflavan-3:4'-diol ('leuco-fisetinidin'). See text.

Table 4. *Yield of anthocyanidins from tannins and flavan-3:4-diols*

Material	Weight (mg.)	Absorption density at 525 or 532 m μ	Conversion* into fisetinidin or robinetinidin (%)
Commercial quebracho	2.9	0.320	5.6†
7:3':4'-Trihydroxyflavan-3:4-diol	6.7	0.438‡	23.8
Commercial wattle	3.7	0.366	6.3†
7:3':4':5'-Tetrahydroxyflavan-3:4-diol	2.9	0.201‡	26.9
Wattle fraction 4A1 ('constituent' E)	2.2	0.169	3.0
Wattle fraction 3A1 ('constituent' C)	2.9	0.378	5.1
Wattle fraction 2A1 ('constituent' A1)	2.6	0.296	4.4

* Conversions for quebracho and leuco-fisetinidin calculated as for fisetinidin, others as for robinetinidin.

† Calculated to a 'pure' tannin basis.

‡ Diluted to 500 ml., others to 50 ml.

Table 5. *Molecular-weight distribution of wattle and quebracho tannins*

Mean R_F	Wattle fraction C_1		Wattle fraction C_3		Quebracho extract	
	Tannins eluted (%)	Mol.wt.	Tannins eluted (%)	Mol.wt.	Tannins eluted (%)	Mol.wt.
0'	—	—	24.3	3240	—	—
0	3.7	1442	15.7	1631	21.0	2350
0.1	20.3	1287	24.3	1606	17.1	1811
0.2	24.3	1033	22.1	1203	16.5	1362
0.3	29.7	908	13.6	933	19.7	1066
0.4	15.6	649	—	—	17.0	910
0.5 + 0.6	6.5	554	—	—	8.7	790
Recovery (%)	71.0	—	74.0	—	95.0	—
Calc. av.mol.wt.	—	971	—	1826	—	1461
Determined av.mol.wt.	—	1284	—	1507	—	1327

Table 6. *Percentage yield of anthocyanidins from wattle- and quebracho-tannin fractions*

Mean R_F of fraction	Wattle fraction C_1	Wattle fraction C_3	Quebracho tannins
0'	—	4.39	—
0.0	4.33	4.96	3.36
0.1	5.78	4.50	3.56
0.2	4.03	5.09	4.00
0.3	4.34	3.89	4.53
0.4	3.63	—	4.08
0.5 + 0.6	2.27	—	3.34

relatively low affinity for cellulose. Only small quantities of residual tannins remained on the strips after elution with ethanol-water mixtures, with the exception of R_F 0 of wattle-fraction C_3 . This band was subsequently eluted with acetone-water (1:1) to give fraction 0'.

Generation of anthocyanidins from tannin fractions. Examination by two-way paper chromatography of each fraction of wattle and quebracho tannins obtained from the chromatographic separation on thick sheets showed that each fraction was broad in direction 1 (butanol-acetic acid-water or water-saturated butan-2-ol) but narrow in direction 2 (2% acetic acid). R_F values of the

bands in direction 2 correspond approximately to those of the same bands when cut from the thick sheets.

The anthocyanidins generated from each fraction of black-wattle tannins contained robinetinidin, fisetinidin and an orange-yellow pigment. The proportions of these pigments varies with the different fractions. Quebracho tannins yield mainly fisetinidin from each fraction. The yields of anthocyanidins from the various fractions are given in Table 6. These are free of 'non-tannin' carbohydrates, and their values are therefore comparable.

DISCUSSION

Since the first recognition of the widespread distribution of leuco-anthocyanins by Robinson & Robinson (1933, 1934) and the use of qualitative colour reactions for the identification of anthocyanidins, the method of recognizing these classes of compounds was enhanced by the use of paper chromatography. Bate-Smith (1954) used this technique for identifying anthocyanidins generated from leuco-anthocyanins conforming to the delphinidin and cyanidin types, and Roux (1957*a, b*) extended the range of anthocyanidins, their

aluminium complexes (Geissman, Jorgenson & Harborne, 1953; Geissman & Jurd, 1955) and a new solvent system being used to assist in their differentiation.

Roux (1957*b*) also used a new chromogenic reagent, toluene-*p*-sulphonic acid, for locating leuco-anthocyanins on paper chromatograms and for differentiating between monomeric and polymeric leuco-anthocyanins. The reagent gives various scarlet colorations with different monomeric (Table 2) and naturally occurring polymeric leuco-anthocyanins, and from their relative positions on paper chromatograms the monomeric (R_F 0.47–0.55 in 2% acetic acid) and polymeric (R_F 0–0.45) forms may be differentiated. Monomeric leuco-anthocyanins, assuming adequate separation has been achieved, will also appear as discrete pink spots, whereas complex leuco-anthocyanins are usually represented as broad pink bands extending often continuously from R_F 0.45 to the origin. The use of these methods is demonstrated in the examination of the heartwoods of a large number of species, where the high proportion of monomeric forms present amongst certain genera of the family Leguminosae is of special interest (Table 3).

For quebracho extract, the overall development of pink colorations with toluene-*p*-sulphonic acid reagent in almost all those areas on two-way paper chromatograms where reducing tannins are located shows the wide distribution of leuco-anthocyanins throughout the tannin fraction. A close inter-relationship between various tannin constituents with different mobilities on paper chromatograms also appears to exist, as independent of their rate of migration and their molecular weights (Table 5) each fraction contains leuco-fisetinidins (Table 6). The complex leuco-fisetinidins all have lower R_F values in 2% acetic acid than monomeric leuco-fisetinidins, and undoubtedly possess greater structural complexity (mol.wt. 790–2350) than the monomeric form (mol.wt. 290).

The lower yield of fisetinidin from quebracho leuco-anthocyanins (5.6%) compared with 7:3':4'-trihydroxyflavan-3:4-diol (23.8%) indicates that approx. 24% of the 'fundamental units' present in the tannins possibly are of the 'leuco-fisetinidin type'. Considering the even distribution of leuco-anthocyanins throughout the tannins on two-way paper chromatograms, this figure may be on the low side. This evidence, however, demonstrates the $C_6-C_3-C_6$ nature of a fairly high proportion of the 'fundamental units' present in quebracho tannins.

The work on wattle extract closely parallels that on quebracho, but the leuco-anthocyanin mixture present in wattle is more complex. Whereas the leuco-anthocyanins of quebracho extract are mainly leuco-fisetinidins, black-wattle tannins

contain leuco-robinetinidins predominantly, but also considerable proportions of leuco-fisetinidins and other constituents capable of forming an unidentified orange-yellow pigment. Furthermore, whereas the leuco-fisetinidins of quebracho extract are fairly evenly distributed throughout the reducing tannins on paper chromatograms, the mobile leuco-robinetinidins, leuco-fisetinidins and other colour-producing compounds of wattle extract may be differentiated amongst the mobile tannins, although a considerable amount of overlap of constituents occurs. Such overlap is to be anticipated in the migration of highly polar units of large molecular weight and high affinity for cellulose, on paper chromatograms. The proportion of fisetinidin generated from the extracts of barks from individual trees reputed to be authentic examples of *Acacia mollissima* varies somewhat. Bark extracts of the closely related species, silver wattle (*A. dealbata*) and green wattle (*A. decurrens*), furnish the same anthocyanidins but in both instances the percentage of fisetinidin generated appears to be higher compared with black wattle. The proportion of leuco-anthocyanins present in the complex wattle tannins (molecular weight 550–3250) appears to be of the same order as those present in quebracho (Table 4).

Both wattle and quebracho tannins on alkaline fusion yield mainly those phenolic nuclei (resorcinol and protocatechuic acid from quebracho tannins, and resorcinol, gallic and protocatechuic acids from wattle; Freudenberg & Maitland, 1934; Roux, 1952) which are potentially present in the anthocyanidins generated from the extract.

For studying the range and distribution of molecular weights in relation to leuco-anthocyanin distribution in wattle and quebracho tannins, the method of fractionating the tannins on cellulose sheets was used. This method has the disadvantage that recoveries of the polyphenolic tannins by elution is not quantitative. Nevertheless, by using a paper-chromatographic method, the fractions isolated may be linked directly to the distribution of complex leuco-anthocyanins on two-way paper chromatograms.

Wattle tannins range in molecular weight mainly from 550 to 1630, but also contain a small fraction of much higher (3240) average molecular weight. Quebracho tannins are evenly distributed throughout the range 800–2350, but the molecular weight of quebracho tannins in each R_F fraction is comparatively higher than the value for the corresponding wattle fraction (Table 5).

The complex wattle and quebracho tannins both show a stepwise increase of molecular weight with increase of affinity for the cellulose substrate in aqueous medium. As wattle tannins show parallel affinity characteristics for both cellulose and

collagen during chromatographic separations (Roux, 1955), it follows that with increase of molecular weight tannins show greater affinity also for collagen. The relationship between affinity and molecular weight is important in the interpretation of the chromatographic behaviour of 'condensed' tannins, and is also of interest in studying the behaviour of tannins towards collagen, where the affinity effects are believed to be mainly due to hydrogen bonding (Shuttleworth & Cunningham, 1948; Ellis & Pankhurst, 1954).

From the two-dimensional chromatography of each fraction and also from the regular increase of molecular weight with decrease of R_f value may be gauged the efficiency of the fractionation procedure. The relatively even distribution of leuco-anthocyanins in these fractions throughout the whole R_f range (Table 6) confirms similar evidence obtained by spraying two-dimensional chromatograms of the tannins with the toluene-*p*-sulphonic acid reagent, and shows that complex leuco-anthocyanins of varying molecular weight, molecular complexity and affinity for various substrates are present in both wattle and quebracho tannins.

SUMMARY

1. Examination of the heartwood extractives of 144 'hardwood' species by methods developed by Roux (1957*a, b*) show that 33% give strongly positive reactions for leuco-anthocyanins. Lower R_f polymeric forms predominate, and in only 12%, mainly amongst the Leguminosae, is evidence found of the presence of monomeric leuco-anthocyanins.

2. Black-wattle and quebracho tannins have been shown to be relatively rich in complex leuco-robinetinidins and leuco-fisetinidins (Roux, 1957*b*). Comparison of the yields of anthocyanidins from complex naturally occurring leuco-anthocyanins and from synthetic monomeric leuco-anthocyanins indicates that about 25% of the 'fundamental units' present in these complex tannins are of the leuco-anthocyanin type.

3. For each tannin mixture, a gradational increase of molecular weight with decrease of R_f value of the tannins was found, when separated on paper chromatograms in 2% acetic acid. The molecular weights of wattle tannins fall mainly in the range 550–1630, although a small fraction of much higher (3240) molecular weight is also present. Values for quebracho tannins fall in the range 800–2350.

4. Similar proportions of leuco-anthocyanins are present throughout the whole molecular-weight range of both tannins, indicating the presence of leuco-anthocyanins (e.g. leuco-robinetinidins and leuco-fisetinidins) of increasing complexity from the higher R_f (0.5–0.6) to the lower R_f (0) regions.

5. With increase in molecular weight condensed tannins show a parallel increase in 'affinity' for cellulose, and also for collagen (Roux, 1955).

Thanks are due to Dr P. Maitland, Department of Chemistry, Cambridge University, for a specimen of ($\pm$)-*epi*-3:7:3':4'-tetrahydroxyflavan; to Dr T. Kubota, Institute of Polytechnics, Osaka, Japan, for a specimen of ampeloptin, and to Miss G. V. Britten and Mr D. G. Comins, of the Botanical Survey, Union Department of Agriculture, Albany Museum, Grahamstown, for their assistance in the identification of plant material.

This work is financed by the annual grant of the South African Wattle Growers' Union to the Leather Industries Research Institute.

REFERENCES

- Bate-Smith, E. C. (1954). *Biochem. J.* **58**, 122.
 Bate-Smith, E. C. & Swain, T. (1953). *Chem. & Ind.* p. 377.
 Bradfield, A. E. & Flood, A. E. (1953). *J. chem. Soc.* p. 4740.
 Ellis, S. C. & Pankhurst, K. G. A. (1954). *Disc. Faraday Soc.* **16**, 170.
 Evelyn, S. R. (1954*a*). *J. Soc. Leath. Tr. Chem.* **38**, 142.
 Evelyn, S. R. (1954*b*). *J. Soc. Leath. Tr. Chem.* **38**, 309.
 Evelyn, S. R., van Berge, P. C. & Cooper, D. R. (1958). *J. Polym. Sci.* (in the Press).
 Freudenberg, K. & Maitland, P. (1934). *Liebigs Ann.* **510**, 193.
 Freudenberg, K. & Roux, D. G. (1954). *Naturwissenschaften*, **41**, 450.
 Freudenberg, K. & Weinges, K. (1955). *Liebigs Ann.* **590**, 140.
 Geissman, T. A., Jorgenson, E. C. & Harborne, J. B. (1953). *Chem. & Ind.* p. 1389.
 Geissman, T. A. & Jurd, L. (1955). *Arch. Biochem. Biophys.* **56**, 259.
 Hathway, D. E. & Seakins, J. W. T. (1957). *Biochem. J.* **67**, 239.
 Hillis, W. E. (1956*a*). *Aust. J. Biol. Sci.* **9**, 362.
 Hillis, W. E. (1956*b*). *Symp. Soc. Leath. Tr. Chem., Vegetable Tannins*, p. 121.
 King, F. E. & Bottomley, W. (1954). *J. chem. Soc.* p. 1399.
 King, F. E. & Clark-Lewis, J. W. (1955). *J. chem. Soc.* p. 3384.
 Nordström, C. G. & Swain, T. (1953). *J. chem. Soc.* p. 2764.
 Pigman, W., Anderson, E., Fischer, R., Buchanan, M. A. & Browning, B. L. (1953). *T.A.P.P.I.* **36**, 4.
 Ray, N. H. (1952). *Trans. Faraday Soc.* **48**, 809.
 Reith, W. S. (1957). *Nature, Lond.*, **179**, 580.
 Robinson, G. M. & Robinson, R. (1933). *Biochem. J.* **27**, 206.
 Robinson, G. M. & Robinson, R. (1934). *Biochem. J.* **28**, 1712.
 Roux, D. G. (1951). *J. Soc. Leath. Tr. Chem.* **35**, 322.
 Roux, D. G. (1952). *J. Soc. Leath. Tr. Chem.* **36**, 393.
 Roux, D. G. (1955). *J. Soc. Leath. Tr. Chem.* **39**, 80.
 Roux, D. G. (1957*a*). *Nature, Lond.*, **179**, 305.
 Roux, D. G. (1957*b*). *Nature, Lond.*, **180**, 973.
 Shuttleworth, S. G. & Cunningham, G. E. (1948). *J. Soc. Leath. Tr. Chem.* **32**, 202.
 Smith, H. (1956). *Trans. Faraday Soc.* **52**, 402.
 Swain, T. (1954). *Chem. & Ind.* p. 1144.
 Vieboch, F. & Brecher, C. (1930). *Ber. dtsch. chem. Ges.* **63**, 2818, 3207.