SUMMARY

1. Pea seeds treated with maleic hydrazide germinated normally until cell enlargement was replaced by cell division, when growth stopped.

2. The ratio of reduced (GSH) to oxidized (GSSG) glutathione, i.e. GSH:GSSG, in cotyledons was not affected by maleic hydrazide but at the onset of cell division GSH increased at the expense of GSSG in treated axes only.

3. Certain enzymes requiring free sulphhydryl groups (starch phosphorylase and β -amylase) were inhibited irreversibly by maleic hydrazide, but other enzymes (trypsin and pea aldolase) were not affected.

4. The results suggest that maleic hydrazide reacts with protein SH groups and thus inhibits the reduction of protein S·S by GSH during mitosis.

We are indebted to Mr H. Holland for the microscopical examinations of the radicle tips and to Dr F. Haworth for helpful discussions during the course of the work.

REFERENCES

- Andrae, W. A. & Andrae, S. R. (1953). *Canad. J. Bot.* **31**, 426.
- Beck, W. S. (1955). *J. biol. Chem.* **212**, 847.
- Benesch, R. E., Lardy, H. A. & Benesch, R. (1955). *J. biol. Chem.* **216**, 663.
- Bertossi, F. (1955). *Année biol.* **LIX**, 231.
- Dohan, J. S. & Woodward, G. E. (1939). *J. biol. Chem.* **129**, 393.
- Elliot, K. A. C. (1946). *Annu. Rev. Biochem.* **15**, 1.
- Grunert, R. R. & Phillips, P. H. (1951). *Arch. Biochem. Biophys.* **30**, 217.
- Hanes, C. S. (1940). *Proc. Roy. Soc. B*, **129**, 174.
- Hopkins, F. G. & Morgan, E. F. (1943). *Nature, Lond.*, **153**, 288.
- Isenberg, F. M. R., Odland, M. L., Prop, H. W. & Jensen, C. D. (1951). *Science*, **113**, 1951.
- Leopold, A. C. & Klein, W. H. (1951). *Science*, **114**, 9.
- Leopold, A. C. & Price, C. A. (1956). *Chemistry of Plant Growth Substances*, p. 271. London: Butterworths Scientific Publications.
- Liener, I. E. (1957). *J. biol. Chem.* **225**, 1061.
- Mapson, L. W. & Goddard, D. R. (1951). *Biochem. J.* **49**, 592.
- Mapson, L. W. & Moustafa, E. M. (1956). *Biochem. J.* **62**, 248.
- Mazia, D. (1954). Symposium on: *Glutathione*, p. 209. New York: Academic Press Inc.
- McLeish, J. (1953). *Heredity*, suppl. **6**, 125.
- Miller, D. M. & White, R. W. (1956). *Canad. J. Chem.* **34**, 1510.
- Morgan, E. J. & Friedman, E. (1938). *Biochem. J.* **32**, 733.
- Muir, R. M. & Hansch, C. (1953). *Plant Physiol.* **28**, 218.
- Nickerson, W. J. & Falcone, G. (1956). *Science*, **124**, 722.
- Peel, J. L. & Loughman, B. C. (1957). *Biochem. J.* **65**, 709.
- Somogyi, M. (1945). *J. biol. Chem.* **160**, 61.
- Stumpf, P. K. (1948). *J. biol. Chem.* **176**, 233.
- Weill, C. E. & Caldwell, M. L. (1945). *J. Amer. chem. Soc.* **67**, 212.
- Weller, L. E., Ball, C. D. & Sell, H. M. (1957). *Plant Physiol.* **32**, 146.
- Yemm, E. W. & Cocking, E. C. (1955). *Analyst*, **80**, 209.

Formation of Antibody by Isolated Perfused Lungs of Immunized Rabbits

THE USE OF [¹⁴C]AMINO ACIDS TO STUDY THE DYNAMICS OF ANTIBODY SECRETION

BY BRIGITTE A. ASKONAS AND J. H. HUMPHREY
National Institute for Medical Research, Mill Hill

(Received 24 February 1958)

Under the usual experimental conditions yields of newly formed proteins secreted by tissue slices are too small to permit their isolation for study without carrier protein, and when whole organs, such as the liver, are perfused with blood (e.g. Miller, Bly, Watson & Bale, 1951) recovery of any newly synthesized plasma proteins is complicated by admixture with similar proteins already present in the perfusion fluid.

Normal lung forms little or no γ -globulin, nor is it known to secrete any other protein. When

rabbits are hyperimmunized by intravenous injections of a particulate antigen, such as pneumococci type 3, the lung may become a major site for antibody formation, as shown both *in vivo* (Humphrey & Sulitzeanu, 1958) and by tissue slices *in vitro* (Askonas & Humphrey, 1958). As a result of hyperimmunization there is enormous proliferation of lymphoid tissue and plasma cells around the small arteries and the lung may increase two to four times in weight. If such lungs are perfused with salt medium containing normal

rabbit serum, antibody is secreted and can be specifically precipitated from the perfusing fluids by adding the appropriate antigen. A considerable net synthesis both of antibody (up to 8 mg./hr.) and of other γ -globulin can be demonstrated.

For a study of protein synthesis this system has, however, two disadvantages common to other perfused organs. The first is that tissues contain several kinds of cells, and the second that it is impossible to wash out all preformed plasma proteins from extracellular spaces so as to prevent their mixing with proteins newly formed during the experiment. The second difficulty can be overcome by injecting into the animal, 1 or 2 days before death, specific antibody labelled with ^{131}I , as described by Humphrey & Sulitzeanu (1958). Such antibody equilibrates with the proteins in the extracellular fluids, and if it is assumed that at the time of death the ^{131}I specific activities of antibody in the plasma and tissue fluids are the same, the proportion of preformed antibody in samples isolated during the experiment can easily be calculated.

In the experiments described below the incorporation of ^{14}C amino acids into intracellular and extracellular antibody was followed, and the specific radioactivity of intracellular and extracellular free amino acids was studied during the perfusion. The bearing of our findings on the secretion of antibody and other proteins is discussed. By secretion we mean the release from within the cell into the surrounding medium of protein derived from the intracellular pool, but not necessarily newly formed during the experimental period. Evidence is also given that the lung from an immune animal may be a site of plasma-protein breakdown.

EXPERIMENTAL

Immunization procedure. Sandylop rabbits (2.5–2.8 kg.), bred at the N.I.M.R., were given intravenous injections of pneumococci type 3 killed with formalin or of alum-precipitated ovalbumin or both. Injections were made on alternate days for 3–4 weeks, the quantity of antigen injected being gradually increased up to 10^{10} pneumococci or 0.5 mg. of ovalbumin. The animals were test-bled at intervals. Antibody estimations were made by precipitating with antigen in the equivalence zone, followed by estimation of γ -globulin in the washed precipitates by dissolving in 0.1N-NaOH and measuring the light absorption at 280 μ .

Administration of ^{131}I antibody. γ -Globulin concentrates were prepared by fractionation with 16% (w/v) Na_2SO_4 of sera from sandylop rabbits hyperimmunized against pneumococci type 3. Of the protein in the preparations, 70–75% was specifically precipitable with pneumococci type 3 capsular polysaccharide, prepared by the method of Heidelberger, Kendall & Scherp (1936). The globulin concentrates were iodinated with ^{131}I by the jet-mixing method of McFarlane (1956), so as to contain not

more than 1 g. atom of I/mole of protein. About $100\mu\text{C}$ of ^{131}I protein (5–16 mg.) was injected intravenously 2 days before the experiment. By precipitating antibody from a serum sample taken at the time of death and measuring its γ -ray radioactivity, the ^{131}I specific radioactivity of preformed extracellular antibody was determined.

Perfusion of the isolated lung. After removal of a preliminary blood sample from the rabbit 500 i.u. of heparin followed by 120 mg. of sodium pentobarbitone were injected intravenously. The heart and lungs were carefully removed and rinsed, and one cannula was inserted into the trachea and another into the pulmonary artery; through the latter 500 ml. of cold balanced-salt solution (Hanks, 1948) was perfused to wash out most of the blood. After trimming away most of the heart and loose tissue, the lung was suspended in the apparatus illustrated in Fig. 1.

The apparatus, for whose design we are indebted to Mr W. E. Brocklehurst, consisted essentially of a jacketed vessel and a pump for recirculating a small volume of perfusion fluid at 39.5° . The lung was alternately inflated and deflated with $\text{O}_2 + \text{CO}_2$ (95:5) gas mixture by means of a respiration pump (Palmer and Co., Brixton). In some experiments the surface of the lung, especially at the bases, was scratched gently with a sharp needle in order to diminish accumulation of fluid which sometimes occurred after perfusion for some hours. In most experiments this was not necessary.

Fig. 1. Arrangement of apparatus for perfusion of rabbit lung. L, Lung; P_1 , small-volume force pump for recirculating about 50 ml. of medium through the lung; P_2 , pump circulating water from a bath at 39.5° ; P_3 , respiration pump supplying $\text{O}_2 + \text{CO}_2$ (95:5) (stroke volume 50 ml.); T, cannulated trachea; O, overflow for perfusate; C, cannula in pulmonary artery; V, jacketed vessel.

The perfusion mixture consisted of 90 ml. of isotonic salt medium (Hanks, 1948), containing 0.27 g. of glucose, 5000 units of sodium penicillin, 1–3 μ C of [14 C]glycine, L-[14 C]lysine or L-[14 C]valine, a complete amino acid mixture (15 mg. of N) prepared from acid-hydrolysed plasma protein with added L-tryptophan and 10 ml. of fresh normal rabbit serum. In experiments where the specific radioactivity of the free valine was determined, the medium contained known amounts of all the L-amino acids (15 mg. of N/100 ml. of medium). The rabbit serum was obtained by collecting blood without anticoagulant into chilled silicone-treated centrifuge tubes (Silicone Fluid M.S. 200/350 ca., Hopkin and Williams Ltd.), and centrifuging the cells and platelets in the cold, and transferring the plasma to glass tubes and allowing it to clot spontaneously. This was done to avoid the high levels of histamine and 5-hydroxytryptamine, which occur in fresh clotted rabbit serum as a result of platelet destruction during clotting and can cause initial pulmonary vasoconstriction.

The lung was perfused with a known volume (about 50 ml.) of fluid at about 30 ml./min. In most experiments the complete medium, but without [14 C]amino acid, was circulated for about 30 min. and was then replaced by labelled medium; after this the perfusion fluid was collected and replaced by new medium every 60–90 min. to maintain the level of radioactivity and of amino acids. To make it easier to judge when one lot of medium had been completely replaced by fresh, and to assess how effectively the lung was perfused, two drops of 5% Pontamine Sky Blue (6 Bx) were added to alternate changes of medium.

Radioactive amino acids. Biosynthetic [14 C]glycine, L-[14 C]lysine and L-[14 C]valine were obtained from the Radiochemical Centre, Amersham, Bucks, each with a specific activity of about 55 μ C/mg.

Isolation of antibody and γ -globulin from perfusates. About 50 mg. of the appropriate inactive amino acid was added to each lot of perfusing fluid, to dilute the radioactive amino acid and decrease non-specific adsorption on to proteins. The perfusates were dialysed under pressure to a volume of 8–10 ml. against 0.6% NaCl solution containing 66 mM-sodium phosphate buffer, pH 7.5. Any sediment was removed by centrifuging for 15 min. at 15 000 g, and more inactive amino acid (30–50 mg.) and Merthiolate (Eli Lilly and Co.) to a concentration of 0.1 mM were added to the supernatant fluid.

Specific antibody was then precipitated by addition of an appropriate amount of antigen. A rough idea of the quantity to be added could be obtained by determining the amount of extracellular antibody present by measurement of the 131 I. Since extracellular antibody usually accounted for less than half of the antibody isolated, enough antigen was added in the first instance to precipitate twice the estimated extracellular antibody. The mixture was incubated for 20 min. at 37° and left overnight in the cold room. Antigen-antibody precipitate was centrifuged down at low speed (about 3 min. at 1600 g). A little more antigen was added to the supernatant and the process repeated until no more precipitate was obtained. The pooled precipitates were washed thrice with 0.9% NaCl solution, once with aq. 50% (v/v) ethanol, twice with ethanol, and dried with ether and weighed. Radioactivity due to 131 I and 14 C was determined as described below.

A sample of the remaining γ -globulin could be isolated

from the perfusing fluid after removal of the specific antigen-antibody precipitates by successive precipitations with Na_2SO_4 at concentrations of 18 and 16% (w/v). The fraction soluble in 0.9% NaCl after dialysis against water was freeze-dried. This fraction contained only γ -globulin as tested by paper electrophoresis. It included both γ -globulin secreted by the lung and the normal rabbit γ -globulin added to the perfusion medium. However, any 14 C radioactivity must have been due to γ -globulin newly formed from [14 C]amino acids in the medium.

Isolation of intracellular antibody from lung. In some cases one lobe of the lung was tied off and removed part way through the experiment and intracellular antibody was isolated from this portion as described below. At the end of each experiment, except when stated otherwise, the lung was perfused for about 3 min. with Hanks's solution without added amino acids, and weighed and frozen at -30° . The frozen lung was weighed, chopped and extracted in a high-speed blender at 0° with 5 ml. of normal rabbit serum and about thrice its weight of saline phosphate buffer, pH 7.5, containing 50 mg. of inactive amino acid. The material sedimenting in 10 min. at 1600 g was re-extracted with about half the previous quantity of saline phosphate buffer, and the combined extracts were centrifuged for 20 min. at 22 000 g. The opalescent supernatant may be expected to contain a representative sample of intracellular lung antibody, although complete extraction can hardly be claimed. After pressure dialysis of the extract against saline phosphate buffer to reduce its volume to about 20 ml., the pH was adjusted to 5.8 in order to aggregate finely suspended particles and some of the nucleoprotein. After centrifuging, the clear supernatant was neutralized and antibody was isolated as described for the perfusion fluids above. Specific radioactivities due to 131 I and 14 C were measured on the pooled antigen-antibody precipitates, since it had been found in early experiments that the specific radioactivities of the first and second precipitates in any one sample did not differ significantly.

Radioactivity of free valine in the perfusate and in cells of the lung. The lung was perfused with 60 ml. of medium containing 6 ml. of normal rabbit serum, 50 mg. of glutamine and 43 mg. of amino acids, of which 3.4 mg. was L-valine, including 1.8 μ C of L-[14 C]valine (18.3 μ C/ μ C). The medium was changed every 80–90 min. during a total period of 5½ hr. Samples (10 ml.) were taken from the original medium, and from the perfusing fluids 10 min. after the beginning and at the end of each perfusion period, and added to equal volumes of 20% (w/v) trichloroacetic acid; precipitated protein was removed, the supernatant was freed from trichloroacetic acid by repeated extractions with ether and its volume reduced to 2–3 ml. by evaporation. After neutralization, the samples were desalted on a column of Zeo-Karb 225 in the H^+ form (1 cm. $\times$ 4.5 cm.). The sample was washed through the column with 30 ml. of water, and the amino acids were eluted with aq. 2N- NH_3 soln. and taken to dryness over H_2SO_4 in a desiccator. About a quarter of the material was applied on chromatography paper (Whatman no. 3) along a line 4 cm. long, and the chromatogram developed for 48 hr. with redistilled collidine-lutidine-water (1:1:2, by vol.). Marker valine was spotted 4 cm. away from each sample. A strip containing the marker valine and a small part of the sample was cut out, and the location of the valine was determined by spraying with ninhydrin reagent. Valine was then eluted

from the unsprayed papers with aq. 50% (v/v) ethanol, and transferred to 2 cm.² polythene disks, and dried and its radioactivity determined. The valine was next eluted from the disks with 3 ml. of water and the amount determined by the ninhydrin method (Jacobs, 1956). Similar-sized pieces of blank chromatographic paper were treated in the same way, and the valine values corrected for the traces of ninhydrin-positive material present in the paper. Specific radioactivities were expressed as counts/min./100 μ g. of valine. The error in this method was found to be about 15% by carrying out determinations on valine with known specific radioactivity.

Intracellular free valine was extracted within 5 sec. of stopping the perfusion by freezing the lung in liquid air, and weighing it and grinding it with 2-3 ml. of 20% (w/v) trichloroacetic acid for each gram of tissue. Isolation of the valine from the trichloroacetic acid extracts and determination of its radioactivity were carried out as described above. [¹⁴C]Valine was isolated also from the intracellular- and extracellular-antibody precipitates after hydrolysis in 6N-HCl for 24 hr. at 100° by the method described above.

Radioactivity determinations. ¹²¹I radioactivities were determined in a well-type scintillation counter, with 3 ml. samples of liquids, or with a known weight of dried protein in the form of a pellet at the bottom of a small pointed tube for antibody or non-specific γ -globulin precipitates. The counter efficiency was about 25% in each case, and the background was about 400 counts/min.

[¹⁴C]Amino acids were plated on 2 cm.² polythene disks and counted at infinite thinness under a Geiger-Müller counter with a mica end-window (1.7 mg./cm.²). [¹⁴C]-Proteins, in the absence of ¹²¹I, and BaCO₃ were counted at infinite thickness on 0.3 or 1 cm.² disks, and their specific radioactivities expressed as counts/min./cm.².

When ¹²¹I was also present, the radioactivity due to ¹⁴C was measured in a proportional counter after combustion to CO₂ (Bradley, Holloway & McFarlane, 1954). Antibody specific radioactivities were not corrected for the presence of antigen, which formed 4 and 10% (w/w) of the C in precipitates of antibodies to pneumococci and ovalbumin respectively. In any one experiment, only specific radio-

activities obtained by a given method were compared with one another.

Measurement of γ -globulin breakdown during perfusion. The perfusates collected at different time intervals and a sample of the rabbit's serum were pressure-dialysed overnight against 400 ml. of phosphate saline buffer at 3°. The diffusates were evaporated to a small volume, after adjustment of the pH to about 9.5, and the ¹²¹I content of the diffusible and non-diffusible material was determined. Increases in the percentage of diffusible ¹²¹I above that found in the serum sample were taken to represent protein breakdown.

RESULTS

Synthesis of antibodies and other γ -globulin. Two typical experiments are described. In Expt. 1 the lung was obtained from a rabbit hyperimmunized by intravenous injections of formalin-killed pneumococci type 3, the last injection having been given 2 days before death. A few minutes after the last injection of antigen 100 μ C of ¹²¹I-labelled rabbit γ -globulin (75% antibody specifically precipitable with capsular polysaccharide) was also administered to enable us later to recognize preformed extracellular antibody by its ¹²¹I-label.

The lung, which was three to four times normal weight, was perfused with unlabelled medium for 30 min.; the fluid was then collected and replaced with a fresh lot containing 2 μ C of L-[¹⁴C]lysine; this process was repeated after 120, 195 and 290 min. After 190 min. one lobe of the lung was tied off and removed for extraction of antibody, and the remainder of the lung was extracted at the end of the experiment. Antibody was also isolated from the perfusates, and the specific radioactivities due to ¹²¹I and ¹⁴C were measured. Table 1 shows the weights of antibody actually isolated, and the proportion of preformed extracellular antibody. It is striking that antibody isolated after perfusion for

Table 1. Amount of intracellular and extracellular antibody isolated during perfusion of lung from a rabbit hyperimmunized against pneumococci type 3

Two days before killing, 100 μ C of [¹²¹I]-labelled globulin concentrate containing 75% of specifically precipitable antibody was injected intravenously. The percentage of extracellular antibody is calculated by comparing the ¹²¹I specific activity of antibody samples with that of the serum antibody at death. Antibody content of serum at death: 43 mg./ml. One lobe of lung was removed at 195 min.

	Time of perfusion (min.)	Amount of antibody precipitate isolated (mg.)	Extracellular antibody (%)	Amount of antibody secreted by lung (mg.)
Perfusates {	0-30	13	83	2.2
	30-120	24.6	56	10.8
	120-195	17.8	48	9.3
	195-290	13.9	40	11.2*
	Total	69.3		33.5
Lung lobe (14 g.) at 195 min.		<14	1.6	—
Remainder of lung (38 g.) at 290 min.		29	1.9	—

* Corrected for lobe removed at 195 min.

4 hr. still contained 40 % of preformed extracellular antibody. A total of 69.3 mg. of antibody was isolated during perfusion for 5 hr., and of this 33.5 mg. was newly secreted by the lung. The rate of secretion (about 7 mg./hr.) was uniform over the whole period of perfusion (Fig. 2). This is probably an underestimate since some losses are apt to occur during pressure dialysis and isolation of the antibody.

The ^{14}C specific radioactivity of newly secreted antibody in the perfusate increased steadily after the first 90 min. Estimations at earlier times in several other experiments showed that ^{14}C never appeared in secreted antibody in less than 30 min., and significant labelling usually did not occur for up to 1 hr.

γ -Globulin was also isolated, though not quantitatively, from the perfusion fluids after removal of all precipitable antibody. Each lot of medium contained about 60 mg. of γ -globulin from the added normal serum, and any γ -globulin secreted by the lung was therefore diluted to an unknown extent. However, since considerable incorporation of ^{14}C lysine occurred, the lung must also have been synthesizing and secreting γ -globulin other than specific antibody. If the specific radioactivity of secreted γ -globulin is assumed to be equal to that of antibody secreted at the same time, the amount secreted during the second perfusion period

(between 120 and 195 min.) would have been 15 mg. During this period 9.3 mg. of antibody was also secreted, making a secretion rate of about 18 mg. of total γ -globulin/hr. The nature of such non-specific γ -globulin has been discussed by Askonas & Humphrey (1958).

In this experiment the antibody extracted from the lung after 190 and 290 min. had almost the same ^{14}C specific radioactivity as antibody isolated from the perfusing fluid at the same time. However, this was not always true, as is shown in the next experiment.

In Expt. 2 the rabbit had received eleven intravenous injections of pneumococci and eight of alum-precipitated ovalbumin during the preceding 4 weeks. Serum-antibody levels had declined by about 30 % during the last week, indicating that antibody synthesis had slowed down markedly. At death the serum contained 14 mg. of anti-capsular polysaccharide and 2 mg. of antiovalbumin/ml. The lungs were greatly enlarged.

Perfusion was carried out as above with medium containing $8\text{ }\mu\text{C}$ of ^{14}C glycine, and the two antibodies were isolated successively by specific precipitation with antigen. The results are summarized in Fig. 3. The two antibodies had similar specific radioactivities, although the amount of antibody against capsular polysaccharide secreted in 5 hr.

Fig. 2. Synthesis of antibody against pneumococci type 3 and incorporation of L- ^{14}C lysine into antibody and other γ -globulin by lung from hyperimmune rabbit. The lung was perfused according to the schedule in Table 1; arrows indicate the times of addition of complete medium containing $2\text{ }\mu\text{C}$ of L- ^{14}C lysine. Radioactivity due to ^{14}C was determined with a proportional gas counter and expressed as counts/min./mg. of C. The radioactivity measurements were corrected for preformed extracellular antibody. x, Antibody precipitate (mg.) isolated from perfusate (corrected for preformed extracellular antibody); O, ^{14}C specific radioactivity of antibody precipitate; Δ , ^{14}C specific radioactivity of residual γ -globulin isolated from perfusates; $\bullet$, specific radioactivity of intracellular antibody from lung after 195 and 290 min.

Fig. 3. Synthesis of antibody and incorporation of ^{14}C -glycine during perfusion of lung from rabbit hyper-immunized against pneumococci type 3 and ovalbumin. Arrows indicate the times at which fresh lots of complete medium containing $8\text{ }\mu\text{C}$ of ^{14}C glycine were added. Radioactivity due to ^{14}C was determined with a proportional gas counter and expressed as counts/min./mg. of C. The radioactivity measurements were corrected for preformed extracellular antibody. x, Antibody (mg.) against capsular polysaccharide isolated from perfusates (corrected for preformed extracellular antibody); O, ^{14}C specific radioactivity of antibody against capsular polysaccharide; $\bullet$, ^{14}C specific radioactivity of intracellular antibody against pneumococcal capsular polysaccharide; $\square$, ^{14}C specific radioactivity of antiovalbumin; $\blacksquare$, ^{14}C specific radioactivity of intracellular antiovalbumin; Δ , mg. of antiovalbumin (cumulative) isolated from perfusing fluids.

Table 2. *Comparison of specific radioactivities of secreted antibody and of antibody isolated from the lungs after perfusion with [¹⁴C]amino acids*

Specific radioactivity of antibody was determined by gas counting, and the values were corrected for the presence of preformed extracellular antibody

Expt.	Radioactive amino acid used	Perfusion time (min.)	Specific radioactivity (counts/min./mg. of C)		Total antibody secreted (mg.)	Wt. of lung after perfusion (g.)	Serum-antibody content (mg./ml.)	
			Antibody from perfusate	Antibody from lung			At death	Week before death
A	Glycine (7 μ C)	230	39 000	8245	—	—	7.2	12
		340	78 000	8200	7.2	38	—	—
B	L-Lysine (1.75 μ C)	340	10 200	9700	4.5	20	4.9	5
C	L-Lysine (1.25 μ C)	325	5 000	4170	5.2	20	4.3	2.9
D	L-Lysine (1.75 μ C)	300	10 200	5850	12	31	10	12.2
E	Glycine (8 μ C)	210	9 300	957	—	—	—	—
		300	10 400	1562	43	74	14.6	19.7
F	L-Lysine (2 μ C)	195	5 200	4400	—	—	—	—
		290	10 000	9850	33.5	52	43	29

(43 mg.) was much more than that of antiovalbumin (approx. 3.4 mg.). Forty-eight hours before death the rabbit had received ¹³¹I-labelled anti-capsular polysaccharide, but no [¹³¹I]antiovalbumin. The radioactivities of both antibodies were corrected for the presence of preformed extracellular antibody on the basis of measurement of ¹³¹I-labelled antibody against capsular polysaccharide. With antiovalbumin the correction is only approximate because the relative proportion of the two antibodies secreted by the lung is not necessarily the same as their proportion in serum.

In this experiment the ¹⁴C specific radioactivities of both types of intracellular antibodies were only 11–17 % of the specific radioactivities of the antibody secreted into the perfusion fluid at the corresponding times.

Comparison of specific radioactivity of intracellular and secreted antibody

Six typical experiments are summarized in Table 2, and show how the findings varied when we compared the specific radioactivities of intracellular antibody (taken at the end of the experiment) and of antibody secreted during the last perfusion period. In each case the lung perfused was from a rabbit immunized by intravenous injections of pneumococci. In half of the experiments the specific radioactivities of the intracellular and secreted antibody were very similar. In the other half intracellular antibody had a very much lower specific radioactivity, 10–50 % of that of the antibody secreted into the perfusate; in these experiments the antibody content of the serum was declining during the days before the experiment. A possible interpretation of these findings will be discussed below.

Table 3. *Specific radioactivity of free valine during perfusion*

Fresh lots of perfusing medium were added after 85, 162 and 240 min.

Material	Time of perfusion (min.)	Specific radioactivity of isolated valine (counts/min./100 μ g.)	Relative radioactivity of free valine (%)
Medium*	0	8630	—
Perfusate*	10	4870	—
Perfusate*	85	1900	—
Perfusate*	95	5550	—
Perfusate*	162	5300	—
Perfusate*	172	7820	—
Perfusate*	240	5640	—
Perfusate*	250	6840	—
Perfusate*	330	6125	100
Lung*	330	3500	57
Antibody† (in perfusate)	240–330	550	9
Intracellular lung antibody†	330	630	10

* Free valine was isolated from trichloroacetic acid filtrates.

† Valine was isolated from antibody hydrolysates.

Radioactivity of intracellular and extracellular free valine and antibody valine during perfusion

L-[¹⁴C]Valine was chosen for this study because valine is not converted into any other amino acid by the cells and is relatively easy to isolate. The lung of an immune rabbit was perfused with complete medium (containing 1.8 μ C of L-[¹⁴C]-valine); this was replaced by fresh medium after 85, 162 and 240 min. Free [¹⁴C]valine was isolated from 10 % trichloroacetic acid filtrates of the perfusates at the beginning and end of each perfusion period, and from the lung at the end of the experiment. The results may be seen in Table 3. During

the first 10 min. the specific radioactivity of the free valine fell from 8630 counts/min./100 μ g. of valine to 4870, by almost half its original value. Since the medium contained 3.4 mg. of L-valine, the valine in the medium must have been diluted by an almost equal amount of intracellular free valine from the 27 g. of lung. Lung contains about 14% of protein (Spector, 1956); if we assume the protein to contain 5% of valine, an intracellular pool of almost 3.5 mg. would represent 2% of the total protein-bound valine. As perfusion proceeded, the specific radioactivity of the valine fell further, and after 85 min. was one-fifth of the original value; this suggests continuous breakdown of unlabelled lung protein to free amino acid in the cells, but only a small percentage of the total lung protein can be involved. When the medium was replaced after 85, 162 and 240 min. with fresh medium containing valine with a specific activity of 8630 counts/min./100 μ g., the specific radioactivity of the valine fell during each perfusion period to about two-thirds of its original level. Hence a small amount of breakdown of lung protein to freely exchangeable valine continues during the entire period of perfusion. The use of Pontamine Sky Blue in the medium showed that all of the lung was being perfused. The intracellular free valine of the lung at the end of the experiment had a specific radioactivity of 3500, i.e. 57% of the radioactivity of the extracellular free valine at the same time.

[14 C]Valine was also isolated from the antibody secreted by the lung (corrected for the presence of extracellular 131 I-labelled antibody) and from intracellular lung antibody (Table 3). It was found to have a specific radioactivity about one-fifth of that of the intracellular free valine isolated from the lung at the end of the experiment. The radioactivity of the total lung protein, precipitated with trichloroacetic acid, was also determined; over 18% of the L-[14 C]valine added to the perfusing fluid was incorporated into the protein of the lung. More than half of this radioactivity was associated with the protein readily extractable with isotonic NaCl from the lung after breaking the cells.

Two other confirmatory experiments were carried out, one with L-[14 C]valine and the other with amino acids obtained by acid hydrolysis of algal [14 C]protein. In the latter experiment the specific radioactivity of the α -amino carboxyl group was determined as BaCO₃ after release of the CO₂ by ninhydrin (Van Slyke, McFadyen & Hamilton, 1941), as described by Humphrey & Sulitzeanu (1958). In both these experiments, similar decreases in specific radioactivity of the free amino acid during the perfusion period were observed. As in the experiment described above, the intracellular free amino acid in the lung had a radioactivity about two-thirds of that of the extracellular amino

acid. Although the specific radioactivity of the antibody secreted into the perfusate during the last hour was almost the same as that of the intracellular antibody, the specific radioactivity of the 14 C of the α -amino carboxyl group of the constituent amino acids was less than one-fifth of that of the free amino acids in the lung.

Breakdown of 131 I-labelled proteins by the lung during perfusion

When 14 C- and 131 I-labelled proteins of the same kind are administered simultaneously to animals, they are catabolized at practically identical rates (Cohen, Holloway, Matthews & McFarlane, 1956; McFarlane, 1956). Perfusion of 131 I-labelled proteins through the isolated liver results in splitting off of non-protein iodine only when protein breakdown occurs (Gordon, 1957). There are good grounds therefore for considering that liberation of non-protein iodine by the lung from 131 I-labelled plasma protein indicates catabolism of the proteins. In our experiments the rabbits had received 131 I-labelled γ -globulin 48 hr. before the lung perfusion, and a small fraction of the dose was retained in the interstitial fluids of the lung. During the perfusion part of the 131 I was split off, the proportion of diffusible iodine in the perfusates being three to five times greater than that in the serum of the same rabbit taken at the time of death. This implies that γ -globulin was being broken down and that, if there was complete mixing of 131 I-labelled γ -globulin and the serum proteins in the perfusates, the lung would have been breaking down during each hour about 5% of the γ -globulin in the medium (about 70 mg.), i.e. up to 3.5 mg. of γ -globulin/hr. To confirm this finding, the lung from a hyperimmunized rabbit, which had not received 131 I-labelled protein *in vivo*, was perfused with medium containing 0.1 vol. of normal serum to which had been added a few milligrams of 131 I-labelled γ -globulin. Under these

Table 4. *Catabolism of 131 I-labelled γ -globulin by hyperimmunized lung*

The lung of a hyperimmunized rabbit was perfused with 60 ml. of medium containing 6 ml. of normal rabbit serum and 3 mg. of rabbit γ -[131 I]globulin. The medium was changed after 35, 110, 190 and 290 min. (All counts were at least three times the background count.)

Perfusion period (min.)	Diffusible 131 I in perfusate (%)	Recovery of 131 I counts in perfusate (%)
Original medium	2.3	—
0-35	5.5	80
35-110	5.8	97
110-190	6.2	78
190-290	5.9	100

conditions also splitting off of non-protein ^{131}I occurred and to a similar extent (Table 4). Since only γ -globulin was labelled it is not possible to state whether lung also catabolized other plasma proteins.

DISCUSSION

Lungs from hyperimmunized rabbits have been perfused and have been shown to synthesize and secrete specific antibody and other γ -globulin at a rate comparable with that found for plasma proteins by livers perfused with blood (Miller *et al.* 1951).

Although such lungs are not obtained as readily as normal livers, they have one great advantage, namely that antibody can easily be isolated from the perfusion medium, and that accurate allowance can be made for any preformed extracellular protein which was present at the beginning of the experiment. Such correction is necessary since about 40% of the antibody appearing in the perfusate was not newly secreted even after perfusing for 2–3 hr. We have calculated the efficiency of conversion of the labelled valine into antibody by the lung in an experiment in which the levels of free valine were determined in all samples of medium. During this experiment about 30 mg. of antibody was newly formed in 5 hr. A total of $4\mu\text{C}$ of valine was added. Of this $0.92\mu\text{C}$ was recovered as free valine in the samples of medium removed and $0.072\mu\text{C}$ was recovered in antibody isolated from the perfusates and from the lung at the end of the perfusion. Thus of the $3.08\mu\text{C}$ which apparently entered the lung tissue, $0.07\mu\text{C}$ (about 2.3%) was converted into antibody. If we assume that the amount of other γ -globulin formed was 1.5 times the amount of antibody (see Expt. 1 above), the total converted into secretable globulins was 5.7%. This is almost certainly an underestimate, since the recovery of free valine and of antibody was probably not complete. In intact rabbits the percentage of administered ^{14}C amino acids which are incorporated into plasma proteins varies between about 3 and 9%, depending on which amino acids are used (J. H. Humphrey & A. S. McFarlane, unpublished work).

The radioactivity found in the lung protein (after extraction with trichloroacetic acid at 100° and with boiling ethanol) at the end of the experiment was 18 and 10% of the total dose when lungs weighing 27 and 15 g. respectively were perfused. About half of this radioactivity was in the tissue protein insoluble in 0.9% sodium chloride solution.

Delayed appearance of label in secreted antibody

A consistent finding, both in these experiments and those with tissue slices (Askonas & Humphrey, 1958), was a definite delay between addition of

^{14}C amino acid to the medium and its appearance in significant amounts in secreted antibody, although secretion of antibody into the medium proceeded linearly from the start. With perfused lungs this delay was half an hour or more and can hardly be attributed to time needed for the tissue to recover from the manipulations involved in setting up the perfusion, since the lung was perfused with complete medium for half an hour before labelled amino acids were added. Two possible explanations may be offered: antibody may be formed from some precursor (be it protein or bound amino acids) or it may be synthesized from an amino acid pool in equilibrium with the extracellular free amino acids, but may be bound in some way when first formed. The second is the explanation preferred by Peters (1957), who reported that in the liver albumin extracted from disrupted microsomes becomes labelled most rapidly and without appreciable lag period. Microsomal fractions isolated from spleen or bone marrow of immunized rabbits do contain some bound antibody extractable with carbonate buffer, pH 9.5, which disrupts the microsome structure (Askonas, 1958). It is therefore likely that antibody is first formed bound to subcellular particles, before it becomes freely extractable with isotonic medium. There is another lag period of about 30 min. between the appearance of radioactive free intracellular antibody and its appearance in the medium during incubation of sliced tissues (Askonas & Humphrey, 1958). The same is true *in vivo*. Whereas radioactive antibody can be extracted from tissues within 15 min. after injection of ^{14}C amino acids, radioactive antibody appears in the blood stream only after 25 min. (Humphrey & Sulitzeanu, 1958). Hence it must take some time for the newly formed antibody to mix with the pool from which secretion occurs, or for the process of secretion, the transport of the protein molecules across the cell membranes, to take place.

Specific activity of free amino acids during perfusion

It is evident from Table 3 that at each stage of the perfusion the L- ^{14}C valine of the medium was diluted with some inactive L-valine, presumably derived from the breakdown of lung protein (and possibly extracellular protein) to free amino acids. However, at the end of the experiment, the specific radioactivities of extracellular and intracellular valine were not identical, even though the tissue was immediately frozen in liquid air to prevent further changes. This suggests either that unlabelled free valine was continually being produced within the cells too fast for equilibrium to be reached with that outside, or that there must be some bound unlabelled intracellular valine not

readily exchangeable with the extracellular valine but easily extracted by the method used.

Since free valine refers to valine extractable with 10% trichloroacetic acid, it is possible that bound valine is released by the trichloroacetic acid and that a mixture of free and bound valine within the cells is determined. In each of three experiments, two with L-[¹⁴C]valine and one with algal [¹⁴C]-protein hydrolysate, the radioactivity of the intracellular amino acids was about 60% of that in the perfusion fluid. These findings are in accord with some observations reported in the literature. Loftfield & Harris (1956) administered 225 μ moles of L-[¹⁴C]leucine to rats by repeated injections at short intervals of time, and found that intracellular leucine in the liver attained only 40% of the radioactivity of the leucine injected (the radioactivity of the blood leucine was not actually determined). Steiner & Anker (1956), using tissue cultures of spleen cells from an immunized donor in a medium containing a relatively large quantity of [¹⁴C]-glycine, found that only about one-half of the glycine residues of the antibody were derived from added glycine after 48 hr. Considerations of this kind clearly must be borne in mind when attempts are made to calculate turnover rates of proteins by measuring incorporations of labelled amino acids.

Relative specific activities of intracellular antibody and free amino acids

It is evident from Table 3 that after perfusion for 5 hr. (1½ hr. after the last change of medium) the specific radioactivity of valine in the secreted antibody, and the antibody remaining in the lung cells, was only about one-sixth of that of intracellular free valine. Similar findings were obtained in two other experiments. If antibody was being synthesized from the intracellular free amino acid pool we examined, this finding implies a rather long turnover time. On the assumption that the mean specific activities of intracellular valine during successive perfusion periods of the experiment described in Table 3 were 800, 2500, 3100 and 3400 counts/min./100 μ g. respectively; and from equation (3) derived by Humphrey & Sulitzeanu (1958), integrated into the form $y = x - ae^{-t/K}$, the turnover time K (time taken for cells to renew their antibody content) in this experiment can be calculated to be about 14 hr. This is at variance with our finding that the amount of antibody secreted in 5 hr. was approximately equal to that extractable from the lung at the end of an experiment, implying a turnover time of about 5 hr., and with the turnover time of 2–3 hr. deduced by Humphrey & Sulitzeanu (1958) for conditions *in vivo*.

Several possible explanations for the discrepancy may be suggested. The measured radioactivity of intracellular free amino acids may not represent

those in the cells actually synthesizing antibody, and may be too high; there may be cells present which continue to secrete antibody, but which are no longer engaged in active synthesis, so that labelled antibody is thereby diluted with unlabelled; our methods of extraction may have led to serious (two- to three-fold) underestimation of the amounts of intracellular antibody present; or our theoretical assumptions may be erroneous. Until a homogeneous preparation of cells synthesizing specific antibody or other γ -globulin becomes available there appears to be no way of deciding between these possibilities. That the turnover time should prove longer in an isolated lung, perfused without blood, than it is *in vivo* is scarcely surprising. Nevertheless, it is possible that the estimate of 2–3 hr. quoted above, which involved numerous approximations, is too short.

Mechanism of antibody secretion

If the intracellular antibody constitutes a pool from which secretion occurs into the medium, the specific radioactivities of newly secreted and of intracellular antibody at any time should be equal. It was not possible to measure instantaneously the radioactivity of antibody as it was secreted, and our experimental values for antibody in the medium represent average values for a perfusion period of 1–1½ hr., whereas those for intracellular antibody refer to antibody extracted at the end of the period. The two are therefore not strictly comparable, especially since equilibrium between the radioactive amino acids in the medium and in the cells was never completely attained. Nevertheless a comparison is of interest.

Table 2 summarizes six typical experiments. In three the specific radioactivity of intracellular antibody is almost equal to that of secreted antibody, and in three it is much lower. In our view, those experiments in which secreted and intracellular antibody had similar radioactivities at the end of the experiment support the idea that liberation of antibody into the medium can be described in terms of secretion from the total intracellular pool. In those experiments where the specific radioactivity of intracellular antibody was much less than that of secreted antibody, metabolically inert antibody must also have been present. There was nothing to suggest that this could be due to failure to perfuse part of the lung. However, a possible explanation may be that some cells contained stored antibody which was not continuously secreted.

The lungs in which the specific radioactivity of intracellular antibody was unexpectedly low all came from rabbits whose serum-antibody levels had begun to fall, despite continued injections of antigen. A section of the lung from such a rabbit,

stained to demonstrate antibody by the sandwich technique of Coons, Leduc & Connolly (1955) is shown in Fig. 4. In this lung there were many cells containing antibody in discrete granules, as well as cells with antibody uniformly distributed throughout the cytoplasm, and it seems reasonable to suppose that the antibody in granular form was metabolically distinct and relatively inert. We have not excluded the possibility that a similar condition may obtain to some extent in the lungs of

all hyperimmunized rabbits, but in those which have not been overimmunized the granular form of antibody is comparatively unimportant.

Breakdown of globulin by immune lungs

The fact that hyperimmunized lungs were able to split iodide from ^{131}I -labelled γ -globulin is of some interest, since it is uncertain where plasma-protein breakdown occurs. Such lungs contain large numbers of macrophages, and these may be the cells responsible; if this is so, normal lungs would be less active, since their macrophage population is smaller. However, the maximum rate of breakdown which could be inferred in the experiment illustrated was 84 mg. in 24 hr., when the lung was perfused with serum diluted 1:10 in the medium. The level of total γ -globulin in the serum of this rabbit was about 50 mg./ml. If we assume that 10% of the 12 g. of total extracellular γ -globulin is broken down daily, the contribution to overall catabolism by the lung would be only 7%, unless catabolism were more rapid in the presence of higher concentrations of plasma protein. Gordon (1957) estimates that in the rat the liver is responsible for not more than 10% of plasma-albumin catabolism.

SUMMARY

1. Isolated lungs from hyperimmune rabbits, perfused *in vitro* with a balanced-salt solution containing amino acids and 10% of normal serum, secreted antibody into the medium at rates up to 8 mg./hr.

Allowance was made for extracellular antibody by injecting ^{131}I -labelled antibody into the rabbits beforehand, so that at the time of death extracellular antibody could be identified by its content of ^{131}I .

2. ^{14}C -Labelled amino acids added to the medium are incorporated into both specific antibody and other γ -globulin. The rate of synthesis of other γ -globulin was calculated to be rather greater than that of specific antibody.

3. A delay of more than 30 min. was observed between the addition of labelled amino acids and the appearance of labelled antibody in the medium. Antibody was extracted from the lung at the end of the experiments, and its radioactivity was compared with that of antibody secreted into the medium during the period immediately preceding. In some experiments the radioactivity of intracellular antibody was equal to, and in others it was less than, that of secreted antibody.

4. It is suggested that newly formed antibody mixes with an intracellular pool, from which secretion occurs into the medium. If this is so, antibody must first be synthesized in a bound form, and some lungs must contain antibody which is relatively inert metabolically.

(a)

(b)

Fig. 4. Section ($\times 360$) of lung from hyperimmunized rabbit whose plasma antibody level was falling, stained (a) by fluorescent antibody to show antibody-containing cells, and (b) the same field stained with haematoxylin and eosin. Note the varied appearance of antibody within the cells surrounding the arteriole.

5. The specific radioactivity of [^{14}C]amino acid in the perfusion fluid was measured at various times. The specific radioactivity of the intracellular free amino acids (L-valine or total amino acids), after perfusing for 5 hr., was about 60 % of that of the corresponding amino acids in the medium at that time. The specific activity of the amino acids in the intracellular antibody was only one-sixth of that of the free amino acids on the intracellular pool. The interpretation of these findings is discussed.

6. Lungs from hyperimmune rabbits can liberate diffusible iodide from ^{131}I -labelled globulin.

We wish to thank Mr W. E. Brocklehurst for advice in the design of the perfusion apparatus, Mr R. C. Holloway for performing the iodinations and Mr G. Dickinson and Mr G. Savage for the gas counting.

It is a pleasure to acknowledge the excellent technical assistance of Miss Marion T. Smith and Miss Rosemary Willson.

REFERENCES

- Askonas, B. A. (1958). *Rec. Trav. chim. Pays-Bas*. (In the Press.)
- Askonas, B. A. & Humphrey, J. H. (1958). *Biochem. J.* **68**, 252.
- Bradley, J. E. S., Holloway, R. C. & McFarlane, A. S. (1954). *Biochem. J.* **57**, 192.
- Cohen, S., Holloway, R. C., Matthews, C. M. & McFarlane, A. S. (1956). *Biochem. J.* **62**, 143.
- Coons, A. H., Leduc, E. H. & Connolly, J. M. (1955). *J. exp. Med.* **102**, 49.
- Gordon, A. H. (1957). *Biochem. J.* **66**, 255.
- Hanks, J. H. (1948). *J. cell. comp. Physiol.* **31**, 235.
- Heidelberger, M., Kendall, F. E. & Scherp, H. W. (1936). *J. exp. Med.* **64**, 559.
- Humphrey, J. H. & Sulitzeanu, B. D. (1958). *Biochem. J.* **68**, 146.
- Jacobs, S. (1956). *Analyst*, **81**, 502.
- Loftfield, R. B. & Harris, A. (1956). *J. biol. Chem.* **219**, 151.
- McFarlane, A. S. (1956). *Biochem. J.* **62**, 135.
- Miller, L. L., Bly, C. G., Watson, M. L. & Bale, W. F. (1951). *J. exp. Med.* **94**, 431.
- Peters, T. (1957). *Fed. Proc.* **16**, 369.
- Spector, W. S. Ed. (1956). *Handbook of Biological Data*, p. 71. London: W. B. Saunders Co. Ltd.
- Steiner, D. & Anker, H. (1956). *Proc. nat. Acad. Sci., Wash.*, **42**, 580.
- Van Slyke, D. D., McFadyen, D. A. & Hamilton, P. (1941). *J. biol. Chem.* **141**, 671.

Transphosphorylations Catalysed by Bivalent Metal Ions

By J. M. LOWENSTEIN*

Department of Biochemistry, University of Oxford

(Received 7 March 1958)

In all enzyme reactions involving the breakdown or the formation of the polyphosphate chain of nucleoside triphosphates there is a requirement for bivalent metal ions. This requirement is absolute in most, if not all, cases. It is most commonly met by Mg^{2+} ions, but these can frequently be replaced by Mn^{2+} ions. A few enzyme reactions which show a requirement for a bivalent metal ion also show an additional requirement for a univalent metal ion. In the examples which have come to light so far, this additional requirement is met by potassium (Hers, 1952; Parks, Ben-Gershom & Lardy, 1957; Webster & Varner, 1954a). Of several reviews the one by Lardy (1951) is the most relevant to the present paper. It also gives references to earlier reviews.

The work presented here arose from a study of enzyme-catalysed exchange reactions. In the course of an experiment designed to test for an enzyme-catalysed exchange between radioactive orthophosphate and adenosine triphosphate, the disappearance of radioactive orthophosphate was observed in controls which contained no enzyme.

* Beit Memorial Fellow.

The radioactive compound formed from orthophosphate was identified as pyrophosphate (Lowenstein, 1957a). The reaction was shown to depend on the presence of bivalent metal ions and adenosine triphosphate (ATP), and was formulated to proceed according to equation (1) (ADP represents adenosine diphosphate).

This paper presents a detailed account and an extension of these observations.

A test of acceptors other than phosphates has shown that acylphosphate formation from nucleoside triphosphate and carboxylic acids also occurs in the absence of enzymes, and that it is catalysed by bivalent metal ions (Lowenstein, 1958).

EXPERIMENTAL

Materials and reagents

Nucleotides. The sodium salts of ATP and ADP were products of the Sigma Chemical Co., St Louis, Mo., U.S.A. The free acid of adenosine monophosphate was obtained