The alkaline-quenching has been found to be a particular case of the quenching of fluorescence by transfer of a H^+ ion from a nitrogen atom belonging to an excited molecule to the OH^- ion (G. Weber & A. White, unpublished work). Such cases were first observed by Boaz & Rollefson (1950). The necessity of a mobile hydrogen atom in NH for this reaction to occur has been checked by observing the lack of quenching of fluorescence of *N*-methyl indole by OH^- ions in 5*N*-sodium hydroxide. The reactions may be represented as:

The reaction $R^* \cdot N^- \rightarrow R \cdot N^-$ is either radiationless (α -naphthylamine) or radiative with low efficiency (β -naphthylamine) (Boaz & Rollefson, 1950). In indole derivatives in *N*-sodium hydroxide no detectable fluorescence has been found to be emitted at any wavelengths.

SUMMARY

1. The influence of pH on the fluorescence of tyrosine, tryptophan and related compounds was investigated.

2. The pH of acidic half-quenching of fluorescence (pH_f) of tyrosine and tryptophan and their derivatives with an intact carboxyl group corresponds approximately to the pK of the carboxyl and falls to a much lower value in those derivatives where the carboxyl group is substituted or replaced.

3. Tyramine, tyrosine and phenol are converted into non-fluorescent species on dissociation of the phenolic hydroxyl group, the quenching of the fluorescence appearing as a sensitive test of this dissociation as shown by the effect of substituents, hydrogen-bonding and chemical substitution of the hydrogen.

4. The existence of a peak in the alkaline pH range is described for two tyrosine and four tryptophan derivatives. A free amino group is apparently a necessary but not a sufficient condition for this maximum to be present.

5. The relative quantum yield in neutral aqueous solution and at a chosen wavelength was measured for the compounds mentioned.

6. The fluorescence-excitation spectra of the majority of the derivatives were determined in neutral and alkaline solutions and found to correspond with the fractional absorption spectra at neutrality.

The author wishes to thank Dr G. Weber for the initiation of this work and his continued encouragement, and Dr F. W. J. Teale for helpful discussion. The financial help of the Medical Research Council and the Medical Research Fund of the University of Sheffield is gratefully acknowledged.

REFERENCES

- Boaz, H. & Rollefson, G. K. (1950). *J. Amer. chem. Soc.* **72**, 3435.
 Bowen, E. J., Barnes, A. W. & Holliday, P. (1947). *Trans. Faraday Soc.* **43**, 27.
 Brenner, M. & Huber, W. (1953). *Helv. chim. acta*, **36**, 1109.
 Crammer, J. L. & Neuberger, A. (1943). *Biochem. J.* **37**, 302.
 du Vigneaud, V. & Meyer, C. E. (1932). *J. biol. Chem.* **98**, 295.
 du Vigneaud, V. & Sealock, R. R. (1932). *J. biol. Chem.* **96**, 511.
 Greenstein, J. P. (1932). *J. biol. Chem.* **95**, 465.
 Katchalski, E. & Sela, M. (1953). *J. Amer. chem. Soc.* **75**, 5284.
 Tanford, C. (1950). *J. Amer. chem. Soc.* **72**, 441.
 Teale, F. W. J. & Weber, G. (1956). *Biochem. J.* **65**, 476.
 Umberger, J. Q. & la Mer, V. K. (1945). *J. Amer. chem. Soc.* **67**, 1099.
 Weber, G. & Teale, F. W. J. (1957). *Trans. Faraday Soc.* no. 413, **53**, part 5, 646.
 Weber, G. & Teale, F. W. J. (1958). *Trans. Faraday Soc.* no. 425, **54**, part 5, 640.
 Winnek, P. S. & Schmidt, C. L. A. (1935). *J. gen. Physiol.* **18**, 889.

The Nature of the Extra Protein Fraction from Myofibrils of Striated Muscle

By S. V. PERRY AND M. ZYDOWO*

Department of Biochemistry, University of Cambridge

(Received 14 July 1958)

When isolated myofibrils are extracted with solutions considered to be selective for myosin most of the material responsible for the difference in refractive index between the A and I bands

* Research Fellow of the Rockefeller Foundation, 1957-8. Present address: Department of Biochemistry, Medical Academy, Gdansk, Poland.

(A substance) is removed. Under these conditions Szent-Györgyi, Mazia & Szent-Györgyi (1955) have claimed that, in addition to the myosin, protein material amounting to approximately 20% of the total myofibrillar nitrogen passes into solution. Evidence has been presented by de Villafraña (1956) that this additional material, the so-called

extra protein fraction, consists mainly of a protein somewhat similar to tropomyosin in properties although not identical with it. Subsequent studies (Perry & Corsi, 1958; Hanson & Huxley, 1957) have indicated that extra protein represents a considerably smaller fraction of well-washed myofibrils than the original results of Szent-Györgyi *et al.* (1955) would suggest. Furthermore, recent electrophoretic evidence obtained by Corsi (1957) has shown that the extra protein fraction contains several components, of which he considers tropomyosin to be the most abundant when the myofibrils are extracted at lower pH values. Independent work in this Laboratory has also indicated that the extra protein fraction obtained from myofibrils isolated from fresh muscle is heterogeneous, and this paper is concerned with its chromatographic resolution into four main fractions, two of which do not appear hitherto to have been isolated from myofibrils. This investigation forms part of a general study of the muscle proteins by the methods of ion-exchange chromatography, with the aim of precise characterization of components present in the myofibril in addition to myosin, actin and tropomyosin.

METHODS

Chromatography. (2-)Diethylaminoethylcellulose was prepared from Solka-Floc BW. 200 (Brown Co., Boston, Mass., U.S.A.) by the method of Peterson & Sober (1956). Total N estimations carried out by the micro-Kjeldahl technique on each batch gave values ranging from 0.95 to 1.36%. Before use the diethylaminoethylcellulose was suspended in about 100 vol. of the buffer in which was dissolved the protein later to be applied to the column. After washing four times by centrifuging and resuspension in fresh buffer, the cellulose was poured into the chromatography column as a thick slurry, and packed in one operation by the application of 10 cm. Hg pressure. Finally the column was washed overnight or until the extinction at 280 m μ of the eluate was negligible and the pH identical with that of the buffer applied. Protein solutions were usually run into the column without pressure, but subsequent elution with buffer was carried out under a pressure of about 60 cm. water. Chromatography was carried out at 0° in all cases. When KCl gradients were applied to the column the total chloride concentration of the eluate was followed by titration with 0.02 N-AgNO₃ and potassium chromate as an internal indicator. As judged by absorption at 280 m μ about 60% of the applied material was recovered from the column when the ionic strength of the eluent was increased to about 2.0.

The diethylaminoethylcellulose was used repeatedly after regeneration by washing by centrifuging and resuspension four times with 1% NaOH followed by four times with water.

Preparation of myofibrils. Myofibrils were prepared from the back and leg muscles of the rabbit by the method already described (Perry & Grey, 1956) with the following modifications. (1) Initial homogenizations of the minced

muscle were carried out in 25 mM-KCl containing 39 mM-sodium borate buffer, pH 7.1 (Perry, 1953), but after removal of the myofibrillar layer from the sedimented whole homogenate all subsequent resuspensions and washings were carried out in 0.1 M-KCl containing 39 mM-sodium borate buffer. (2) Homogenization and resuspension of myofibrils was carried out in 900 ml. of suspension for each 100 g. of the original minced muscle. Once the crude myofibrils had been separated from the fibrous material and unbroken residue of the muscle they were washed eight times by centrifuging, instead of five times as in the earlier method (Perry & Grey, 1956). Thus the myofibrils from 300 g. of minced muscle were washed with 20–25 l. of buffer. Yield was about 600–800 ml. of an approx. 2% suspension.

Larger amounts of less well-washed myofibrils, so-called crude preparations, were obtained in a similar manner by homogenizing and resuspending the myofibrils in a total volume of 700 ml. of buffer/100 g. of minced muscle and restricting the washing to four long centrifugings in addition to the short centrifuging to remove coarse material. In this case, myofibrils from 300–400 g. of muscle were washed with about 10 l. of buffer.

Preparation of extra protein. Extra protein was prepared by extracting 1 vol. of a concentrated myofibril suspension with 3 vol. of a solution containing 0.47 M-KCl, 0.1 M-potassium phosphate buffer, pH 6.5, mM-MgCl₂ and 10 mM-sodium pyrophosphate (Hasselbach & Schneider, 1951). After standing for 20 min. at 0° the insoluble material was centrifuged for 30 min. at 600 g and the turbid supernatant dialysed against 0.04 M-KCl containing 6.7 mM-potassium phosphate buffer, pH 7.2, until precipitation was complete. The gelatinous precipitate was removed by centrifuging for 20 min. at 40 000 g, and the resulting clear supernatant containing extra protein ($E_{1\text{ cm.}}$ at 280 m μ usually about 0.6–0.7) concentrated by freeze-drying or salted out by saturating with ammonium sulphate.

Preparation of fractions II and IV. Fraction II was prepared from the extra protein in the following way. The dilute supernatant obtained after the (acto)myosin precipitation was adjusted to 0.1 M-KCl by the addition of solid KCl and the pH brought to 7.6 by the addition of m-2-amino-2-hydroxymethylpropane-1:3-diol (tris) solution. This solution of extra protein was run straight into a column of diethylaminoethylcellulose (10–15 g. for about 1 g. of extra protein) previously equilibrated against 0.02 M-tris HCl buffer, pH 7.6, containing 0.1 M-KCl. After the protein had been washed in the column with the latter buffer until $E_{1\text{ cm.}}$ at 280 m μ of the eluate reached a negligible value, fraction II was eluted by the stepwise increase of the concentration of the KCl in the buffer solution to 0.22 M. Globulin was isolated from this fraction by dialysis against water.

Fraction IV was likewise separated from the same sample of dilute extra protein solution by collecting the fraction eluted between 0.4 M-KCl and 1.5 M-KCl.

Preparation and extraction of muscle granules. Minced rabbit skeletal muscle was homogenized for 2 min. with 6 vol. of 0.1 M-KCl containing 39 mM-sodium borate buffer, pH 7.1. The homogenate was centrifuged for 20 min. at 600 g, the turbid supernatant removed carefully and filtered through a very thin layer of glass wool. This suspension was again centrifuged at 600 g for 20 min. to sediment any myofibrils remaining in suspension and the

granules were sedimented out of the supernatant by centrifuging for 2 hr. at 15 000 *g*. After resuspension in the original KCl-borate medium the granules were spun down again to remove the bulk of contaminating sarcoplasm and finally treated as for the extraction of extra protein (see above). After dialysis against 0.04 M-KCl containing 6.7 mM-phosphate buffer, pH 7.2, the precipitate was centrifuged off and the clear supernatant was dialysed against 0.1 M-KCl containing 0.02 M-tris HCl buffer, pH 7.6, and applied directly to the diethylaminoethylcellulose column.

Sarcoplasmic extracts. Freshly minced muscle was homogenized with 3 vol. of 0.039 M-sodium borate buffer, pH 7.1, containing 0.025 or 0.10 M-KCl. The bulk of the myofibrils was removed by centrifuging for 30 min. at 600 *g* and the supernatant subsequently re-centrifuged for 2 hr. at 15 000 *g* to remove granular material. The clear supernatant was dialysed against 0.02 M-tris HCl buffer, pH 7.6, containing 0.1 M-KCl and applied directly to the column.

Preparation of other muscle proteins. L-Myosin was prepared as described by Perry (1955), F-actin by the method of Straub (1943) as modified by Bailey & Perry (1947) and twice-crystallized tropomyosin by the method of Bailey (1948). These proteins were isolated from rabbit skeletal muscle in all cases.

Viscometry. Actomyosin formation was evaluated by determining the fall in viscosity obtained when adenosine triphosphate (ATP) was added to a final concentration of 0.8 mM to a solution containing L-myosin and the protein under investigation. The response was compared with that of F-actin. Viscosity estimations were carried out at 0° in a solution containing 0.5 M-KCl and 0.05 M-potassium phosphate buffer, pH 7.1.

Tropomyosin assays were also carried out viscometrically by a method described earlier (Perry, 1953; Perry & Corsi, 1958), which depended on the fall in viscosity obtained on the addition of KCl to a solution of tropomyosin in a buffer of low ionic strength.

Enzymic assays. All determinations were carried out at 25° with short incubation times (usually 5 min.). Adenosine triphosphatase (ATPase) measurements were carried out by the standard procedure used in this Laboratory (Perry & Grey, 1956), and aldolase assays under the experimental conditions described by Sibley & Lehninger (1949). ATPase activity was expressed in μ l. of P gas liberated in 1 hr. by 1 ml. of protein solution of $E_{1\text{ cm}}^{1\%}$ 1.0 at 280 $m\mu$. This is identical with the Q_p value (Bailey, 1942) when $E_{1\text{ cm}}^{1\%}$ was 10.0 at 280 $m\mu$.

5'-Adenylic deaminase determinations were carried out in a total volume of 2 ml. containing 0.05 M-sodium succinate buffer, pH 5.9, and 5 mM-adenosine monophosphate. The reaction was stopped with 1 ml. of 15% (w/v) trichloroacetic acid. After nesslerization the NH_3 present was determined by measuring the absorption at 490 $m\mu$ in the Beckmann spectrophotometer.

Materials. ATP and adenosine monophosphate (AMP) were supplied by Sigma Chemical Co. Fructose diphosphate was prepared as the dibarium salt by Mr D. Gardiner of this Department by a modification of the method of Neuberg & Lustig (1942). The nucleic acids used for purposes of comparison in the identification of fraction IV were commercial preparations (British Drug Houses Ltd.) of the sodium salts of yeast and thymus nucleic acid. This was recrystallized twice from hot ethanol before use.

RESULTS

Fractionation of extra protein

Extra protein from well-washed rabbit myofibrils was separated into four main fractions by chromatography on diethylaminoethylcellulose (Fig. 1). Fraction I, representing about 30% of the total absorption at 280 $m\mu$ by the eluted material, was not held by the cellulose when the extra protein dissolved in 0.1 M-KCl buffered with 0.02 M-tris,

Fig. 1. Chromatography of extra protein extracted from a crude preparation of rabbit myofibrils. Protein solution (46.5 ml.; $E_{1\text{ cm}}^{1\%}$ 9.69 at 280 $m\mu$) in 0.1 M-KCl containing 0.02 M-tris buffer, pH 7.6, was applied to a column 27 cm. $\times$ 2 cm. diam. Gradient to 0.35 M-KCl, 0.02 M-tris buffer at point marked A and step to 2.0 M-KCl, 0.02 M-tris buffer at B.

pH 7.6, was applied to a column which had been previously equilibrated against this solution. On slowly increasing the concentration of KCl in the buffer, two additional peaks (fractions II and III) were eluted, the maxima in each case corresponding to chloride concentrations of about 0.20 and 0.3 equiv./l. respectively. Fraction IV was obtained as a sharp peak on subsequent stepwise or gradient application of M-KCl buffered with 0.02M-tris, pH 7.6.

Although the myofibrils used in the earlier stages of this work were exhaustively washed to remove the soluble sarcoplasmic proteins it was thought possible that the preparations might still contain proteins of this type. This view was confirmed by enzymic studies (see below) on repeatedly washed myofibrils, which showed that such preparations contained aldolase, an enzyme normally associated with the sarcoplasm. For purposes of comparison, sarcoplasmic extracts made in buffer solutions identical with those used for the myofibril preparations and freed from particulate material (see Methods) were chromatographed on diethylaminoethylcellulose under the same conditions as those used for fractionation of the extra protein. A typical result is illustrated in Fig. 2. Relatively little of the sarcoplasmic protein was held on when it was applied dissolved in 0.1M-KCl buffered with

0.02M-tris, pH 7.6. It was concluded that the components represented by peaks II, III and IV of the extra protein fraction were not present in the sarcoplasm in significant proportions and were concentrated in the myofibrils. During the chromatography of the sarcoplasmic proteins, coloured fractions presumably corresponding to chromoproteins were observed in the fractions eluted at the points marked X and Y in Fig. 2. Further investigations were not carried out on these coloured fractions.

As fractions II, III and IV were absent from whole sarcoplasm and could readily be separated from the bulk of the sarcoplasmic proteins on diethylaminoethylcellulose, in the later stages of this investigation crude myofibril preparations were used for the preparation of extra protein. This speeded up the preparation considerably and enabled larger amounts to be made. Chromatography of such preparations showed that, apart from some slight differences in the relative amounts of the main fractions, this so-called crude extra protein was virtually identical with that obtained from more exhaustively washed myofibrils. Fractionation of extra protein extracted from a single preparation of myofibrils isolated from hen-breast muscle yielded similar results to those obtained with rabbit myofibrils.

Fig. 2. Chromatography of sarcoplasmic extract of rabbit skeletal muscle made in 25 mM-KCl containing 39 mM-sodium borate buffer, pH 7.1. Protein solution (20 ml.; $E_{1\text{ cm.}} 15.0$ at 280 mμ) in 0.1M-KCl containing 0.02M-tris buffer, pH 7.6, applied to a column 22 cm. x 2 cm. diam. Gradient to 0.35M-KCl, 0.02M-tris applied at point A and steps to M-KCl, 0.02M-tris and 2.0M-KCl, 0.02M-tris at points B and C respectively. In the regions marked X and Y the protein eluted was coloured. ●, Extinction at 280 mμ; □, chloride concentration.

Enzymic studies

In view of the method of preparation the extra protein would be expected to be completely free of myosin, nevertheless assay for calcium-activated ATPase with large amounts of protein (2–3 mg./assay) indicated that traces of myosin were probably present. The amounts were very small as the specific ATPase activity of the concentrated extra protein solutions was less than 1.0 % of that of L-myosin preparations. This ATPase activity was concentrated in fraction II, where it could be detected by using large amounts of protein for enzymic assay (see section on fraction II).

It has been observed earlier (S. V. Perry, unpublished work) that it is extremely difficult to wash rabbit myofibrils free from the last traces of the sarcoplasmic aldolase. This enzyme was found to be present in the extra protein fraction and assay of the fractions obtained with diethylaminoethylcellulose showed that aldolase was concentrated at the front of peak I (material not held in 0.1 M-KCl buffered with 0.02 M-tris, pH 7.6). A relatively low level of activity was also associated with peak III. These studies indicated that the bulk of the aldolase was a constituent of fraction I.

5'-Adenylic deaminase was also not readily removed from repeatedly washed rabbit myofibrils and the extra protein fraction likewise possessed appreciable 5'-adenylic deaminase activity. In 0.1 M-KCl buffered with 0.02 M-tris, pH 7.6, this

enzyme appeared somewhat retarded on diethylaminoethylcellulose. Activity was associated with the tail of peak I and high specific activities were obtained when the bulk of this peak had been eluted. Complete elution was not achieved until the gradient was applied (Fig. 3). The general impression was that 5'-adenylic deaminase probably represented one of the minor components of the extra protein fraction.

Identification of extra protein fractions

Fraction I. The chromatographic behaviour of rabbit sarcoplasm and the limited enzymic studies suggested that fraction I was complex and contained some if not most of the proteins found in sarcoplasm. It was not possible to decide at this stage whether this fraction also contained protein localized exclusively in the myofibrils.

Fraction II: water-insoluble component. The peak obtained on elution of fraction II although usually fairly sharp was not absolutely symmetrical, but in our experience with the chromatography on diethylaminoethylcellulose of muscle proteins, particularly those with asymmetric molecules, such slight irregularities in elution taken without other evidence did not necessarily imply heterogeneity. Evidence for the composite nature of this fraction was provided by the fact that on dialysis of fraction II against water, 70–80 % of the E_{280} -absorbing material came out of solution. This globulin fraction could be redissolved readily by the addition of KCl to ionic strength 0.05–0.10 and usually gave a

Fig. 3. Distribution of 5'-adenylic deaminase activity during chromatography of the extra protein fraction isolated from well-washed myofibrils. Protein solution (25 ml.; $E_{1\text{ cm.}}$ 12.3 at 280 $m\mu$) in 0.02 M-tris, pH 7.6, containing 0.1 M-KCl applied to column 19 cm. $\times$ 2 cm. diam. Gradient to 0.35 M-KCl, 0.02 M-tris at point A and step to 2.0 M-KCl, 0.02 M-tris at point B. Approximately 8 ml. of fractions was collected. Specific deaminase activity was measured as $\mu\text{g. of N}$ liberated as $\text{NH}_3/\text{hr.}$ by 1 ml. of protein solution, $E_{1\text{ cm.}}$ 1.0 at 280 $m\mu$. ●, Extinction at 280 $m\mu$; △, 5'-adenylic deaminase specific activity; □, chloride concentration.

clear solution from which the protein could be reprecipitated by dialysis.

Although fraction II was the only one from which the globulin was precipitated on dialysis against water, the protein could not be satisfactorily prepared by dialysis of the whole original extra protein because non-specific precipitation occurred and almost the whole of the extra protein precipitated under these conditions.

Preliminary studies on the chromatography of myosin, actin and tropomyosin indicated that these proteins were eluted from diethylaminoethyl-cellulose over the range of salt concentration which eluted fractions II and III. In view of this fact and the persistence of low ATPase activity in the globulin fraction the properties of the latter were carefully compared in turn with those of each of the recognized myofibrillar proteins.

It was apparent that the globulin differed markedly from tropomyosin for the following reasons. When it was dissolved in 78 mM-sodium borate buffer, pH 7.1, the viscosity of the solution was relatively low, and unlike tropomyosin showed no significant drop on addition of KCl to 0.5M (Perry, 1953; Perry & Corsi, 1958). Furthermore, tropomyosin was eluted from diethylaminoethyl-cellulose at a higher chloride concentration (Fig. 5), was not precipitated on dialysis against water under conditions which precipitated the globulin and the electrophoretic mobilities of the two proteins were different (Table 1).

Non-identity with F-actin was indicated by the inability of the globulin to form actomyosin. The globulin was not precipitated by 0.75M-phosphate buffer, pH 7.2, and hence did not appear to be identical with inactive actin. Actin inactivated by prolonged dialysis against low-ionic-strength buffers or treatment with KI is precipitated under these conditions (Perry & Corsi, 1958). Treatment of F-actin under conditions identical with those used for the preparation of extra protein, and subsequent chromatography on diethylaminoethyl-cellulose, failed to yield a component corresponding to fraction II. The electrophoretic mobilities of inactive actin and the main component of the globulin preparation were different (Table 1).

Solutions of the globulin differed in appearance from and were clearly less viscous than solutions of L-myosin of similar concentration. The globulin was soluble at 1 0.04, pH 7.0, whereas under these conditions 99 % of myosin solution of similar concentration was precipitated. On electrophoresis in 0.25M-KCl buffered with 0.05M-phosphate, pH 6.9, the main peak obtained with solutions of the globulin moved more slowly and was usually less sharp than the L-myosin boundary.

When relatively large amounts were taken for assay (2-3 mg./incubation tube) ATPase activity

was detectable in the globulin. The specific activity was in the range 50-100 units, whereas under similar conditions L-myosin gave values of 1000-2000 units.

If the extra protein was concentrated by chromatography or by freeze-drying the globulin isolated from it gave a rather consistent electrophoretic pattern (Fig. 4a). There was one main boundary from which separated smaller amounts of one or two faster components on prolonged electrophoresis. On one occasion the main peak split off a smaller slower component, which became

Table 1. *Mobilities of the globulin isolated from fraction II and other myofibrillar proteins*

Conditions: 0.25M-KCl, 0.05M-potassium phosphate buffer, pH 6.9. Mobility is expressed as 10^{-5} cm.²/v/sec.

Protein	Source	No. of experiments	Mobility	
			Asc.	Desc.
Globulin	Rabbit	3	2.1	1.9
Globulin	Hen	1	2.2	2.0
L-Myosin	Rabbit	2	2.6	2.5
Tropomyosin*	Rabbit	3	4.1	3.9
Inactive actin*	Rabbit	3	3.4	3.0

* Perry & Corsi (1958).

Fig. 4. (a) Electrophoretic diagrams of the globulin isolated by dialysis of fraction II from the extra protein of rabbit skeletal myofibrils. Conditions: 0.25M-KCl; 0.05M-potassium phosphate buffer, pH 6.9; 480 min. (b) Ultracentrifuge diagrams of the globulin from fraction II. Conditions: 0.1M-KCl, 0.05M-potassium phosphate buffer, pH 7.15; 5.98×10^4 rev./min. (i) 16 min. (ii) 64 min. (iii) 112 min.

apparent after electrophoresis for 8 hr. Further evidence for non-identity of the main component of the globulin with L-myosin was obtained from the fact that whereas the former protein could be subjected to electrophoresis on filter paper at pH 8.6 in veronal acetate buffer (I 0.1) myosin stayed at the origin. In the ultracentrifuge the number of components which could be recognized was somewhat variable and appeared in some cases to be related to the previous history of the globulin. In some cases of freshly prepared globulin which readily redissolved on addition of salts after precipitation by dialysis one main peak with small amounts of faster and slower components were apparent (Fig. 4b). Attempts to remove the minor components by reprecipitation by dialysis gave a product containing several faster components in addition to the original component. Globulin preparations also became more turbid on standing and handling and the general impression was that aggregation readily occurred and was responsible for the variation in ultracentrifuge records obtained with different globulin preparations.

Globulin corresponding to that isolated from fraction II could not be extracted from granular preparations (see Methods) isolated from sarcoplasm prepared from rabbit skeletal muscle, by the procedure used to prepare the extra protein from myofibrils.

Fraction II: water-soluble component. Owing to the difficulty in isolating sufficient material, little further investigation has been carried out on the water-soluble component of fraction II. One ultracentrifuge run on the pooled freeze-dried protein obtained from three different extra protein preparations revealed one peak with small amounts of a faster component. This faster component had a similar sedimentation velocity to the main component obtained on ultracentrifuging the globulin.

Fraction III: The peak representing fraction III was eluted at 0.3 equiv. of Cl^- ion/l., and when the whole fraction was bulked and concentrated viscometric assay gave values of 50–60% tropomyosin. Fraction III was the only component of the extra protein in which significant amounts of tropomyosin could be detected. When a twice-recrystallized sample of rabbit tropomyosin was chromatographed under similar conditions to those used for the extra protein a sharp peak was eluted at a chloride concentration of 0.29 equiv./l. (Fig. 5). Apart from this peak only traces of material absorbing at $280\text{ m}\mu$ were eluted elsewhere when the KCl concentration in the buffer was changed from 0.1 to 2.0 M. Assay of the tropomyosin content of extra protein isolated from well-washed rabbit myofibrils indicates that the amount of this protein present ranged from 4 to 22%. The relative size of

peak III correlated well with the tropomyosin content of the original extra protein. There was some evidence that the tropomyosin content of extra protein prepared from crude myofibrillar preparations was rather higher than that obtained from well-washed myofibrils.

Fraction IV: nucleoprotein component. After removal of fractions I–III and washing the diethylaminoethylcellulose column with 0.4 M-KCl containing tris buffer until E at $280\text{ m}\mu$ (E_{280}) of the eluate reached a low value, subsequent application of a gradient eluted a symmetrical peak (maximum at ~ 0.7 equiv. of Cl^- ion/l.). Fraction IV differed from the others in that the ratio E_{280}/E_{260} was always less than 1 and usually approached 0.5, a fact which suggested the presence of nucleic acid. On the basis of the following findings it was concluded that fraction IV consisted of ribonucleoprotein.

(1) The u.v.-absorption spectrum measured at pH 7.6 and 3.2 was characteristic for nucleic acid with a maximum at $260\text{ m}\mu$ and a sharp minimum in the region of $240\text{ m}\mu$ (Beaven, Holiday & Johnson, 1955).

(2) The material was non-dialysable.

(3) The diphenylamine test for deoxypentose (Dische, 1930) was negative whereas positive tests for pentose were obtained with the orcinol reaction

Fig. 5. Chromatography of twice-recrystallized rabbit tropomyosin. Tropomyosin solution (6 ml.; $E_{1\text{ cm.}}$ 4.2 at $280\text{ m}\mu$) in 0.15 M-KCl, 0.02 M-tris buffer, pH 7.6, applied to column 23 cm. $\times$ 1 cm. diam. Gradient to 0.5 M-KCl, 0.02 M-tris applied at point A and step to 2.0 M-KCl, 0.02 M-tris buffer at point B. ●, Extinction at $280\text{ m}\mu$; □, chloride concentration.

(Mejbaum, 1939) and the cysteine- H_2SO_4 reaction (Dische, 1949).

(4) A positive biuret test and the shift of the u.v.-absorption minimum towards higher wavelengths suggested the presence of protein.

(5) On chromatography of the products of hydrolysis with 72% perchloric acid (Wyatt, 1951) four spots were obtained, identical with those obtained from a yeast nucleic acid treated under the same conditions (Fig. 6).

It seemed unlikely that the ribonucleoprotein fraction was derived from contamination of the myofibril preparations by nucleic acid-rich elements from muscle cytoplasm such as granules or sarcoplasmic reticulum. As would be expected from the mode of preparation, few granules could be detected

on microscopic examination of myofibril suspensions. To confirm the myofibrillar origin of the ribonucleoprotein, granular material sedimented from sarcoplasmic extracts by centrifuging for 2 hr. at 15 000 *g* was treated by the method used for the preparation of extra protein and the protein solution obtained fractionated on diethylaminoethylcellulose. These preliminary studies indicated that although some nucleic acid could be extracted from the granules by this procedure, the bulk of the nucleic acid isolated from the extra protein was derived from the myofibrils.

DISCUSSION

As obtained from myofibrils prepared from fresh skeletal muscle the extra protein fraction is complex and clearly should not be considered as a homogeneous protein. Tropomyosin is present in this fraction, and although Corsi (1957) considers it to be the main component when extraction is carried out at low pH values, in our hands tropomyosin rarely amounted to more than 20% of the total extra protein nitrogen. In either case the tropomyosin present in the extra protein fraction only accounts for a part of the total amount present in the myofibril. It is possible that the protein isolated by de Villafranca (1956) is in fact tropomyosin in a complex with the other components present in the extra protein fraction. Such an association might modify the solubility properties of tropomyosin but leave unchanged the effect of salt on its viscosity.

Part of the extra protein fraction consists of proteins resembling sarcoplasmic components in their enzymic and chromatographic behaviour, which remain associated with the myofibrils even after exhaustive washing of these structures. Fraction I, in which the sarcoplasmic proteins are concentrated, is very probably identical with the slow-moving component C present in the extract obtained when myofibrils are extracted with buffers of low ionic strength (Perry, 1953; Perry & Corsi, 1958). The persistence of aldolase activity in the myofibril preparations is somewhat comparable with the findings of Roodyn (1956), who studied a similar association of aldolase activity with isolated liver nuclei. In this case, as with the myofibrils, it is difficult to decide whether the association is of metabolic significance or merely an example of non-specific adsorption. The extra protein also possesses appreciable 5'-adenylyc deaminase activity, but most of this enzyme is not associated with the main fractions. It can be concluded that 5'-adenylyc deaminase is only a minor component of the extra protein.

The remaining other two components from the extra protein which have been characterized have

Fig. 6. Chromatography of purine and pyrimidine bases obtained on hydrolysis of fraction IV by heating for 2 hr. at 100° in 72% (w/w) perchloric acid. Propan-2-ol-HCl-water solvent of Wyatt (1951). (a) Fraction IV; (b) yeast ribonucleic acid; (c) fraction IV + yeast ribonucleic acid; (d) thymus deoxyribonucleic acid.

not hitherto been isolated from the myofibril. The globulin possesses certain similarities to the Y-protein fraction (Dubuisson, 1950; Louis, 1954). Y-protein, which is heterogeneous in the ultracentrifuge, is prepared from whole muscle by a method which would be expected to yield a similar fraction to the whole extra protein. Like the Y-protein, most of the whole extra protein fraction precipitates on dialysis against water. This precipitate may be identical with the Y-protein but only part of it consists of the globulin isolated by chromatography in diethylaminoethylcellulose. The evidence suggests that the other components of the extra protein can interact with the globulin to form a water-insoluble complex. It seems probable that the low ATPase activity associated with the globulin is due to myosin contamination, for the physical properties of the former are very different from those of myosin. Nevertheless, the possibility that the globulin is some modified form of myosin of low ATPase activity cannot yet be excluded. Further investigations are now in progress to establish the true significance and nature of the globulin isolated from fraction II.

In the past, histochemical evidence has been produced for the presence of nucleic acid in the myofibril (Gerendas & Matoltzky, 1948; Clavert, Mandel, Mandel & Jacob, 1949). With histochemical studies on sections of whole muscle it is difficult to decide whether the effects obtained are due to overlying granular or reticular matter rather than the myofibril itself. This criticism does not apply to isolated myofibrils, for most of the granules and probably also the sarcoplasmic reticulum fraction are removed during washing. Nevertheless, the ribonucleic acid-rich fraction IV is invariably obtained on chromatography of the extra protein, clearly indicating that ribonucleic acid is a component of well-washed myofibril preparations, and it is probable that it is the source, in part at least, of the nucleic acid moiety of nucleotropomyosin (Hamoir, 1951) and that found associated with myosin (Mihalyi, Bradley & Knoller, 1957; Mihalyi, Laki & Knoller 1957). As separated in diethylaminoethylcellulose the nucleic acid is associated with protein which does not appear to be tropomyosin or myosin. It is possible that this ribonucleic acid is concerned with protein synthesis in the myofibril.

SUMMARY

1. The extra protein fraction of skeletal-muscle myofibrils has been shown to consist of a complex mixture of proteins which can be resolved into four main fractions by chromatography on diethylaminoethylcellulose.

2. About 30 % of the extra protein consisted of a complex fraction containing aldolase activity and

resembling the sarcoplasmic proteins in chromatographic behaviour.

3. Tropomyosin was a somewhat variable component of the extra protein representing 4–22 % of the total protein nitrogen.

4. About 20 % of the extra protein consisted of a protein insoluble in water which may be a component of the Y-protein of Dubuisson (1950).

5. A ribonucleoprotein fraction of myofibrillar origin has been isolated from the extra protein.

We are grateful to Miss F. Johnson for skilled technical assistance throughout this investigation, and would also like to express our thanks to Mr B. Boon and Mr B. Slater for their help with the ultracentrifuge and electrophoresis experiments. This investigation was in part supported by a research-expenses grant (to S.V.P.) from the Medical Research Council.

REFERENCES

- Bailey, K. (1942). *Biochem. J.* **36**, 121.
 Bailey, K. (1948). *Biochem. J.* **43**, 271.
 Bailey, K. & Perry, S. V. (1947). *Biochim. biophys. Acta*, **1**, 506.
 Beaven, G. H., Holiday, E. R. & Johnson, E. A. (1955). In *The Nucleic Acids*, vol. 1, p. 493. Ed. by Chargaff, E. & Davidson, J. N. New York: Academic Press Inc.
 Clavert, J., Mandel, P., Mandel, L. & Jacob, M. (1949). *C.R. Soc. Biol., Paris*, **143**, 539.
 Corsi, A. (1957). *Biochim. biophys. Acta*, **25**, 640.
 de Villafranca, G. W. (1956). *Arch. Biochem. Biophys.* **61**, 378.
 Dische, Z. (1930). *Mikrochemie*, **8**, 4.
 Dische, Z. (1949). *J. biol. Chem.* **181**, 379.
 Dubuisson, M. (1950). *Nature, Lond.*, **166**, 1116.
 Gerendas, M. & Matoltzky, A. G. (1948). *Hung. acta Physiol.* **1**, pp. 124, 128.
 Hamoir, G. (1951). *Biochem. J.* **48**, 146.
 Hanson, J. & Huxley, H. E. (1957). *Biochim. biophys. Acta*, **23**, 250.
 Hasselbach, W. & Schneider, G. (1951). *Biochem. Z.* **321**, 462.
 Louis, P. (1954). *Experientia*, **10**, 258.
 Mejbbaum, W. (1939). *Hoppe-Seyl. Z.* **258**, 117.
 Mihalyi, E., Bradley, D. F. & Knoller, M. I. (1957). *J. Amer. chem. Soc.* **79**, 6387.
 Mihalyi, E., Laki, K. & Knoller, M. I. (1957). *Arch. Biochem. Biophys.* **68**, 130.
 Neuberg, C. & Lustig, H. (1942). *J. Amer. chem. Soc.* **64**, 2722.
 Perry, S. V. (1953). *Biochem. J.* **55**, 114.
 Perry, S. V. (1955). In *Methods in Enzymology*, vol. 2, p. 582. Ed. Colowick, S. P. & Kaplan, N. O. New York: Academic Press Inc.
 Perry, S. V. & Corsi, A. (1958). *Biochem. J.* **68**, 5.
 Perry, S. V. & Grey, T. C. (1956). *Biochem. J.* **64**, 184.
 Peterson, E. A. & Sober, H. A. (1956). *J. Amer. chem. Soc.* **78**, 751.
 Roodyn, D. B. (1956). *Biochem. J.* **64**, 361.
 Sibley, J. A. & Lehninger, A. L. (1949). *J. biol. Chem.* **177**, 859.
 Straub, F. B. (1943). *Stud. Inst. med. Chem. Univ. Szeged*, **3**, 23.
 Szent-Györgyi, A. G., Mazia, D. & Szent-Györgyi, A. (1955). *Biochim. biophys. Acta*, **16**, 339.
 Wyatt, G. R. (1951). *Biochem. J.* **48**, 584.