action of thioglycollate on insulin (Schellman, 1958c).

The results suggest that a rapid uncoiling of the ovalbumin molecule occurs after a certain degree of denaturation has been exceeded, which is in agreement with the serial ethanol-denaturation study previously described.

SUMMARY

1. The degree of denaturation produced in ovalbumin by urea solutions under various conditions has been followed by physical means.

2. During denaturation optical rotation and viscosity show parallel changes and indicate that there is a critical urea concentration. This urea concentration is thought to be associated with the rupture of a linkage important for maintaining the configuration of the native protein.

3. Changes in extinction coefficient at 295 m μ and, after iodination, at 310 m μ also exhibit a similar dependence upon urea concentration.

REFERENCES

- Anfinsen, C. B. & Redfield, R. R. (1956). *Advanc. Protein Chem.* **11**, 1.
 Crammer, J. L. & Neuberger, A. (1943). *Biochem. J.* **37**, 302.
 Frensdorff, H. K., Watson, M. T. & Kauzmann, W. (1953). *J. Amer. chem. Soc.* **75**, 5157.
 Glazer, A. N., McKenzie, H. A. & Wake, R. G. (1957). *Nature, Lond.*, **180**, 1286.
 Li, C. H. (1942). *J. Amer. chem. Soc.* **64**, 1147.
 Li, C. H. (1945). *J. Amer. chem. Soc.* **67**, 1065.
 Maurer, P. H. & Heidelberger, M. (1951). *J. Amer. chem. Soc.* **73**, 2070.
 Schellman, J. A. (1955). *C.R. Lab. Carlsberg, Sér. chim.* **29**, 229.
 Schellman, J. A. (1958a). *C.R. Lab. Carlsberg, Sér. chim.* **30**, 363, 439.
 Schellman, J. A. (1958b). *C.R. Lab. Carlsberg, Sér. chim.* **30**, 429.
 Schellman, J. A. (1958c). *C.R. Lab. Carlsberg, Sér. chim.* **30**, 415.
 Simpson, R. B. & Kauzmann, K. (1953). *J. Amer. chem. Soc.* **75**, 5139.
 Steven, F. S. & Tristram, G. R. (1958). *Biochem. J.* **70**, 179.

Kinetics and Equilibria in Systems Containing Haem, Carbon Monoxide and Pyridine

By M. H. SMITH*

Department of Biochemistry, University of Sheffield

(Received 25 November 1958)

The object of the work described here was to compare the kinetic behaviour of haem compounds of varying degrees of complexity. A single reaction, combination with carbon monoxide, was studied to determine if the rate of combination was variable and the structural factors upon which any variation depended. Three groups of haem compounds were chosen: free haem, the haemochromogens and the haemoproteins. Myoglobin was taken as representative of the last-named group, being formally similar to the simpler haem compounds in having only one haem group per molecule. It also combines reversibly with carbon monoxide, and is very similar in many of its properties to haemoglobin. For the haemochromogen group the compounds with pyridine were chosen, since pyridine has been very widely used and more information was available than for other possible ligands. The effect of changing the side chains on the porphyrin nucleus has also been examined, and mesohaem and deuterohaem and their derivatives

have been prepared in addition to protohaem. The main problem therefore was to determine the rates of combination of carbon monoxide with three closely related haems, three pyridine haemochromogens and three myoglobins (nine reactions in all), the whole representing a gradual increase in the complexity of the haem-conjugated molecule.

The results with myoglobin conveniently stand alone and are dealt with in the next paper (Smith & Gibson, 1959). The present paper is divided into two parts: the first deals with experiments designed to establish the nature of the materials being studied, and the conclusions drawn about their structure; the second gives the results of the kinetic experiments.

It has been generally accepted that the iron atom in a haem molecule is able to combine with only two other groups, one on each side of the haem ring (cf. Lemberg & Legge, 1949). Although the concept of facultative hepta-co-ordination could resolve certain experimental and theoretical difficulties, its use seems to lead, in the end, to even greater perplexities than exist at present. In this work it

* Present address: John Harrison Laboratory of Chemistry, University of Pennsylvania, Philadelphia, 4.

has been assumed that only a sixfold linkage of the iron need be considered, and this is sufficient to explain the observed results and to relate them to previous findings.

MATERIALS AND METHODS

Haemins. Protohaemin was prepared by Drabkin & Austin's (1935) modification of the method of Schälfejeff (1885). A sample of haemin from British Drug Houses Ltd. containing 8.46% of Fe was also used. Mesohaemin was prepared from protohaemin with the reagents of Fischer & Gibian (1941), but no reaction took place when the mixture was allowed to stand for 24 hr. at room temperature. Satisfactory yields were obtained when the reaction mixture was refluxed gently for 15–20 min. Deuterohaemin was prepared from protohaemin by the method of Treibs (1933) as described by Walter (1952). All the samples of haemin were crystallized from acetic acid by the method of Hogness, Zscheile, Sidwell & Barron (1937).

Sodium dithionite. A sample of Manox dithionite was obtained from Holdman and Harden, Miles Platting, Manchester, 1.

Spectrophotometry. The concentration of solutions was determined with a Unicam SP. 600 spectrophotometer. Drabkin (1942) found that the maximum extinction coefficients for several derivatives of meso- and coprohaem had the same numerical values as those for the corresponding derivatives of protohaem. It therefore seemed reasonable to use the value of Hogness *et al.* (1937) for the extinction coefficient of protohaemin, $\epsilon_{482\text{ m}\mu}^{\text{max}} = 6.55$ for all three haematin. Readings were taken at 482 m μ for meso- and deutero-haematin, however. All readings were taken in a 1 cm. cell unless otherwise stated.

The millimolar extinction coefficients are defined as $\epsilon = E/cl$, where C is millimolarity of the haem solution, l is optical depth and E (observed extinction) is $\log I_0/I$, where I_0 and I are the intensities of the incident and transmitted light respectively.

Preparation of solutions. As neither haematin nor dithionite is appreciably soluble in glycol alone, the following standard procedure was used throughout. Stock solutions of haematin in 0.02N-NaOH were made up as required in sufficient strength to give the required extinction after dilution. To 0.5 ml. of a stock solution of haematin (0.15 mm), 1 ml. of 2.5% (w/v) $\text{Na}_2\text{S}_2\text{O}_4$ in 0.02N-NaOH was added and the mixture diluted to 10 ml. with 0.02N-NaOH. This gave a haem solution suitable for spectrophotometric or kinetic experiments in the Sorlet region with a 1 cm. optical cell. If pyridine was to be introduced into the system this was done by adding the necessary small volume of a stock solution in water before making up to 10 ml. with 0.02N-NaOH. Where ethylene glycol was to be used, the haem solution was made up to 10 ml. with ethylene glycol instead of NaOH. There was always at least 70% (v/v) of glycol and usually 80–90%. Some preliminary experiments suggested that about 60% (v/v) was required to produce the full effect on the spectrum.

Carbon monoxide-haem dilution test. Carbon monoxide-haem was examined over the range 0.075–65 μm by recording the change in extinction at 406.5 μm , the Sorlet maximum of the substance. Various depths of cell from 0.1

to 4.0 cm. were employed to obtain measurable extinctions at all concentrations. These were all subsequently converted into the corresponding values for a 1 cm. cell.

Carbon monoxide-haem dissociation curve. The methods used to establish the curve consisted of (i) direct equilibration of haem solutions with varying partial pressures of CO, and (ii) mixing aqueous solutions of known concentrations of haem and CO in a closed system and observing the change in extinction. In both the extinction at 406.5 μm was recorded, the fractional saturation then being given by the increase of extinction over that of the free haem, divided by the increase on forming fully saturated CO-haem from free haem.

Flash-photolysis kinetic measurements were carried out with the apparatus and procedure of Gibson (1959).

Photochemical absorption spectrum. The construction of the flash-photolysis apparatus used in the kinetic work made it possible to calculate the change in extinction of the solution being examined, at the moment of flashing. If the initial spectrum was known, then the absolute spectrum immediately after flashing could be calculated, as well as the difference spectrum. The former corresponds to the 'photochemical absorption spectrum' of Warburg & Negelein (1929) and Lemberg & Legge (1949), but is now obtained by direct measurement.

RESULTS

Structure of haem and carbon monoxide-haem

Spectrophotometric observations. Much of the available information about the simpler haem compounds has come from spectrophotometric work. So far as haem and its derivatives are concerned, it seems (Shack & Clark, 1947) that a distinction can be drawn between monomeric and dimeric forms in aqueous solution on the basis of their absorption spectra. These criteria appear to be applicable also to haem derivatives in non-aqueous solvents. The general finding is that the monomer has a much sharper spectrum than that of the dimer (which is usually markedly rounded-off and often shifted to a longer wavelength than that of the monomer), and that the monomer has a much greater extinction at the absorption maximum.

Comparison of the absorption spectra of free haem and of CO-haem in aqueous solution (Figs. 1 and 2) shows that on this basis the former is dimeric and the latter monomeric. Fig. 1 shows the dramatic difference between the absorption spectra of free haem in aqueous solution and in glycol. On the premise that aqueous haem is dimeric, the considerable increase in extinction and shift in position of the Sorlet band, together with the form of the new curve, lead to the conclusion that the solution in glycol contains monomeric haem.

Use of ethylene glycol. It is generally recognized that many substances which occur as polymers in aqueous solution tend to split up into lower polymers and eventually monomers in organic solvents

(Rabinowitch & Epstein, 1941; Lewis, Magel & Lipkin, 1942; Lewis, Goldschmid, Magel & Bigeleisen, 1943). The use of ethylene glycol in this respect is cited in the paper by Lewis *et al.* (1943), and it has recently been used for the same purpose in this Department (Weber & Teale, 1958). The findings could not be fully confirmed with other solvents as several common organic solvents, including ethanol and acetone, were found to be unsuitable because of the formation of a white precipitate, presumably as a result of reaction with the dithionite (a sulphurous smell was usually noticeable). The general effect of these solvents, however, appeared to be similar to that of glycol.

With CO-haem (Fig. 2) there is only a slight increase in absorption at the maximum and no change in the form of the curve. It is not clear whether the increase is due directly to the change in the solvent, or whether it is due to the presence of small amounts of free (dimeric) haem in aqueous solutions of CO-haem. Observations made in this Department by Dr G. Weber (personal communication), on the effect of different solvents on the spectra of various aromatic nitrogenous compounds, suggest that the former is very probable.

Dilution test. When the observed extinctions at various dilutions were converted into the corresponding values for 10 cm. cell, the plot of extinction against concentration gave a straight line passing through the origin. Although confirming that the material obeyed Beer's law down to very low concentrations, this result could not by itself be taken as evidence that the CO-haem was monomeric. Reproducible results could not be obtained in similar experiments with free haem.

Dissociation curve. Some time was spent on establishing the dissociation curve of CO-haem, in view of the information which this might give about the system. It was found difficult to obtain consistent results, partly owing to the high affinity of haem for CO, which tended to render observations difficult by the methods available, and partly because of unexplained differences between the earlier and subsequent results. In each case several different solutions of haem were used. The differences are at present tentatively attributed to variations in the state of aggregation of the haem. The earlier results indicated that in 0.02N-NaOH at room temperature (about 20°) the haem was 50% saturated at an equilibrium value of 0.75 mm. Hg CO or 1.0 μ M. Thirty-three points obtained over the whole range of the dissociation curve in six experiments were analysed by the method of least squares, giving $K \cdot 0.92 \times 10^6$ ($K' = [\text{CO-H}]/[\text{CO}][\text{H}]$ concentrations in moles, where H is the haem molecule and $n = 1.17$). The results were compatible with a value of $n = 1$. (In the Hill equation, the value n gives a measure of the degree of

Fig. 1. Absorption spectra in the Soret and visible regions of alkaline protohaem, in 0.02N-NaOH (—) and in 80% (v/v) ethylene glycol (---). [Haem] 7 μ M; light path 1 cm. pH of aqueous solution 11.5.

Fig. 2. Absorption spectra in the Soret and visible regions of alkaline CO-protohaem, in 0.02N-NaOH (—) and in 80% (v/v) ethylene glycol (---). [Haem] 7 μ M; light path 1 cm.

interaction of the haems: $n = 1$ means either that the substance is monomeric or that it is polymeric but with negligible interaction.) Later experiments gave 50% saturation in the range 3.5–4.5 μM -CO (mean about 3 mm.Hg CO with $K = 0.33 \times 10^6$). Both sets of results were obtained under apparently identical conditions.

Since this section of the work was completed, Wang, Nakahara & Fleischer (1958) have presented what appears to be the first published set of data for the dissociation of CO-haem. Their figures indicate a half-saturation value of about 3.1 μM (2.4 mm.Hg CO), at approximately 24.6°. The curve plotted from their figures is a rectangular hyperbola, passing through the origin, from which n may be calculated to equal 1.0. These figures, which agree very well with the second set of those given above, are apparently based upon a single set of observations, and there is no mention that variations can occur.

Photochemical absorption spectrum. The photochemical absorption spectrum of CO-haem solutions showed peaks at 406 $m\mu$ and at 420 $m\mu$. The latter corresponds in position to the absorption maximum of haem in glycol. The changes are not yet fully explained since it appeared that even in glycol an appreciable fraction of the CO-haem was not dissociated; yet increasing the intensity of the flash did not give the expected increase in photolysis. It is presumed that the reactive portion yields monomeric haem even in aqueous solution, and that this then recombines with CO, possibly with dimeric haem as an intermediate.

Structure of pyridine and carbon monoxide-pyridine haemochromogens

Changes in extinction at a single wavelength. By varying the concentration of pyridine in equilibrium with a fixed amount of haem it may be shown that, at the concentration of haem used, there must be more than one compound formed. The changes were most easily followed in the Soret region, and in order to use 1 cm. cells the haem solutions were made up to between 5 and 10 μM . (No variation in the results was noted which could be ascribed to differences in concentration over this range.) Although the conclusions drawn from experiments with this dilute haem have been tentatively assumed to apply also to somewhat stronger solutions (up to at least 0.1 mm), only a few observations have so far been made with the stronger solutions. The general findings presented in this and other sections of this paper therefore refer directly to the more dilute solutions of haem, and some care must be taken in extending them to stronger solutions.

To show the changes in extinction with increase in pyridine concentration, the composite figure

shown in Fig. 3 was built up from a number of single observations with varying amounts of pyridine. It was found that as the concentration of pyridine was increased, the extinction observed at a single wavelength increased in a regular manner. At first there was a small increase, approaching completion at about mm-pyridine. This change was best observed at 410 $m\mu$ (see next section); subsequent changes were best observed at 420 $m\mu$. As the pyridine concentration was raised more, there was a further and larger increase in absorption, which reached a maximum at about 60 mm-pyridine. One further increase in extinction took place on raising the pyridine concentration, and this was maximal at about 10M-pyridine. No further increase occurred, even in 80% (v/v) pyridine. The last two stages are quite clear-cut and easily reproducible, and although replicate experiments did not necessarily agree exactly, their general form was very similar. The initial curve was not always easy to demonstrate, partly owing to the small change in extinction, and possibly also owing to differences in the state of aggregation of the haem.

Dr Philip George and Mr Alan Adler of the John Harrison Laboratory, University of Pennsylvania, have confirmed (personal communication) that the spectrophotometric titration of pyridine against haem gives a stepped curve of the type described. Their experiments were done within 3 hr. of making up the haemin solution and at pH 7.0. As in the work reported above, they found observations of the initial stages to be the most difficult. In a

Fig. 3. Change in extinction at a single wavelength on adding pyridine to alkaline protohaem (data from a single experiment). Extinction was observed at 420 $m\mu$. [Haem] 12 μM throughout; light path 1 cm. Pyridine concentration in final solution varied from 0 to 10M. Maximum millimolar extinction at 418 $m\mu$ was more generally found to be about 130. Inset scale applies to first step only, i.e. to 3.25 units on abscissa.

typical experiment maximum formation of the last two compounds occurred at about 70 mm- and 2.5M-pyridine. The millimolar extinction of haem at 418 m μ at these points was 67.0 and 188. The latter value is considerably higher than was found in the present work (see Figs. 3 and 4). Adler & George report that, in general, higher values for the extinction of pyridine haemochromogen were found than are given in the literature, and that this may possibly be correlated with the age of the haem solution employed. On the other hand, however, experiments involving the oxidized form from the same stock solution gave extinction coefficients similar to, or slightly lower than, those previously reported. Experimental conditions used by these workers were: haematin, 8 μ M in 0.2M-phosphate buffer, pH 7.0; reduction by solid dithionite; whole spectra in the Soret region were obtained with a Beckman Recording Spectrophotometer with 1 cm.

The curve shown in Fig. 3 demonstrates that the stepwise addition of pyridine forms no less than three compounds with haem. The changes were most easily observed in the Soret region owing, as will be shown in the next section, to the wide separation of the absorption maxima of the three forms in this region. They can also be observed, but with more difficulty, in the visible region, where all three forms show the typical two-banded haemochromogen spectrum. Investigation of the visible absorption spectra has not yet been detailed enough to show with certainty whether or not a shift in wavelength accompanies the increase in extinction. If there is a shift it is only of the order of 2 m μ . Shack & Clark (1947), using 0.1 mm-haem, reported that 'on further addition of pyridine [to haem] the peak near 560 m μ progressively shifted to lower wavelengths'. The extent of the shift was not stated. In the present work the absorption maximum of (monomeric) pyridine haemochromogen was found to be at 555 m μ , as measured in the Unicam SP. 600 spectrophotometer [but cf. Drabkin's (1942) value of 558 m μ]. The Soret extinction of free haem in glycol, at 420 m μ , so closely resembles the peak of pyridine haemochromogen in glycol in position and extinction coefficient that observations on the change from one to the other are difficult to make. In the visible region with glycol only one stage has been observed, corresponding to the last stage in NaOH; the change is maximal at 0.1M-pyridine. A few experiments with the meso- and deuterohaem were less satisfactory, since the absorption bands of the different compounds are closer together than with protohaem, and there is overlapping of absorption bands in the reactions with pyridine. There is, however, no reason to suppose that the properties of meso- and deuterohaem in

these systems are different from those of protohaem (cf. Walter, 1952).

Absorption spectra of pyridine haemochromogen with varied pyridine concentration. Examination of the absorption spectra in the Soret region during the determination of the pyridine dissociation curves shows three distinct types of spectrum, which merge to give composite spectra at intermediate concentrations of pyridine. Fig. 4 shows the observed spectra in the Soret region of free haem, and of the solution at stages corresponding to the maximum formation of each intermediate compound.

Further spectrophotometric observations in the Soret region are set out in Figs. 5-8. In the visible region the usual maximal formation of each of the three pyridine haemochromogen compounds and CO-haemochromogen spectra have been observed, but the changes have not yet been followed so closely as in the Soret region.

In the presence of CO the spectral changes are simpler (all experiments on the effect of CO were carried out by equilibrating the system with the gas at a pressure of 1 atmosphere). The CO-haem peak at 406 m μ (Fig. 5) shifted to about 410 m μ and became more rounded (Fig. 6). With 50-100 mm-pyridine the maximum extinction was

Fig. 4. Absorption spectra of free alkaline protohaem and haem-pyridine compounds in the Soret region, in 0.02N-NaOH. pH of solution 11.5. The three spectra in the presence of pyridine represent the observed spectrum at maximum formation of each intermediate haem-pyridine compound (cf. Fig. 3). Pyridine concentration in the observed solution: curve 1, 0 (free haem); curve 2, 1-2 mm; curve 3, 50-100 mm; curve 4, 10-13M. Maximum millimolar extinction at 418 m μ was more generally found to be about 130. [Haem] 12 μ M; light path 1 cm.

Fig. 5. Absorption spectra in the Soret region of the γ -band of alkaline CO-protohaem, in 0.02N-NaOH (—) and in 80% (v/v) ethylene glycol (---). (No pyridine was present.) [Haem] $7 \mu\text{M}$; light path 1 cm.

Fig. 7. Absorption spectra in the Soret region of alkaline CO-protohaem with 50–100 mM-pyridine, in 0.02N-NaOH (—) and in 80% (v/v) ethylene glycol (---). The spectrum in aqueous solution demonstrates incomplete conversion into the monomeric CO-haemochromogen (peak at $418 \text{ m}\mu$), the shoulder at about $430 \text{ m}\mu$ being due to the presence of dimeric dipyridine haemochromogen (see Fig. 4). [Haem] $7 \mu\text{M}$; light path 1 cm.

Fig. 6. Absorption spectra in the Soret region of alkaline CO-protohaem with 1–2 mM-pyridine, in 0.02N-NaOH (—) and in 80% (v/v) ethylene glycol (---). [Haem] $7 \mu\text{M}$; light path 1 cm.

Fig. 8. Absorption spectra in the Soret region of alkaline CO-protohaem with 10–13 mM-pyridine, in 0.02N-NaOH (—) and in 80% (v/v) ethylene glycol (---). [Haem] $7 \mu\text{M}$; light path 1 cm.

about 65% of the original value. There was little further change until the pyridine concentration was raised above 0.4M, when a sharper peak emerged at 418 m μ (Fig. 8). It was found to be essential to add pyridine to CO-haem thoroughly saturated with CO, not to bubble CO through pyridine haemochromogen solution. In the latter case the peak at 430 m μ , formed in the intermediate range of pyridine concentration, remained even after passing CO (Fig. 7). It was necessary to ensure thorough saturation with CO because this same peak could appear even when the pyridine was added last, presumably through combination of pyridine with free haem.

Ethylene glycol solutions were again used to supplement the observations with aqueous solutions. In the absence of CO there was only a single peak at 418 m μ , higher and sharper than corresponding peaks in NaOH, at all concentrations of pyridine. There appeared to be a slight increase in the height of the peak with increasing pyridine concentration. At the highest pyridine concentration the peak in glycol resembled the corresponding one in NaOH very closely, but gave a slightly higher extinction. In the presence of CO exactly the same behaviour was observed, but the extinctions were in all cases greater than those in absence of CO and did not vary very much with change in pyridine concentration.

Photochemical absorption spectrum. Photochemical absorption spectra were taken at 1.0 mM- and 2.3M-pyridine. In each case the spectrum showed a maximum at 415–420 m μ . In the first case this represented an increase in extinction over this range; in the second it represented a decrease.

Kinetic results

Carbon monoxide-haem. Carbon monoxide-haem is commonly regarded (cf. Lemberg & Legge, 1949) as being photosensitive, yet the first experiments appeared to show that no reaction was taking place. Examination of the literature showed that the published evidence for photosensitivity derived from the observations of Warburg & Negelein (1929) and Krebs (1929). These workers described manometric experiments in which haem catalysed the oxidation of cysteine to cystine. The presence of cysteine at once rendered their interpretations suspect since the work was done before it was shown (Keilin, 1950) that a number of amino acids can form haemochromogens, and the authors did not appear to have ruled out this possibility. However, it has not proved possible to obtain such a haemochromogen under the conditions employed in the present work, nor is there any sign of the formation of a mixed CO-cysteine compound with the haem. Presumably a similar situation existed in the experiments cited, where the haem was dis-

solved in phosphate and borate buffers at pH 6.8 and 10.3.

It was subsequently possible to demonstrate the photolysis of the CO-haem, the reason that this had not been observed earlier being due to the rapidity of the recombination reaction. This was found to be some 40 or more times faster than the CO-myoglobin reactions already studied (Smith & Gibson, 1959), and could only be observed in the existing apparatus by reducing the partial pressure of CO to not more than 10–20 mm. Hg.

In view of the technical difficulties in measuring these very fast reaction rates, which are among the fastest reactions known involving biochemical substances, the results in Table 1 must be regarded as approximate only.

Carbon monoxide-haemochromogens. So far, no evidence has been found that any of the pyridine haemochromogens is photosensitive in the absence of CO. In experiments in which increasing amounts of pyridine were added to CO-haem it was found that the very rapid CO-haem reaction gradually became biphasic, eventually giving a much slower, monophasic reaction. At still higher pyridine concentration this too became biphasic and finally there was a very slow, apparently monophasic reaction at M or greater concentrations of pyridine.

Table 1. *Rate of combination of proto-, meso- and deuterio-haem with carbon monoxide*

Concentration of the haem was 7 μ M, of CO $\sim$ 20 μ M, temperature 20°. Observations were made over the range 410–430 m μ , usually at 430 m μ . Solutions with glycol contained 80% (v/v); in all cases the concentration of Na₂S₂O₄ was 0.25% (w/v). Flash-photolysis energy 400J; light path 1 cm.

Haem	Solvent	10 ⁻⁷ × Velocity constant (M ⁻¹ sec. ⁻¹)
Proto-	0.02N-NaOH	1.6
	Glycol	1.4
Meso-	0.02N-NaOH	8.7
	Glycol	5.1
Deutero-	0.02N-NaOH	8.4
	Glycol	—

Table 2. *Rate of combination of monopyridine haemochromogens with carbon monoxide*

Haem (7 μ M) was dissolved in 0.02N-NaOH and reduced with dithionite as described in the section on Methods. [CO] 140 μ M. Temperature 20°: Flash-photolysis energy 100J; reaction was followed by the change in absorption at 420 m μ .

Haem	10 ³ × Pyridine concentration (M)	10 ⁻⁶ × Velocity constant (M ⁻¹ sec. ⁻¹)
Proto-	4.0	0.42
Meso-	6.5	1.04
Deutero-	10.5	1.30

These findings may be explained by supposing that the first biphasic reaction is due to the presence in the same solution of CO-haem and CO-haemochromogen (both monomeric). The reaction with the haemochromogen is much slower than the reaction with haem, and only when sufficient pyridine has been added to convert all of the haem into haemochromogen does the reaction become monophasic. This result can only be observed over a small range of pyridine concentration, further increase in the amount of pyridine causing the tracing to become biphasic again. The appearance of a second component in the reaction at this point is not believed to be due to a new reacting species but to result from competition between pyridine and CO for a place on the haem molecule. The relative affinities appear to be such that with millimolar CO (1 atm. partial pressure) a fairly high proportion of the haem is in combination with CO even at the highest pyridine concentrations. This equilibrium state is upset by flash photolysis and with high pyridine concentrations dipyrindine haemochromogen, rather than monopyridine-water-haemochromogen, is formed immediately after flashing. Carbon monoxide is unable to combine directly with dipyrindine haemochromogen and the rate of the observed reaction is no longer determined by the combination velocity constant for the reaction: monopyridine haemochromogen + CO $\rightarrow$ CO-pyridine haemochromogen, but by the dissociation velocity constant of the reaction dipyrindine haemochromogen $\rightarrow$ monopyridine haemochromogen + pyridine. This replacement reaction, which has a half-time of several seconds, forms the slowest component of the reaction at high pyridine concentrations.

In practice it was found to be difficult to establish the precise values for the reaction: monopyridine haem + CO $\rightarrow$ CO-pyridine haemochromogen, owing to the tendency for the results to be influenced either by the CO + haem reaction (fast) or by the replacement reaction (slow). The reaction could only be observed over a relatively small range of pyridine concentrations and the values in Table 2 are taken from the results of a single experiment carried out after the explanation given above had been worked out, but they are in general agreement with between three and six experiments for each haem which had been previously made under less fully suitable conditions.

DISCUSSION

The work of Shack & Clark (1947) and of Walter (1952) has established that in aqueous solution free haem is present in the form of dimeric units aggregated into micelles. These authors did not, however, investigate the influence of carbon monoxide

upon such systems. Carbon monoxide-haem was represented by Hill (1926) as CO-H-X, 'where X is either another molecule of the complex or a molecule of water', the symbol H representing the haem nucleus. If the former possibility were correct, then the formula would in fact be CO-H-H-CO, i.e. two carbon monoxide molecules combined with dimeric haem. Although this possibility does not seem to have been definitely excluded by later workers (cf. Keilin, 1949*a, b*) it has been generally assumed that carbon monoxide-haem is monomeric. Combination of pyridine with haem was shown by Hill (1926, 1929) to lead to the formation of a monomeric haemochromogen consisting of one haem nucleus combined with two pyridine molecules. The later work of Shack & Clark (1947) and of Walter (1952) has suggested that the formation of the monomeric haemochromogen from dimeric haem may proceed stepwise, although no definite evidence has been presented that this is so. With the combination of haematin with pyridine, these workers have shown that there is in fact a pH-dependent transition from dimer to monomer, each form being combined with two molecules of the base. The effect of carbon monoxide on haemochromogen systems has not yet been studied from the point of view of structure, and the available evidence has always been interpreted as due to the reaction of carbon monoxide with a monomeric haemochromogen. The final product is known (Hill, 1926) to contain iron, carbon monoxide and pyridine in equal proportions and can only be monomeric.

The general findings of the present spectrophotometric work have served to confirm the tentative division of haem compounds into dimers and monomers, postulated by Shack & Clark (1947) on the basis of the observed absorption spectra, as well as the general shift of the absorption bands of compounds of meso- and deuterio-haem towards the blue, as compared with those of protohaem, already demonstrated by Drabkin (1942). The finding that solutions of haem reduced with dithionite may be readily and safely prepared in ethylene glycol has made it possible to provide direct spectrophotometric evidence for such a classification, and has proved a useful tool for the investigation of complex haem systems.

Haem and carbon monoxide-haem. The experiments with haem and carbon monoxide-haem support the belief of earlier workers that these substances in aqueous solution are normally dimeric and monomeric respectively. A change from dimeric haem to a monomeric form can be demonstrated by changing the solvent (cf. Maehly & Åkeson, 1958). With carbon monoxide-haem there is very little spectral difference in the two solvents, and the findings in the dilution test and

dissociation curve also support the view that the substance is normally monomeric. The photochemical absorption spectrum serves as a link between carbon monoxide-haem and free haem, by suggesting that monomeric haem is formed from carbon monoxide-haem even in aqueous solution. This change may be represented by the equation:

where H represents the haem nucleus. (Presumably the monomeric haem combines immediately with a water molecule, not remaining with a free valency.) In view of the general agreement with Hill's (1926) finding that the ratio of iron to carbon monoxide was 1:1 in carbon monoxide-haem, indicating the presence of monomeric haem, it seems likely that there is a straightforward conversion into monomeric carbon monoxide-haem at all haem concentrations.

Pyridine haemochromogen and carbon monoxide-pyridine-haemochromogen. The dissociation curves and absorption spectra of pyridine haemochromogen systems have shown that the addition of pyridine to haem can take place in a stepwise manner. Since the starting material (haem) is dimeric, only two dimeric and two monomeric haem-pyridine compounds are theoretically possible. In ethylene glycol only the two monomeric forms can occur.

Scheme 1

Experiment showed that in aqueous solution there were three compounds, and in glycol only one. Scheme 1 is therefore proposed to account for the various observations. It is well known (Hill, 1929) that pyridine haemochromogen contains two pyridine molecules per iron atom. Shack & Clark (1947) and Walter (1952) agree that this form is monomeric. This makes it quite clear that the final pyridine-haem compound (I) has the formula indicated. In order to account for its formation in the number of steps given by the dissociation curves, it is necessary to postulate that the first pyridine molecule adds to the haem to give (III), which by addition of a second pyridine molecule passes to (II). This sequence is preferred to the

alternative formulation of the monomeric monopyridine compound as the intermediate, on the following grounds: (i) the intermediate spectrum appeared to be that of a dimeric, rather than a monomeric, molecule, and has not been observed in glycol solutions, where only monomeric haem compounds have so far been identified; (ii) it did not appear in solutions of carbon monoxide-haem to which pyridine was added. On the other hand, its spectrum persisted as a secondary peak (see Fig. 7) if carbon monoxide was passed into solutions of (II). These qualitative observations suggest that (II) is not easily formed from the monomeric carbon monoxide-haem, nor does it readily give rise to this compound. Moreover, the dissociation curve of the pyridine compound in glycol appeared to be indicative of a change from free haem directly to haemochromogen I. The evidence therefore favoured the supposition that a monomeric monopyridine haemochromogen was not a normal intermediate either in aqueous solutions or in ethylene glycol. It is presumably formed on the photolysis of carbon monoxide-haemochromogen, and is tentatively assumed to have an absorption spectrum very similar to that of the dipyridine compound.

The explanation given above can be correlated with observations on systems containing carbon monoxide, as is shown in scheme 2, which is applicable both to aqueous systems and those containing ethylene glycol. Here, the only possibility in either solvent is for the carbon monoxide-haem to form the carbon monoxide-haemochromogen, and this is in accordance with spectrophotometric observation. It is well known that pyridine can compete with carbon monoxide for a place on the haem molecule. Under certain conditions therefore the system may pass over wholly or partially to haemochromogen I. This possibility is believed to explain the kinetic results at high pyridine concentrations, as discussed above.

Previously published work has not been found which can be directly correlated with these findings, other than the suggestions of Shack & Clark (1947) and of Walter (1952) that association might be stepwise. Most workers have either used strong haem solutions, or else have investigated systems containing oxidized haem only. The closest analogy appears to be with the haem-cyanide system, where stepwise association undoubtedly occurs (Hogness *et al.* 1937; Keilin, 1949*a*). However, there does not seem to be

any possibility that dimeric compounds exist in the cyanide system, and it is actually more directly comparable with the carbon monoxide-haem system than with the haem-pyridine system, with the distinction that a compound of haem with two carbon monoxide molecules has not yet been identified. The conclusions drawn about the structure of various components of the systems examined are in good agreement with the findings of other workers. Although it remains to be seen if these conclusions are fully applicable to the stronger solutions used in potentiometric work, for instance, there is no doubt that they have to be taken into account when working with haem solutions up to 20–30 mm. In the common case of the estimation of haem as the pyridine haemochromogen the conventional use of 10–20% (v/v) pyridine (1.2–2.4M) has doubtless obviated anomalous results, as well as delaying the recognition of the changes in the haem-pyridine system.

As no kinetic data have hitherto been available for the combination of carbon monoxide with the simple haem compounds investigated, it is not possible to compare the results with those of previous workers. Nevertheless, they seem to be on the whole so clear-cut that definite conclusions may be drawn from them.

Comparison of the results given in Table 1 for the free haems with those in Table 2 shows a reduction in the combination rate on passing from free haem to the pyridine chromogen. This reduction is so large as to suggest that something more fundamental than a mere steric effect is operating. In any case, the results of the present investigations have demonstrated that the respective substances are both monomeric and, although it is not possible altogether to rule out the possibility that there is loose aggregation of the molecules through the side chains of the porphyrin nucleus, there is no reason to suppose that the photochemical reaction or the recombination reaction are affected by this.

Apart from steric and concentration effects, the velocity of a chemical reaction depends upon the ease with which bonds may be broken and formed. Where the co-ordination number of the iron appears to be less than six, as in the products of the photochemical reaction, it has been assumed that water fills the remaining place. In the reactions to be considered, the bond between the haematin iron and a water molecule must presumably be broken before the carbon monoxide compound can be reformed. Since this applies in all the cases cited, and since there is at present no way of evaluating the extent of such a hindrance to recombination, the reactions will for convenience of description be considered as taking place directly between iron and carbon monoxide.

The rate of reaction of certain aromatic compounds can be predicted theoretically (cf. Dewar, 1949), but no information appears to be available for any haem compounds. The results presented above now make it possible to appraise these particular systems comparatively, taking the rate of formation of carbon monoxide-haem as a standard. The rates of reaction will depend upon the availability of the valency electrons of the iron atom, which will participate in the bond with carbon monoxide, and this will be affected by the nature of the haem itself as well as by the nature of the conjugated group which is not dissociated by light. (Carbon monoxide is itself intensely reactive owing to the fact that the electronic configuration of the carbon atom is depleted, whereby the molecule acquires an electron-acceptor or cationoid character.)

Since the only difference between protohaem and protopyridine haemochromogen lies in the replacement of a water molecule by a pyridine group, this must be the cause of the marked difference in the velocity constants of the two forms. The pyridine molecule is strongly cationic by comparison with water, i.e. it tends to attract electrons towards itself. It appears therefore that the direct linkage of pyridine to the central iron atom reduces the accessibility of the valency electrons to carbon monoxide (as shown by the reduction in the velocity constant), by the attraction which it exerts upon those electrons. Assuming that there actually is combination with a water molecule after photolysis, it could be considered that a 'tighter' bond was being formed, i.e. that the volume of the electron cloud was being reduced through the effect of the pyridine. The same phenomenon is observed with natural myoglobin (Smith & Gibson, 1959). The two rates are in fact virtually identical, suggesting that the effect is already maximal even upon combination with such a relatively small molecule as pyridine.

A similar argument can be applied to the differences observed within each group, where the structure of the individual haem is being varied. The haems differ only in the constitution of the side chains at positions 2 and 4 of the porphyrin ring, namely, vinyl, ethyl and hydrogen respectively for proto-, meso- and deuterio-haem. Such groups may be arranged in a series in which the cationoid character progressively increases. The three groups of immediate interest can be arranged in the series hydrogen < ethyl < vinyl. Hence protohaem, which contains two vinyl groups, might be expected to show the slowest reaction rates owing to the attraction exerted on the valency electrons through the structure of the molecule. Similarly, deuterohaem should show the fastest rates since

hydrogen atoms will exert little if any effect; meso-haem should be slower than deuterohaem. This relationship does apply to the pyridine haemochromogen reactions, which have rates in the ratio of 1:2.5:3.1. (The myoglobin reactions give anomalous results in the mesomyoglobin reaction.) The free haem reactions at first sight appear to be anomalous in that meso- and deuterohaem apparently have the same rate, although this is still much faster than that of protohaem. However, it must be noted that these reactions are extremely fast, being of the same order of magnitude as some of the peroxidase reactions (Chance, 1949), and it is possible that the two fastest rates may require some revision when further improvements in technique become possible. It may be that there is a similar progression with the free haems but that this is obscured by the present difficulties in measuring the very fastest rates.

One obvious drawback in applying such an interpretation to individual haems is the electro-metric effect, known to occur in the vinyl group, whereby electrons may be donated to other parts of the molecule by resonance. This would oppose the inductive effect invoked above, and is usually found to be more important. On the other hand, the differences between the observed reaction rates are not great enough to rule out a predominating inductive effect in this instance. It is to be hoped that further work with substituted haems will be undertaken to resolve this point. For the present it appears that variation in the inductive attraction of the side groups represents one characteristic property of the molecule which can account for the observed data; and this suggestion is put forward as a basis for further investigation.

SUMMARY

1. Spectrophotometric observations of haem and carbon monoxide-haem in aqueous solution and in ethylene glycol suggest that carbon monoxide-haem is monomeric in both solvents, whereas haem itself is dimeric in aqueous solution but is depolymerized in ethylene glycol.

2. The dissociation curve of carbon monoxide-haem in aqueous solution was found to be a rectangular hyperbola. Different haem solutions gave 50% saturation at 0.75 and 3.5–4.5 mm.Hg carbon monoxide at 20°. The material also obeyed Beer's law over the range of carbon monoxide concentration 0.075–65 μM .

3. The photochemical absorption spectrum of carbon monoxide-haem indicated that only a part of the material was being dissociated on flash photolysis. A new peak appeared at 420 $\text{m}\mu$, corresponding in position to the peak of monomeric haem in glycol. It is suggested that this new

peak is caused by the absorption spectrum of the haem nucleus subsequent to the photodecomposition.

4. The dissociation of pyridine haemochromogen shows three stages. Spectrophotometric evidence supports the belief that three compounds of haem and pyridine are formed consecutively, the last of which is the well-known 'pyridine haemochromogen'. It is suggested in explanation of these observations that two of the compounds are dimeric, only 'pyridine haemochromogen' itself being monomeric.

5. The rate of combination of carbon monoxide with three closely related haems was measured by the flash-photolysis technique. The rates of combination with proto-, meso- and deuterohaem respectively were found to be approximately 16, 85 and 85 $\mu\text{M}^{-1} \text{sec}^{-1}$ in 0.02N-sodium hydroxide at room temperature.

6. The rate of combination of carbon monoxide with the pyridine haemochromogens of the same three haems was similarly measured. The rates were found to be respectively 0.42, 1.04 and 1.30 $\mu\text{M}^{-1} \text{sec}^{-1}$ under the same conditions.

7. A theoretical explanation of the variation in these results, and also of those reported in the next paper, is attempted.

This work was carried out during the tenure of a Medical Research Council Scholarship for Training in Research Methods and this support is gratefully acknowledged.

REFERENCES

- Chance, B. (1949). *Arch. Biochem.* **22**, 224.
- Dewar, M. J. S. (1949). *The Electronic Theory of Organic Chemistry*. Oxford: Clarendon Press.
- Drabkin, D. L. (1942). *J. biol. Chem.* **146**, 605.
- Drabkin, D. L. & Austin, J. H. (1935). *J. biol. Chem.* **112**, 89.
- Fischer, H. & Gibian, H. (1941). *Liebigs Ann.* **548**, 183.
- Gibson, Q. H. (1959). *Biochem. J.* **71**, 293.
- Hill, R. (1926). *Proc. Roy. Soc. B*, **100**, 419.
- Hill, R. (1929). *Proc. Roy. Soc. B*, **105**, 112.
- Hogness, T. R., Zscheile, F. P., jun., Sidwell, A. E., jun. & Barron, E. S. G. (1937). *J. biol. Chem.* **118**, 1.
- Keilin, J. (1949a). *Biochem. J.* **45**, 440.
- Keilin, J. (1949b). *Biochem. J.* **45**, 448.
- Keilin, J. (1950). *Nature, Lond.*, **165**, 151.
- Krebs, H. A. (1929). *Biochem. Z.* **204**, 343.
- Lemberg, R. & Legge, J. W. (1949). *Hematin Compounds and Bile Pigments*. New York: Interscience Publishers Inc.
- Lewis, G. N., Goldschmid, O., Magel, T. T. & Bigeleisen, J. (1943). *J. Amer. chem. Soc.* **65**, 1150.
- Lewis, G. N., Magel, T. T. & Lipkin, D. (1942). *J. Amer. chem. Soc.* **64**, 1774.
- Maehly, A. C. & Åkeson, A. (1958). *Acta chem. scand.*, **12**, 1259.

Rabinowitch, E. & Epstein, L. F. (1941). *J. Amer. chem. Soc.* **63**, 143.
 Schallfejeff, M. (1885). *J. Soc. phys.-chim. russe*, p. 30. Quoted in *Ber. dtsch. chem. Ges.* **18** (Referate) 232.
 Shack, J. & Clark, W. M. (1947). *J. biol. Chem.* **171**, 143.
 Smith, M. H. & Gibson, Q. H. (1959). *Biochem. J.* **73**, 101.

Treibs, A. (1933). *Liebigs Ann.* **506**, 196.
 Walter, R. I. (1952). *J. biol. Chem.* **196**, 151.
 Wang, J. H., Nakahara, A. & Fleischer, E. B. (1958). *J. Amer. chem. Soc.* **80**, 1109.
 Warburg, O. & Negelein, E. (1929). *Biochem. Z.* **204**, 495.
 Weber, G. & Teale, F. W. J. (1958). *Trans. Faraday Soc.* **54**, 640.

The Preparation and some Properties of Myoglobin Containing Meso- and Deutero-Haem

By M. H. SMITH* AND Q. H. GIBSON
Department of Biochemistry, University of Sheffield

(Received 25 November 1958)

Although experiments in which the protohaem of haemoglobin was replaced by other haems or porphyrins were first carried out by Hill & Holden (1926), the number of papers dealing with similar experiments is comparatively small, and even fewer have been concerned with the quantitative study of reactions of these 'synthetic' haemo-proteins. Myoglobin, unlike haemoglobin, is free from effects due to haem-haem interaction and therefore seems particularly suited for work on the relation between the constitution of the haem group and the properties of the haemoprotein, yet little work has been reported. Drabkin (1945) has described the reconstitution of myoglobin with protohaem and has found that the reconstituted material has the same spectral absorption and crystal habit as the native protein. The preparation and properties of reconstituted myoglobin have recently been more fully described by Theorell & Åkeson (1955).

Since this work was completed a preliminary report has appeared from Rossi-Fanelli & Antonini (1957), who have prepared native human myoglobin apo-protein and have coupled it with protohaem and deuterohaem. They report the spectra of various derivatives, and found that there was a threefold increase in oxygen affinity on replacing protohaem with deuterohaem.

The object of the present experiments was to prepare myoglobins containing meso- and deuterohaem, to examine their spectra and to investigate the kinetics of their combination with carbon monoxide. A preliminary account has been given by Gibson & Smith (1957).

* Present address: John Harrison Laboratory of Chemistry, University of Pennsylvania, Philadelphia, 4.

EXPERIMENTAL

Haemins. The haemins used were prepared as described in the preceding paper (Smith, 1959).

Sodium dithionite. A sample of Manox dithionite was obtained from Holdman and Harden, Miles Platting, Manchester, 1.

Buffers. Phosphate buffers, pH 6.5–7.5 (Sørensen, 1909); for pH 5.0–6.3, 0.2N-citric acid–0.2N-NaOH (Britton & Welford, 1937); for pH 7.8–9.1, 0.2M-H₃BO₃ in 0.2N-KCl with 0.2N-NaOH (Clark, 1928).

Whale myoglobin. Whale meat from a local fishmonger was prepared both by the methods of Theorell & de Duve (1947) and of Bowen (1948). No crystals were obtained from the samples of whale meat available to us by either of these methods, although considerable yields of myoglobin were obtained in amorphous form.

Horse-heart myoglobin. Horse hearts were obtained fresh from the local abattoir and prepared by the method of Bowen (1948). Four hearts were used at various times, of which three yielded the typical wheatsheaf clusters of crystals at the final stages of the preparation. The fourth gave amorphous material only.

Preparation of native globin. Haematin was removed from the myoglobin molecule by a method devised by Dr F. W. J. Teale (personal communication). A solution of metmyoglobin was dialysed for 24 hr. against running distilled water in a rocking dialyser in 0.5 in. cellophan tubing. To 1 vol. of an approximately mM-solution of the salt-free myoglobin at 0°, 0.5 vol. of 2-butanone containing 10% (v/v) of 0.1N-HCl was added. The mixture was shaken gently and allowed to stand for 15 min. If a layer of 2-butanone had separated, it was removed. A second portion of acid butan-2-one equal in volume to the first was then added and the mixture again shaken and allowed to stand for 15 min. The process of removal of 2-butanone and its replacement with a fresh portion was continued until no further haematin passed into the upper layer. The lower solution of globin was dialysed against water or 0.02M-phosphate buffer, pH 7.1. The solution was stable for several days when stored at 0° and pH 7.1.