had 5.9 times the phosphate and 2.7 times the acetate incorporation of fraction 3. Both these fractions developed strong purple colours when treated with Feulgen reagent, indicating the presence of plasmalogens (Feulgen & Bersin, 1939). Collins & Wheeldon (1957) isolated two classes of mixtures of methyl esters of dinitrophenylkephalins from both rat liver and egg yolk. The first class had a phosphorus:dinitrophenol ratio of 1.0 and the second of 2.0. The second class, when derived from rat-liver homogenates incubated with [^{32}P]phosphate, had incorporated more radioactivity than the first class or the corresponding choline-containing lipids. It is possible that the same complexities of kephalin structure occur in human blood cells. In these studies no attempts were made to distinguish between the cellular components of blood, e.g. red cells, white cells and platelets, and it is possible that the different contributions of these may be partly responsible for the metabolic heterogeneity observed.

SUMMARY

1. The simultaneous incorporation *in vitro* of [^{32}P]orthophosphate and [$\text{Me-}^{14}\text{C}$]acetate into the phospholipids of human blood cells has been measured.

2. The doubly labelled phospholipids obtained after incubation with the radioactive isotopes for 6.5 hr. were fractionated by chromatography on silica gel, into ethanolamine-containing kephalins, serine-containing kephalins, phosphatidylcholine, sphingomyelin and other phospholipids.

3. All fractions were labelled with both ^{32}P and ^{14}C . Of the isolated phospholipid ^{32}P the kephalins contained the most (67%), followed by the lecithins (12%) and the sphingomyelins (6.5%). The incorporation of ^{14}C followed the same order, the kephalins containing 54%, the lecithins 11% and the sphingomyelins 10% of the isolated lipid ^{14}C .

4. The fatty acids of the fractionated phospholipids were analysed by vapour-phase chromatography and different distributions of fatty acids for each fraction were observed.

5. Two ethanolamine-containing phospholipid fractions were isolated. These differed in the rates of phosphate and acetate incorporation, and in fatty acid composition.

The author is grateful to Dr J. E. Lovelock for helpful advice and the use of a gas-chromatography apparatus. Thanks are due to Mr P. Simmonds and Mr C. Widnell for technical assistance.

REFERENCES

- Altman, K. I. (1953). *Arch. Biochem. Biophys.* **42**, 478.
 Altman, K. I., Whatman, R. N. & Salomon, K. (1951). *Arch. Biochem. Biophys.* **33**, 168.
 Chargaff, E., Levine, C. & Green, C. J. (1948). *J. biol. Chem.* **175**, 67.
 Collins, F. D. & Wheeldon, L. W. (1957). *Biochem. J.* **66**, 441.
 Erf, L. A., Tuttle, L. W. & Lawrence, J. A. (1941). *Ann. intern. Med.* **15**, 487.
 Feulgen, R. & Bersin, T. (1939). *Hoppe-Seyl. Z.* **260**, 217.
 Finean, J. B. (1957). *Metabolism of the Nervous System*. Ed. by Richter, D. London: Pergamon Press Ltd.
 Fiske, C. H. & Subbarow, Y. (1925). *J. biol. Chem.* **66**, 375.
 Hack, M. H. (1947). *J. biol. Chem.* **169**, 137.
 Hanahan, D. J., Dittmer, J. C. & Warashina, E. (1957). *J. biol. Chem.* **228**, 685.
 James, A. T. (1958). *Amer. J. clin. Nutr.* **6**, 494.
 James, A. T., Lovelock, J. E. & Webb, J. (1957). *Biochem. J.* **66**, 60 p.
 Lawrence, J. A., Erf, L. A. & Tuttle, L. W. (1941). *J. appl. Phys.* **12**, 333.
 Lea, C. H., Rhodes, D. N. & Stoll, R. D. (1955). *Biochem. J.* **60**, 353.
 Lipsky, S. R. & Landowne, R. A. (1958). *Biochim. biophys. Acta*, **27**, 666.
 Lovelock, J. E. (1955a). *Biochem. J.* **60**, 692.
 Lovelock, J. E. (1955b). *Brit. J. Haematol.* **1**, 117.
 Lovelock, J. E. (1958). *J. Chromat.* **1**, 35.
 Tuttle, L. W., Scott, K. G. & Lawrence, J. H. (1939). *Proc. Soc. exp. Biol., N.Y.*, **41**, 20.

Studies on Sulphatases

25. THE DETERMINATION OF $\text{BaS}^{16}\text{O}_3^{18}\text{O}$ BY INFRARED SPECTROSCOPY*

By B. SPENCER

Department of Biochemistry, University of Wales, Cardiff

(Received 18 March 1959)

In vitro, the various enzymes of the sulphatase group hydrolyse their specific sulphate ester substrates with the formation of inorganic sulphate and the corresponding desulphated hydroxy compound.

Whether this is also their action *in vivo* or whether they can act as transferases is not yet known with certainty. Two transferring activities, involving the transfer of either the sulphate group or the desulphated residue, can be envisaged. A knowledge of the position of cleavage of the sulphate esters by

* Part 24: Lloyd (1959).

the enzymes would be helpful in deciding which of these transfer reactions could occur; splitting between the oxygen and sulphur atoms would indicate the possibility of sulphate transfer whereas C—O bond rupture would indicate transfer of the asulphate.

The position of cleavage can be studied by allowing the sulphatase to act on its substrate in the presence of water containing excess of ^{18}O . Splitting of the C—O bond will result in incorporation of ^{18}O into the hydroxy compound formed (1), whereas in O—S bond-splitting the ^{18}O will be found in the liberated sulphate (2). The sulphate can be precipitated as barium sulphate and its ^{18}O content assessed.

In a previous study with arylsulphatases the ^{18}O content of barium sulphate was measured by combustion with carbon and mass-spectrographic determination of the ^{18}O content of the carbon dioxide formed (Spencer, 1958). Extension of this study to other sulphatases was hampered by the lack of ready access to a mass spectrometer and alternative methods of determining the ^{18}O content of barium sulphate were therefore considered. Infrared-absorption spectrometry seemed promising since the substitution of the heavier ^{18}O atom for ^{16}O in barium sulphate should lead to a decrease in the frequencies of vibration in the molecule with consequent shifting of the infrared absorption peaks to longer wavelengths. The magnitude of the shift can be calculated from the Teller-Redlich equation. The free sulphate ion is a tetrahedral XY_4 molecule belonging to the point group T_d and for the fully isotopically substituted $\text{XY}_4^{(i)}$

$$\frac{\omega_1}{\omega_1} = \frac{\omega_2^{(i)}}{\omega_2} = \sqrt{\frac{m_Y^{(i)}}{m_Y}} \quad (3)$$

for the totally symmetrical (ν_1) and doubly degenerate (ν_2) vibrations where ω_1 and ω_2 are the frequencies of the ν_1 and ν_2 vibrations respectively, m_Y is the mass of the Y atom and all the quantities marked with the superscript (i) refer to the isotopic material (Herzberg, 1945, p. 233). For S^{16}O_4 and S^{18}O_4 $\omega_1^{(i)}/\omega_1 = 0.942$ and since ν_1 for the free S^{16}O_4 ion is 981 cm^{-1} (Table 1) the expected shift of this vibration in S^{18}O_4 will be 57 cm^{-1} . As a rough approximation the shift of ν_1 for $\text{S}^{16}\text{O}_3^{18}\text{O}$ will be one-quarter of this value, i.e. 14 cm^{-1} , and a shift of this magnitude in the region around

1000 cm^{-1} should be readily discernible with most infrared spectrophotometers. To check whether these predicted shifts might be of practical use for the detection of $\text{BaS}^{16}\text{O}_3^{18}\text{O}$, a sample of barium sulphate containing 87% of $\text{BaS}^{16}\text{O}_3^{18}\text{O}$ was prepared and its infrared-absorption spectrum examined.

EXPERIMENTAL

Materials. Potassium *p*-nitrophenyl sulphate was prepared by the method of Burkhardt & Lapworth (1926). Water containing 10 and 90 atoms % excess of ^{18}O was obtained from The Radiochemical Centre, Amersham, Bucks and the Weizmann Institute of Science, Rehovoth, Israel.

Preparation of $\text{BaS}^{16}\text{O}_3^{18}\text{O}$. Potassium *p*-nitrophenyl sulphate (52 mg.) was dissolved in $54.8 \mu\text{l}$. of water containing 90 atoms % excess of ^{18}O and acidified with $2.8 \mu\text{l}$. of 10N-HCl . The arylsulphate was completely hydrolysed by heating the solution, which was contained in a 3 mm. int. diam. tube in a boiling-water bath for 15 min. The non-immersed part of the tube was cooled in a stream of air to prevent excess of evaporation. After cooling, the hydrolysed material was diluted with 1 ml. of water, extracted with 1 ml. of ether ($\times 4$) to remove *p*-nitrophenol and then warmed to expel dissolved ether. The liberated sulphate was precipitated by the addition of 1 ml. of 5% (w/v) BaCl_2 and the barium sulphate formed was washed on the centrifuge with 2 ml. of N-HCl ($\times 2$), 2 ml. of water ($\times 5$), 2 ml. of ethanol ($\times 2$) and finally dried *in vacuo* over P_2O_5 for 2 days. It has previously been established that 100% splitting of the O—S bond occurs in the acid hydrolysis of *p*-nitrophenyl sulphate (Spencer, 1958) and, taking into account the water added in the 10N-HCl , the barium sulphate isolated in the present experiment was calculated to contain 21.75 atoms % excess of ^{18}O or 87% of $\text{BaS}^{16}\text{O}_3^{18}\text{O}$. The $\text{BaS}^{16}\text{O}_3^{17}\text{O}$ content was 1.2% and the $\text{BaS}^{16}\text{O}_4$ content, by difference, was 11.8%. The majority of the work was carried out on this specimen, but some confirmatory spectra were obtained with a sample containing 90% of $\text{BaS}^{16}\text{O}_3^{18}\text{O}$. This was prepared by passing sulphur trioxide into water containing 90 atoms % excess of ^{18}O and precipitating and washing the barium sulphate as before.

Samples (20 mg.) containing different amounts of $\text{BaS}^{16}\text{O}_3^{18}\text{O}$ were prepared by mixing the 87% isotopic material with normal barium sulphate. To obtain homogeneous mixtures on this small scale the components were weighed into a small centrifuge tube, suspended in 2 ml. of ethanol and the tube was vibrated against a flattened nail rotating at about 1500 rev./min. After centrifuging, the ethanol was removed and the mixture dried *in vacuo* over P_2O_5 .

Infrared-absorption spectrophotometry. Spectra were studied on a double-beam Perkin-Elmer Infracord with a rock-salt prism. Accurate measurements of the frequencies

of the absorption peaks were obtained on a double-beam Perkin-Elmer Model 21 with a rock-salt prism for frequencies above 700 cm^{-1} and a single-beam Perkin-Elmer model 12C with a KBr prism for frequencies below 700 cm^{-1} . Liquid paraffin (Nujol) mulls were used throughout.

Arylsulphatase of Aspergillus nidulans. *Aspergillus nidulans* (C.M.I. 16643) was grown on Czapek-Dox synthetic medium for 6 days with aeration. The mycelium was harvested, washed with water and starved by aeration in water for 5 hr. The mycelium was then ground in a mortar with acetone at 0° and allowed to stand in acetone at this temperature for 3 hr. The mycelium was separated by filtration, washed well with cold acetone and dried *in vacuo*. The supernatant of an aqueous suspension (2.5%, w/v) of the acetone-dried material was dialysed against several changes of water at 2° and then freeze-dried. The lyophilized material hydrolysed 3.91 μmoles of *p*-nitrophenyl sulphate/hr./mg. at 38° in 0.1M-acetate buffer, pH 6.2, with 0.015M-substrate.

Arylsulphatase assay. A final volume of 20 μl . of 0.5M-acetate buffer, pH 6.2, containing potassium *p*-nitrophenyl sulphate (0.015M), the enzyme and the inhibitor where appropriate, was incubated at 38° for 1 hr. The reaction was stopped by the addition of 100 μl . of 0.5M-NaOH and, after centrifuging, the absorption at 400 $\text{m}\mu$ was measured in microcells with the Hilger Uvispek spectrophotometer. The amount of *p*-nitrophenol liberated was calculated by the use of ϵ_{400} 18 200.

RESULTS

Method developed for the determination of $\text{BaS}^{16}\text{O}_3^{18}\text{O}$

The frequencies of the infrared absorption peaks of normal barium sulphate and barium sulphate con-

taining 87% of $\text{BaS}^{16}\text{O}_3^{18}\text{O}$ are shown in Table 1. The significance of the spectra are discussed later but, of practical interest, it was noted that whereas the 981 and 1076 cm^{-1} frequencies of $\text{BaS}^{16}\text{O}_4$ were shifted to 961 and 1059 cm^{-1} in $\text{BaS}^{16}\text{O}_3^{18}\text{O}$, a weak peak at 981 cm^{-1} and a shoulder at 1072 cm^{-1} were also shown in the isotopic sample (Table 1 and Fig. 1). Since the isotopic sample contained 11.8% of $\text{BaS}^{16}\text{O}_4$ in addition to the $\text{BaS}^{16}\text{O}_3^{18}\text{O}$ it seemed reasonable to attribute the subsidiary peak and the shoulder to the presence of the normal barium sulphate.

This detection of 11.8% of $\text{BaS}^{16}\text{O}_4$ in the presence of 87% of $\text{BaS}^{16}\text{O}_3^{18}\text{O}$ suggested that observations at 981, 961, 1076 and 1059 cm^{-1} might be used to estimate the amounts of normal and isotopic barium sulphate in a given sample. The peaks at 1076 and 1059 cm^{-1} , however, proved of little use for this purpose since even the best resolution, obtained with the sample containing 11.8% of $\text{BaS}^{16}\text{O}_4$ and 87% of $\text{BaS}^{16}\text{O}_3^{18}\text{O}$ with the Perkin-Elmer 12C spectrophotometer, succeeded only in showing the presence of the minor component as a slight shoulder on the major peak. In a sample containing 60% of $\text{BaS}^{16}\text{O}_4$ and 40% of $\text{BaS}^{16}\text{O}_3^{18}\text{O}$ the specific peaks at 1076 and 1059 cm^{-1} respectively were indistinguishable and a new flat peak between the two appeared (Fig. 1). On the other hand, the ν_1 peak in both $\text{BaS}^{16}\text{O}_4$ and $\text{BaS}^{16}\text{O}_3^{18}\text{O}$ was sharp and in various mixtures of normal and isotopic barium sulphate peaks at 981 and 961 cm^{-1} were clearly resolved (Fig. 1).

Table 1. Infrared spectra of normal barium sulphate and barium sulphate containing 87% of $\text{BaS}^{16}\text{O}_3^{18}\text{O}$

Assignments for barium sulphate are from Ramdas (1954). The fundamental frequencies of the free sulphate ion are known from Raman spectra in solution (Kohlrausch, 1943).

Barium sulphate			Free $\text{S}^{16}\text{O}_4^{2-}$ ion		
Frequencies (cm^{-1})		Assignment	Assignment	Degree of degeneracy	Frequencies (cm^{-1})
$\text{BaS}^{16}\text{O}_4$	$\text{BaS}^{16}\text{O}_3^{18}\text{O}$ sample				
981	961	$\left\{ \begin{matrix} \omega_1^1 (B_{1u}) \\ \omega_1^2 (B_{3u}) \end{matrix} \right\}$	$\nu_1 (a_1)$	1	981
?	?				
	(981 weak)				
?	?	$\left\{ \begin{matrix} \omega_2^2 (B_{2u} \text{ or } B_{1u}) \\ \omega_2^3 (B_{1u} \text{ or } B_{2u}) \end{matrix} \right\}$	$\nu_2 (e)$	2	451
?	?				
?	?				
1076	1059	$\left\{ \begin{matrix} \omega_3^1 (B_{1u}) \\ \omega_3^2 (B_{3u}) \end{matrix} \right\}$			
1112	1095				
1187	1189	$\left\{ \begin{matrix} \omega_3^3 (B_{1u}) \\ \omega_3^4 (B_{3u}) \end{matrix} \right\}$	$\nu_3 (f_2)$	3	1104
?	?				
?	?	$\left\{ \begin{matrix} \omega_3^5 (B_{3u}) \\ \omega_3^6 (B_{2u}) \end{matrix} \right\}$			
	(1072 shoulder)				
611	604	$\left\{ \begin{matrix} \omega_4^1 (B_{1u}) \\ \omega_4^2 (B_{1u}) \end{matrix} \right\}$			
637	634				
?	?	$\left\{ \begin{matrix} \omega_4^3 (B_{1u}) \\ \omega_4^4 (B_{3u}) \end{matrix} \right\}$	$\nu_4 (f_2)$	3	613
?	?				
?	?				

A method for the determination of $\text{BaS}^{16}\text{O}_3^{18}\text{O}$ in barium sulphate was developed by utilizing the resolution of the ν_1 frequency. The principle is illustrated in Fig. 2, which shows part of the spectrum of a paraffin oil (Nujol) mull of barium sulphate containing 8.7 % of $\text{BaS}^{16}\text{O}_3^{18}\text{O}$. The values of *A* and *B* (Fig. 2) from the $\log I/I$ scale were taken to represent the absorption due to normal and isotopic barium sulphates respectively. The percentage of $\text{BaS}^{16}\text{O}_3^{18}\text{O}$ in the sample is given by $100 B/A+B$ and in the case illustrated the measurements gave 9 %. Although based on the Lambert-Beer law this method can only be approximate since it also depends to some extent upon the freehand completion of the $\text{BaS}^{16}\text{O}_3^{18}\text{O}$ and $\text{BaS}^{16}\text{O}_4$ spectra as shown by the dotted lines in Fig. 2. To make the method more sensitive concentrated mulls requiring about 20 mg. of sample have been used and a 0.05 mm. spacer has been placed between the rock-salt plates. In this way the extinction of the ν_1 peaks is effectively increased but at the same time the base of the peaks is broadened and the freehand completion of the spectra becomes more approximate. In practice a compromise is made by limiting the extinction of the strongest ν_1 peak to 0.5. A linear magnification of *A* and *B* (Fig. 2) can be obtained by adjusting the spectrophotometer so that the trace at the approximate position *C* has an extinction of about 0.05.

Fig. 1. Infrared-absorption spectra of Nujol mulls of barium sulphate samples. *A*, Normal barium sulphate; *B*, barium sulphate containing 87 % of $\text{BaS}^{16}\text{O}_3^{18}\text{O}$ and 11.8 % of $\text{BaS}^{16}\text{O}_4$; *C*, 40 % of $\text{BaS}^{16}\text{O}_3^{18}\text{O}$, 59.5 % of $\text{BaS}^{16}\text{O}_4$; *D*, 8.7 % of $\text{BaS}^{16}\text{O}_3^{18}\text{O}$, 91.2 % of $\text{BaS}^{16}\text{O}_4$.

A straight line was obtained when the known $\text{BaS}^{16}\text{O}_3^{18}\text{O}$ content of prepared samples was plotted against the percentage of the isotopic material calculated from the spectra (Fig. 3). The presence of as little as 2 % of $\text{BaS}^{16}\text{O}_3^{18}\text{O}$ (i.e. 0.5 atom % excess of ^{18}O) could be detected and an accuracy varying between 2 atoms % of $\text{BaS}^{16}\text{O}_3^{18}\text{O}$

Fig. 2. A portion of the spectrum of a Nujol mull of a barium sulphate sample containing 8.7 % of $\text{BaS}^{16}\text{O}_3^{18}\text{O}$. The mull was prepared with 20 mg. of sample and a 0.05 mm. spacer was placed between the rock-salt plates of the cell. For details see text.

Fig. 3. Spectra of prepared samples of barium sulphate containing various amounts of $\text{BaS}^{16}\text{O}_3^{18}\text{O}$ were examined; 20 mg. samples and a 0.05 mm. spacer were used. The percentages of $\text{BaS}^{16}\text{O}_3^{18}\text{O}$ calculated from the spectra are plotted against the known concentration.

for low values of isotopic barium sulphate and 4 atoms % of $\text{BaS}^{18}\text{O}_3^{18}\text{O}$ for high values can be considered a reasonable claim.

Biological application of the method

Mechanism of action of the arylsulphatase of Aspergillus nidulans. To check the practical efficiency of the infrared method of $\text{BaS}^{18}\text{O}_3^{18}\text{O}$ estimation the position of cleavage of potassium *p*-nitrophenyl sulphate by the arylsulphatase of *A. nidulans* was studied. The reaction mixture contained 12 mg. of lyophilized *A. nidulans* preparation, 30 mg. of potassium *p*-nitrophenyl sulphate and 0.136 g. of sodium acetate trihydrate in 2 ml. of ^{18}O -enriched water and the whole was adjusted to pH 6.2 with a measured amount of acetic acid by using a glass electrode. The final concentration of ^{18}O in the water of the reaction mixture was calculated to be 8.7 atoms % excess. After incubating at 38° for 3 hr. the liberated inorganic sulphate was precipitated as barium sulphate and washed free of protein, *p*-nitrophenol and unhydrolysed substrate as detailed previously (Spencer, 1958). The barium sulphate (yield 18.2 mg.) was analysed by the infrared spectroscopic method and was found to contain 7.8 % of $\text{BaS}^{18}\text{O}_3^{18}\text{O}$. The theoretical value for O-S cleavage of *p*-nitrophenyl sulphate was 8.7 % of $\text{BaS}^{18}\text{O}_3^{18}\text{O}$.

Effect of ions on the arylsulphatase of Aspergillus nidulans. Arylsulphatases have been divided into two types on the basis of their relative substrate specificities and their behaviour towards certain ions (Dodgson & Spencer, 1957). Type I enzymes are scarcely affected by phosphate, sulphate or fluoride but are markedly inhibited by cyanide whereas the type II enzymes behave in the opposite manner. The effect of these and other ions on the arylsulphatase activity of *A. nidulans* (Table 2) shows that this enzyme can be classified as a type I arylsulphatase.

Table 2. *Inhibition of the arylsulphatase activity of Aspergillus nidulans*

Activity was measured against 0.015M-potassium *p*-nitrophenyl sulphate in 0.5M-acetate buffer, pH 6.2, over 1 hr. at 38°. The inhibitors (0.02M final concentration) were adjusted to the assay pH by means of a glass electrode before being added to the incubation mixture.

Compound added	Inhibition (%)
KCN	65.5
KH_2PO_4	1.3
K_2SO_4	1.2
NaF	19.9
Na_2SO_3	48.7

DISCUSSION

The conventional crystallographic unit cell of barium sulphate contains four molecules of barium sulphate and this is also the primitive or Bravais cell (Bhagavantam & Venkatarayudu, 1951) from which selection rules may be deduced. The free sulphate ion, which belongs to the tetrahedral point group T_d , has nine fundamentals which give rise to four distinct frequencies, a_1 (single), e (doubly degenerate) and f_2 (two, both triply degenerate). In the barium sulphate crystal the symmetry of a single sulphate molecule is changed to the monoclinic local symmetry C_2 , and the result is a perturbation of the internal vibrations of the sulphate ion thereby removing the degeneracy and giving rise to nine separate frequencies. The number of internal modes of the four sulphate ions in the primitive barium sulphate cell will therefore be 36. Since there is a centre of symmetry the mutual exclusion rule applies and it can be shown that of the expected frequencies 15 will be infrared active, 18 Raman active and 3 inactive. Thus in the infrared spectrum of barium sulphate two bands near 981 cm^{-1} , three bands near 451 cm^{-1} , five bands near 1104 cm^{-1} and five bands near 613 cm^{-1} can be expected. Of these predicted bands only six were actually observed (Table 1). None of the weakly infrared-active components of ν_2 was seen either in the normal or isotopic barium sulphate. Duval & Lecomte (1954) with barium sulphate and Hass & Sutherland (1956) with gypsum also failed to observe the ν_2 bands, but they are evident in the sulphates of copper, magnesium and manganese (Duval & Lecomte, 1954). Ramdas (1954), using barytes crystals in the 001 and 110 cleavage sections, observed nine of the 15 expected infrared fundamentals including two components of ν_2 , and a further band is known from reflexion spectra. In the present study slight differences in the frequencies of the overtone and combination bands of the normal and isotopic barium sulphates were observed in the region 2000–3000 cm^{-1} but an accurate analysis was not attempted.

Absorption spectra of mulls do not allow a more specific assignment of the observed frequencies than that given in Table 1, and an exact comparison of the observed shifts with the theoretical shifts calculated from the Teller-Redlich equation is not possible. However, it is generally considered (Herzberg, 1945, p. 238) that for both small and large mass differences no crossing over of frequencies of the same species can occur because of the repulsion in consequence of Fermi resonance, and it is therefore permissible to correlate the absorption frequencies of the normal and isotopic barium sulphates as shown. The band frequencies observed for the isotopic material may not

necessarily be the actual fundamental frequencies of $\text{BaS}^{16}\text{O}_3^{18}\text{O}$ but may be intermediate peaks between those of $\text{BaS}^{16}\text{O}_3^{18}\text{O}$ (87%) and the $\text{BaS}^{16}\text{O}_4$ (11.8%) present in the sample. This is more likely for the strongly absorbing broad ν_3 peaks but, on the other hand, the ν_1 peaks of $\text{BaS}^{16}\text{O}_4$ and $\text{BaS}^{16}\text{O}_3^{18}\text{O}$ are clearly resolved.

The infrared spectrophotometric method for the determination of $\text{BaS}^{16}\text{O}_3^{18}\text{O}$ is insensitive in comparison with the mass-spectrographic method (Hoerring, 1952; Spencer, 1958), which can readily distinguish differences of 0.01 atom % excess of ^{18}O . Clearly, the infrared method can be of little use with samples containing low concentrations of ^{18}O or when small differences in isotope content between two samples are required to be measured, but for the study of the position of cleavage of sulphate esters by sulphatases, where an 'all or none' response can be expected, the method offers the advantages of speed and simplicity. For experiments of this type it is advisable to carry out the enzyme reactions in water containing at least 5 atoms % excess of ^{18}O , and the availability of such material constitutes the major limiting factor in the use of the method.

Application of the method to the study of the hydrolysis of *p*-nitrophenyl sulphate by the arylsulphatase of *Aspergillus nidulans* showed that the enzyme acts by cleaving the O-S bond of the arylsulphate. On theoretical grounds (cf. Spencer, 1958) it may therefore be deduced that if the enzyme can act as a transferase it will transfer the sulphate rather than the phenol group. Inhibition studies indicate that the enzyme, which is similar to the arylsulphatase of Taka-diaxase (*Aspergillus oryzae*), is a type I arylsulphatase and its method of action is also the same as other type I enzymes, namely rat-liver arylsulphatase C and the arylsulphatase of *Alcaligenes metalcaligenes*, which are already known to split the O-S bond. No actual transferring activity has yet been demonstrated for any type I arylsulphatase.

SUMMARY

1. The infrared-absorption spectra of normal barium sulphate and barium sulphate containing 87% of $\text{BaS}^{16}\text{O}_3^{18}\text{O}$ have been examined.

2. A method of estimating $\text{BaS}^{16}\text{O}_3^{18}\text{O}$ in barium sulphate has been developed, based on the measurement of relative extinctions at 981 and 961 cm^{-1} .

3. The method has been used to analyse the barium sulphate prepared from the inorganic sulphate liberated by *Aspergillus nidulans* arylsulphatase acting on potassium *p*-nitrophenyl sulphate in the presence of water containing 8.7 atoms % excess of ^{18}O . The $\text{BaS}^{16}\text{O}_3^{18}\text{O}$ content of the barium sulphate was found to be 7.8%, showing that the type I arylsulphatase of *Aspergillus nidulans* splits the O-S bond of arylsulphates.

I am most grateful to Dr E. Whittle and Mr C. G. Cannon for help with the infrared spectra. The investigation was supported in part by a Royal Society Grant and by a research grant (A-1982) from the U.S. Public Health Service.

REFERENCES

- Bhagavantam, S. & Venkatarayudu, T. (1951). *Theory of Groups and Application to Physical Problems*, 2nd ed. p. 127. Waltair, India: Andhra University.
- Burkhardt, G. N. & Lapworth, A. (1926). *J. chem. Soc.* p. 684.
- Dodgson, K. S. & Spencer, B. (1957). In *Methods of Biochemical Analysis*. Ed. by Glick, D. New York: Interscience Publishers.
- Duval, C. & Lecomte, J. (1954). *C.R. Acad. Sci., Paris*, **239**, 249.
- Hass, M. & Sutherland, G. B. B. M. (1956). *Proc. Roy. Soc. A*, **236**, 427.
- Herzberg, G. (1945). *Molecular Spectra and Molecular Structure*. New York: Van Nostrand Co.
- Hoerring, T. (1952). Ph.D. Thesis: Washington University, U.S.A.
- Kohlrausch, K. W. F. (1943). *Ramanspektren. Hand- und Jahrbuch der chemischen Physik*, vol. 9, pt. 6, p. 399. Leipzig: Akademische Verlagsgesellschaft.
- Lloyd, A. G. (1959). *Biochem. J.* **72**, 133.
- Ramdas, A. K. (1954). *Proc. Indian Acad. Sci. A*, **39**, 81.
- Spencer, B. (1958). *Biochem. J.* **69**, 155.

The Acyl-Group Specificity of Choline Acetylase

BY J. F. BERRY* AND V. P. WHITTAKER

A.R.C. Institute of Animal Physiology, Babraham, Cambridge

(Received 13 March 1959)

Although much is known about the specificity of the cholinesterases, that of choline acetylase, the enzyme responsible for the synthesis of acetylcholine, has received relatively little attention. The

* Present address: Biochemistry Research Division, Department of Medicine, Sinai Hospital of Baltimore, Inc., Baltimore, Md., U.S.A.

alkyl-group specificity of purified rat-brain choline acetylase has been investigated by Burgen, Burke & Desbarats-Schonbaum (1956) with a number of choline analogues; limited studies of the acyl-group specificity of the enzyme in pigeon brain have been made by Gardiner & Whittaker (1954) and Whittaker (1953) and in invertebrates by