2. Glutamic, malic and β -hydroxybutyric dehydrogenases, together with alkaline deoxyribonuclease (DNase I), were found to be associated essentially with the true mitochondria. In addition, the microsomal fraction contained about 10% excess of alkaline deoxyribonuclease over what could be expected from mitochondrial contamination, and a similar excess of malic dehydrogenase was found in the final supernatant. As observed before, acid deoxyribonuclease (DNase II), which was measured here by an improved technique, showed a lysosomal distribution.

3. About two-thirds of the easily detachable iron of liver tissue was recovered in the final supernatant. The remainder was distributed diffusely amongst all particulate fractions, with a significant peak in the light mitochondrial fraction, which is richest in lysosomes. The iron content of this fraction was correlated with its acid phosphatase activity. These results support the contention, based on electron-microscope observations, that particles loaded with ferritin are concentrated together with lysosomes.

4. After administration of iron, the iron content of all fractions was augmented; the greatest increase occurred in the nuclear and heavy mitochondrial fractions, in which the iron was probably present in the form of gross haemosiderin granules.

This work was supported by grants from the Centre National de Recherches sur la Croissance normale et pathologique, the Centre National de Recherches Enzymologiques, the Rockefeller Foundation and the Lilly Research Laboratories. One of us (D.S.B.) is indebted to

the Belgian Government for a scholarship and to the Administrators of the H.E. Durham Fund, King's College, Cambridge, for a grant.

REFERENCES

- Allard, C., de Lamirande, G. & Cantero, A. (1957). *Exp. Cell Res.* **13**, 69.
 Appelmans, F., Wattiaux, R. & de Duve, C. (1955). *Biochem. J.* **59**, 438.
 Baudhuin, P. (1959). *Arch. int. Physiol. Biochim.* **67**, 106.
 de Duve, C. & Beaufay, H. (1957). *Arch. int. Physiol. Biochim.* **65**, 160.
 de Duve, C., Pressman, B. C., Gianetto, R., Wattiaux, R. & Appelmans, F. (1955). *Biochem. J.* **60**, 604.
 Gabrio, B. W., Shoden, A. & Finch, C. A. (1953). *J. biol. Chem.* **204**, 815.
 Hogeboom, G. H. & Schneider, W. C. (1953). *J. biol. Chem.* **204**, 233.
 Horecker, B. L. & Kornberg, A. (1948). *J. biol. Chem.* **175**, 385.
 Kitzes, G., Elvehjem, C. A. & Schnette, H. A. (1944). *J. biol. Chem.* **155**, 653.
 Kuff, E. L. & Dalton, A. J. (1957). *J. Ultrastruct. Res.* **1**, 62.
 Lehninger, A. L. & Greville, G. D. (1953). *Biochim. biophys. Acta*, **12**, 188.
 Novikoff, A. B., Beaufay, H. & de Duve, C. (1956). *J. biophys. biochem. Cytol.* **2**, Suppl. 179.
 Rotherham, J., Schottelius, D. D., Irvin, J. L. & Irvin, E. M. (1956). *J. biol. Chem.* **223**, 817.
 Shack, J. (1957). *J. biol. Chem.* **226**, 573.
 Stimson, M. M. & Reuter, M. A. (1945). *J. Amer. chem. Soc.* **67**, 487.
 Thiers, R. E. & Vallee, B. L. (1957). *J. biol. Chem.* **226**, 911.
 Watson, D. (1952). *J. med. Lab. Tech.* **10**, 3.
 Wattiaux, R. & de Duve, C. (1956). *Biochem. J.* **63**, 606.

Tissue Fractionation Studies

13. ANALYSIS OF MITOCHONDRIAL FRACTIONS FROM RAT LIVER BY DENSITY-GRADIENT CENTRIFUGING

By H. BEAUFAY,* D. S. BENDALL,† P. BAUDHUIN, R. WATTIAUX AND C. DE DUVE
Department of Physiological Chemistry, University of Louvain, Belgium

(Received 24 March 1959)

In the investigations described in this paper an attempt has been made to analyse in a more detailed manner the distribution of various enzymes in mitochondrial fractions from rat liver, with the help of density-gradient centrifuging. The results obtained, which have been briefly reported before (Beaufay, Baudhuin & Bendall, 1958; Baudhuin, 1959), fully confirm the distinction between mitochondria and lysosomes and

provide additional information about both groups of particles.

Density-gradient centrifuging, especially when applied to cell particles, requires specialized equipment and fairly elaborate mathematical treatments. These aspects, as well as the theoretical bases of the method, have recently been examined in detail by de Duve, Berthet & Beaufay (1959). To avoid lengthy duplication of material, we must refer to the above mentioned paper for all details which are not strictly essential to the understanding of the present account.

* Chargé de recherches du FNRS.

† Present address: Department of Biochemistry, Cambridge.

METHODS

Density-gradient centrifuging

All experiments were performed on washed mitochondrial fractions prepared according to Appelmans & de Duve (1955). When the gradients were made with deuterium oxide, the last washing was carried out with 0.25M-sucrose in heavy water and followed by centrifuging at 800 000 *g.* min. Equal quantities of the particles were added to two solutions of different density and these two suspensions were then used to make a linear gradient with the help of a twin-syringe machine. Thus at the beginning of the experiment the particles themselves were always distributed homogeneously within the tube whereas other components were not. In most experiments the equivalent of 0.6 g. of original tissue was suspended in this manner in a total volume of 5 ml. The tubes had a cross-sectional area of 1.22 cm.² and the column of fluid could thus be assimilated (neglecting the sphericity of the bottom) to a cylinder 4.1 cm. high. In calculating the physical properties of the suspension media, it was assumed that particles from 1 g. of liver occupy a volume of 0.25 ml. in 0.25M-sucrose and that they lose 0.075 and 0.125 ml. of water respectively in 0.5M-sucrose and in highly hyperosmotic sucrose, as a result of osmotic dehydration (Tedeschi & Harris, 1955).

The tubes were centrifuged with the swinging-bucket SW-39 rotor in a Spinco model L preparative ultracentrifuge equipped with special devices designed to ensure slow acceleration and deceleration, to prevent wobbling of the rotor at low speed and to record the angular velocity.

After centrifuging, the column of fluid was divided into approximately ten successive portions, of height varying between 0.2 and 1.5 cm. depending on the observed distribution of material. A special cutter was used for this purpose and the amount of fluid separated by the knife was sucked off each time, weighed and then diluted with a known volume of a solution containing mM-sodium bicarbonate, mM-ethylenediaminetetra-acetic acid and 0.01% of Triton X-100. The volume of the undiluted fractions was calculated by dividing their weight by the average density corresponding to their position in the gradient. Thus the precise dilution could be known for each fraction. In addition, their theoretical volume was estimated from the height of the cut and the cross-sectional area of the tube. As a rule, the difference between theoretical and measured volume, representing losses due to leakage and surface contamination of the cutter and pipette, amounted to about 10%. The sediment adhering to the bottom of the tube was also recovered and resuspended in a known volume of the diluent described above.

All these operations were performed with the equipment described by de Duve *et al.* (1959) and at a temperature only slightly above the freezing point of the solvent (0° for water and 3.82° for deuterium oxide). Some degree of heating of the rotor during prolonged centrifuging at high speed could, however, not be prevented. The conditions of centrifuging are expressed by the time integral of the squared angular velocity: $W = \int_0^t \omega^2 dt$.

Analytical procedures

Cytochrome oxidase was measured as described by Appelmans, Wattiaux & de Duve (1955). Glutamic and

malic dehydrogenase as well as alkaline deoxyribonuclease were assayed according to Beaufay, Bendall, Baudhuin & de Duve (1959). Except in one case, the back reaction (oxidation of reduced diphosphopyridine nucleotide) was used in all glutamic dehydrogenase assays. Uricase was determined as reported by de Duve, Pressman, Gianetto, Wattiaux & Appelmans (1955), but at 37° and in the presence of 0.1% of Triton X-100 to increase the sensitivity of the assay. For the lysosomal hydrolases, the methods of Wattiaux & de Duve (1956) were adapted to the use of long incubation periods (24 hr.) and small amounts of enzyme. All substrate mixtures contained 0.01% of sodium thiomersalate (Eli Lilly and Co., Indianapolis, Ind., U.S.A.) to prevent bacterial contamination. With acid phosphatase and β -glucuronidase, the preliminary fixation and filtration were omitted and the analytical procedures were carried out immediately. For cathepsin and the nucleases, the usual procedure was followed. Protein was determined by the method of Lowry, Rosebrough, Farr & Randall (1951).

Calculation and presentation of results

As a rule, the results are given in the form of a graph and a complementary table. In the graphs, the data are expressed in terms of the ratio of the concentration found (*C*) to the initial concentration (*C_i*) and plotted graphically as a function either of the mean distance separating the analysed fraction from the meniscus or of the corresponding density. The concentration *C* was obtained by multiplying the result of the analysis performed on the diluted sample by the dilution factor determined as explained above. The initial concentration *C_i* was calculated by dividing the total amount recovered by the volume of suspension. To compute the total amount recovered, each value of *C* was multiplied by the corresponding theoretical volume and these products were added together and with the total amount found in the sediment. By calculating *C_i* in this manner, the ratios *C/C_i* were automatically corrected for faulty recoveries.

The amounts found in the sediments and in some of the top fractions are not represented in the graphs, since they correspond to discontinuities in the gradient. They are given separately in the tables, together with the actual recoveries. These values are expressed as percentages of the total amounts determined on a mixture of equal parts of the two suspensions used to prepare the gradients.

RESULTS

Sedimentation velocity

In these experiments, the density of the suspension medium was kept below that of the particles at all levels and the centrifuging conditions were chosen so as to cause incomplete sedimentation of the particles, the gradient serving simply a stabilizing purpose. The gradient chosen extended linearly from 0.25 to 0.5M-sucrose over a radial distance of 5.5–9.6 cm. In this system, as shown by de Duve *et al.* (1959), the velocity of sedimentation of osmotically active particles having the same equilibrium density (1.22, see below) as rat-liver

mitochondria is practically independent of their position and given in first approximation by

$$\frac{dx}{dt} = 5.75 \cdot S_s \cdot \omega^2, \quad (1)$$

in which x is the radial distance from the axis, S_s is the standard sedimentation coefficient (in 0.25 M-sucrose at 0°) and ω is the angular velocity.

By integration and with the meniscus at 5.5 cm., (1) gives:

$$x - 5.5 = 5.75 \cdot S_s \int_0^t \omega^2 dt = 5.75 \cdot S_s \cdot W. \quad (2)$$

Fig. 1. Sedimentation boundaries of particle-bound enzymes. Linear gradient of 0.25–0.5 M-sucrose over radial distance of 5.5–9.6 cm. $W = 2.63 \times 10^8 \text{ rad.}^2 \text{ sec.}^{-1}$. Complementary data are given in Table 1. ●, Cytochrome oxidase; ○, glutamic dehydrogenase; ▼, alkaline deoxyribonuclease; ×, acid phosphatase; +, β-glucuronidase; ■, cathepsin; ▽, acid deoxyribonuclease; □, protein.

Table 1. Complementary data on experiment of Fig. 1

Values given refer to the activity of the sediment and to the sum of the activities of all the fractions, expressed in percentages of the total activity included in the gradient.

Enzyme	Percentage of total activity in gradient	
	Sediment	Total recovery
Cytochrome oxidase	71.6	108.0
Glutamic dehydrogenase	65.2	100.5
Alkaline deoxyribonuclease	54.9	88.2
Acid phosphatase	34.6	104.0
β-Glucuronidase	40.5	96.1
Cathepsin	40.1	98.6
Acid deoxyribonuclease	45.6	99.3
Protein	56.7	107.2

The results of three such experiments performed under comparable conditions, and of a fourth carried out with about twice the total force, are shown in Figs. 1–4 and in Tables 1–4. In Fig. 5 are represented the sedimentation-coefficient distribution curves, obtained by differentiating the observed sedimentation boundaries according to de Duve *et al.* (1959). An approximate size scale, calculated on the assumption that the particles are spherical and have a density of 1.10 in 0.25 M-sucrose, is appended for comparison with the results of other workers. For reasons which are

Table 2. Complementary data on experiment of Fig. 2

Values given refer to the activity of the sediment and to the sum of the activities of all the fractions, expressed in percentages of the total activity included in the gradient.

Enzyme	Percentage of total activity in gradient	
	Sediment	Total recovery
Cytochrome oxidase	51.3	98.2
Glutamic dehydrogenase	50.0	96.8
Acid phosphatase	25.4	96.3
β-Glucuronidase	31.1	93.3
Cathepsin	25.2	89.7
Acid deoxyribonuclease	24.5	79.6
Uricase	26.4	96.5
Protein	43.7	92.1

Fig. 2. Sedimentation boundaries of particle-bound enzymes. Experimental conditions were as in Fig. 1. $W = 2.69 \times 10^8 \text{ rad.}^2 \text{ sec.}^{-1}$. Complementary data are given in Table 2. ●, Cytochrome oxidase; ○, glutamic dehydrogenase; ×, acid phosphatase; +, β-glucuronidase; ■, cathepsin; ▽, acid deoxyribonuclease; ▲, uricase; □, protein.

explained in the discussion, the losses evidenced by the decrease below unity of the plateau values of C/C_i have been attributed to agglutination or adsorption artifacts rather than to rapid sedimentation after side-wall impactation. Consequently, in the computation of the distribution curves, the geometrical factor γ (de Duve *et al.* 1959) has been taken equal to zero and the ordinate values have been corrected for losses by multiplying by the inverse of the plateau value of C/C_i . Thus in view of equation (2):

Abscissa: $S_s = \frac{(x-5.5)}{5.75W}$ (3)

Ordinate: $F(S_s)$
 $= \frac{dC}{C_i dS_s} = \frac{dC}{C_i dx} \cdot 5.75 \cdot \frac{C_i}{C(\text{plateau})}$ (4)

Fig. 3. Sedimentation boundaries of particle-bound enzymes. Experimental conditions were as in Fig. 1. $W = 2.70 \times 10^8 \text{ rad.}^2 \text{ sec.}^{-1}$. Complementary data are given in Table 3. ●, Cytochrome oxidase; ×, acid phosphatase; △, acid ribonuclease.

Table 3. Complementary data on experiment of Fig. 3

Values given refer to the activity of the sediment and to the sum of the activities of all the fractions expressed in percentages of the total activity included in the gradient.

Enzyme	Percentage of total activity in gradient	
	Sediment	Total recovery
Cytochrome oxidase	61.5	101.5
Acid phosphatase	34.3	101.4
Acid ribonuclease	44.1	84.2

It is clear from the results obtained in these experiments that the enzymes studied may be divided into two groups: a heavier, mitochondrial group comprising cytochrome oxidase, glutamic dehydrogenase and alkaline deoxyribonuclease, and a more slowly sedimenting one including uricase and the five lysosomal acid hydrolases. The shape of the distribution curves varies little from one enzyme to the other in the former group; it is more variable in the latter, being sharpest for uricase and most spread out for acid ribonuclease and β -glucuronidase. The comparisons of Fig. 5

Table 4. Complementary data on experiment of Fig. 4

Values given refer to the activity of the sediment and to the sum of the activities of all the fractions, expressed in percentages of the total activity included in the gradient.

Enzyme	Percentage of total activity in gradient	
	Sediment	Total recovery
Cytochrome oxidase	85.0	96.1
Glutamic dehydrogenase	85.1	102.2
Acid phosphatase	42.4	99.4
β -Glucuronidase	51.1	91.5
Cathepsin	46.5	91.9
Acid deoxyribonuclease	48.4	99.7
Uricase	33.6	80.0

Fig. 4. Sedimentation boundaries of particle-bound enzymes. Experimental conditions were as in Fig. 1. $W = 5.03 \times 10^8 \text{ rad.}^2 \text{ sec.}^{-1}$. Complementary data are given in Table 4. ●, Cytochrome oxidase; ○, glutamic dehydrogenase; ×, acid phosphatase; +, β -glucuronidase; ■, cathepsin; ▽, acid deoxyribonuclease; ▲, uricase.

also make it possible to judge the reproducibility of the method which, though not perfect, may, however, be considered as fairly satisfactory.

Density equilibration

In these experiments, the limits of the gradient were chosen so as to encompass the equilibrium density of the particles. Centrifuging was pursued until, as predicted by de Duve *et al.* (1959), particles of the size of mitochondria had moved up or down the tube to a position corresponding closely to their own density. Particles of smaller size were not expected to be equilibrated as completely, but the duration of centrifuging necessary to achieve this purpose would have been prohibitive.

In Fig. 6 and Table 5 are given the results of such an experiment performed in aqueous sucrose,

and the results of two others, in which deuterium oxide was used as solvent, are shown in Figs. 7 and 8 and in Tables 6 and 7. In all three experiments, the mitochondrial group of enzymes stands out clearly from the others and shows almost perfect parallelism with the total proteins. The five lysosomal hydrolases show similar flattened curves which are progressively shifted towards higher densities in the order: acid phosphatase, β -glucuronidase, cathepsin, acid ribonuclease and acid deoxyribonuclease. These differences are not very marked in water but are exaggerated considerably in deuterium oxide, which causes a much greater shift of the equilibrium density of lysosomes than of mitochondria. This shift is particularly marked for uricase, which also shows a sharper curve than the lysosomal hydrolases.

As shown by the results of Table 8, it is possible by an appropriate choice of conditions to separate in heavy water a sediment practically free of mitochondria and in which the lysosomal enzymes and

Fig. 5. Sedimentation-coefficient distribution curves derived from Figs. 1-4 as described in the text. —, Curves of Fig. 1; — —, curves of Fig. 2; ----, curves of Fig. 3; — — —, curves of Fig. 4.

Fig. 6. Density equilibration in a linear gradient of 1.42-1.88M-sucrose in water over a radial distance of 5.5-9.6 cm. $W = 1.2 \times 10^{11}$ rad.² sec.⁻¹ (2 hr. at 39 000 rev./min.). Complementary data are given in Table 5.

Table 5. Complementary data on experiment of Fig. 6

Values given refer to the activities of the top fraction and of the sediment, and to the sum of the activities of all the fractions, expressed in percentages of the total activity included in the gradient.

Enzyme	Percentage of total activity in gradient		
	Top fraction	Sediment	Total recovery
Cytochrome oxidase	3.4	0.1	104.5
Glutamic dehydrogenase	2.4	0.2	—
Alkaline deoxyribonuclease	5.0	1.3	123.6
Acid phosphatase	7.8	3.6	94.0
Acid deoxyribonuclease	6.1	13.9	87.3
Uricase	1.5	24.4	74.5

Fig. 7. Density equilibration in a linear gradient of 0.87-1.46 M-sucrose in D₂O over a radial distance of 5.5-9.6 cm. $W = 1.8 \times 10^{11}$ rad.² sec.⁻¹ (3 hr. at 39 000 rev./min.). Complementary data are given in Table 6.

Fig. 8. Density equilibration in a linear gradient of 0.95-1.61 M-sucrose in D₂O over a radial distance of 5.5-9.6 cm. $W = 1.8 \times 10^{11}$ rad.² sec.⁻¹ (3 hr. at 39 000 rev./min.). Complementary data are given in Table 7.

Table 6. Complementary data on experiment of Fig. 7

Values given refer to the activities of the top fraction and of the sediment, and to the sum of the activities of all the fractions, expressed in percentages of the total activity included in the gradient.

Enzyme	Percentage of total activity in gradient		
	Top fraction	Sediment	Total recovery
Cytochrome oxidase	1.4	0.1	122.0
Glutamic dehydrogenase	1.1	0.2	109.0
Alkaline deoxyribonuclease	6.0	6.1	142.5
Acid phosphatase	8.3	4.8	98.2
Acid deoxyribonuclease	7.2	29.2	123.0
Uricase	0	33.5	59.5
Protein	3.0	2.5	130.7

Table 7. Complementary data on experiment of Fig. 8

Values given refer to the activities of the top fraction and of the sediment, and to the sum of the activities of all the fractions, expressed in percentages of the total activity included in the gradient.

Enzyme	Percentage of total activity in gradient		
	Top fraction	Sediment	Total recovery
Cytochrome oxidase	4.2	0	—
Malic dehydrogenase	4.0	0.1	92.2
Acid phosphatase	12.2	3.1	102.7
beta-Glucuronidase	13.1	3.5	100.3
Cathepsin	11.1	5.1	99.7
Acid ribonuclease	7.6	11.2	102.8
Acid deoxyribonuclease	7.9	14.0	93.5
Uricase	1.7	14.6	95.7
Protein	4.1	1.6	98.0

Table 8. *Isolation of a lysosome-rich sediment*

Sediment from mitochondrial fraction centrifuged for 3 hr. at 39 000 rev./min. ($W = 1.8 \times 10^{11}$ rad.² sec.⁻¹) in linear gradient of 0.9–1.51 M-sucrose in D₂O over radial distance of 5.5–9.6 cm. Values given refer to the activity of the sediment and to the sum of the activities of all the fractions, expressed in percentages of the total activity included in the gradient.

Enzyme	Percentage of total activity in gradient		Percentage of recovered activity Percentage of recovered protein
	Sediment	Total recovery	
Cytochrome oxidase	0.1	92.5	0.24
Acid phosphatase	9.7	98.2	22.4
β -Glucuronidase	13.0	92.5	31.8
Cathepsin	10.9	91.9	26.7
Acid ribonuclease	10.8	95.8	25.5
Acid deoxyribonuclease	11.9	80.9	33.3
Uricase	35.4	102.4	78.2
Protein	0.4	90.4	1.0

particularly uricase are purified to a considerable extent. Remembering that the original preparation used in this experiment contained about 70 % of the total tissue content of these enzymes purified approximately threefold, we find that the hydrolases of the sediment were purified 75- to 100-fold with a 10 % yield, and uricase some 250-fold with a 25 % yield.

DISCUSSION

Technical and theoretical considerations

The experiments reported in this paper are the first to be performed according to the principles set forth in the theoretical paper of de Duve *et al.* (1959) and it is therefore of interest to examine the results closely for all deviations from the predicted behaviour.

Sedimentation velocity. The gradient chosen is one in which it was predicted, on the basis of an approximate formula taking into account osmotic effects, that particles having an equilibrium density of 1.22 should sediment at a speed depending only on the angular velocity of the centrifuge and independent of their position. The consequence of this is that the plateau concentration below the boundary should depend only on the fate of the particles which hit the wall of the tube as a result of the radial direction of the centrifugal force. If these particles continue sedimenting at the same rate as the freely moving ones, the plateau concentration should remain unchanged; it should decrease as sedimentation proceeds, if the particles slide down rapidly to the bottom as soon as they have touched the tube wall, and should increase if they trail behind.

Unfortunately, the plateau concentration found in different experiments is not the same, but varies between 0.8 and 1.0 times the initial concentration. This may possibly be due to the presence of some agglutinated material in the preparations or to the co-sedimentation of adsorbed lysosomal enzymes

together with the mitochondria. Such artifacts should lower the plateau concentration, and since the latter may in some cases reach almost unity we conclude, in agreement with Kuff, Hogeboom & Dalton (1956), that impact with the side-wall probably has little effect on the rate of sedimentation of the particles. Accordingly, in computing the differentiated curves of Fig. 5 we have introduced no geometrical correction ($\gamma = 0$, de Duve *et al.* 1959), but have considered instead the plateau concentration as representing the 100 % base line of the population analysed.

In our opinion, these differentiated curves have some advantages in that they provide a more easily grasped representation of the experimental results and give a direct estimate of the range of sedimentation coefficients or sizes characterizing each particle population. However, too much significance should not be attached to their exact shape, since the boundaries themselves are not known with enough precision to permit accurate differentiation. In addition, it should be stressed that the curves apply to the distribution of enzymic activity and not that of the particles themselves, being directly applicable to the latter only if each particle of a given population contains the same amount of enzyme, a rather improbable assumption.

In design, though not in some of the calculations, our method is similar to that used by Kuff *et al.* (1956) in comparable experiments. In contrast, Thomson and his co-workers (Thomson & Mikuta, 1954; Thomson & Moss, 1956; Thomson & Klipfel, 1957, 1958) have applied another technique, in which the particles are layered above the gradient. The latter technique has the advantage that it furnishes the differentiated pictures as the direct experimental result. However, the thickness of the layer causes an artificial broadening of the bands and the existence of an interface may be responsible for some artifacts, as shown by Anderson (1955).

Density equilibration. Both by their high viscosity and by their very high osmotic pressure, aqueous sucrose solutions are far from ideal in isopycnic experiments. Our principal aim in using heavy water was to reduce the concentration of sucrose required for equilibration. This aim has been largely fulfilled with the mitochondria, but less with the lysosomes and uricase-containing particles owing to an unexpected increase of the equilibrium density of the particles in heavy water. According to de Duve *et al.* (1959), the equilibrium density is theoretically independent of the density of the solvent to the extent that the density of the framework of the particle is itself independent of the density of the solvent. Consequently, the observed deviations from theoretical behaviour are presumably due to the presence of exchangeable bound water of hydration in this framework. The hydration ratio must be particularly high for the uricase-containing particles, for which the peak equilibrium density is 1.228 in water and 1.273 in deuterium oxide. On the other hand, the mitochondria either contain very little bound water or their bound water does not exchange as rapidly. This latter interpretation could explain the second mitochondrial peak which is seen in heavy water and which could be due to particles in which the exchange has taken place, possibly as the result of some damage.

Properties of particles

Wherever comparison is possible, our results are in agreement with those of Thomson and co-workers (Thomson & Mikuta, 1954; Thomson & Moss, 1956; Thomson & Klipfel, 1957, 1958) and of Kuff *et al.* (1956). They fully confirm the distinction between mitochondria and lysosomes and provide additional information about the properties of these particles.

Mitochondria. As shown by the graphs of Figs. 9–11, in which have been included the fractionation results described by Beaufay *et al.* (1959), there is a remarkable correlation between the content of mitochondrial subfractions in glutamic dehydrogenase, malic dehydrogenase or alkaline deoxyribonuclease and their content in cytochrome oxidase. The same correlation exists between cytochrome oxidase and total protein in the fractions separated by gradient centrifuging (Fig. 12), except that in this case the regression lines do not pass through the origin, owing to the existence of an approximately constant amount of extraneous proteins in all subfractions. From Fig. 12 it may be concluded that between 80 and 90 % of the proteins of the analysed fractions belonged to mitochondria and that in all the mitochondrial subgroups which were separated in these

experiments, the specific activity of cytochrome oxidase, and consequently also of glutamic dehydrogenase, malic dehydrogenase and alkaline deoxyribonuclease, were essentially the same. In view of the numerous factors, size, density, osmotic behaviour, uptake of deuterium oxide, which intervene in determining the distributions observed, such results provide strong additional support to the hypothesis that mitochondria are essentially homogeneous in enzymic content (de Duve *et al.* 1955; de Duve, 1957).

Lysosomes. The five lysosomal hydrolases show typical differences from the mitochondrial group as well as obvious similarities in behaviour, which further confirm the contention that they belong to a distinct group of particles. The enzymic heterogeneity of this group, which contrasts with the homogeneity of the mitochondria, has been repeatedly discussed (de Duve *et al.* 1955; de Duve, 1957; Beaufay & de Duve, 1959) and is again evident in the present experiments. The greatest differences observed are in the ratio of acid phosphatase to acid deoxyribonuclease, which may vary from 1.5 in the less dense fractions to 0.2 in the denser ones.

Obviously, it is possible that such differences reflect the existence of distinct classes of lysosomes, perhaps as many as there are lysosomal enzymes. But it is equally possible that these particles are intrinsically heterogeneous with respect to their enzyme content, for reasons connected, for instance, with their mode of formation and of breakdown within the cells. In this respect, a comparative study of the lysosomes and of the presumably related phagosomes of Straus (1958) should prove very rewarding. A morphological study of the purified sediments which can be isolated in heavy water might also throw some light on the problem.

Uricase-containing particles. Uricase presents another vexing problem which has been discussed before. According to de Duve *et al.* (1955), this enzyme belongs either to the insoluble framework of the lysosomes or to yet another class of particles. With the new evidence available, we can only repeat this statement, with perhaps a slightly greater emphasis on the second possibility. One feature of the uricase-containing particles which has been noted before and stands out clearly in all the present experiments is their relative homogeneity with respect to sedimentation coefficient and density. This is in rather sharp contrast with the heterogeneity of the lysosomes. In addition, the uricase particles sediment somewhat more slowly and are denser, especially in heavy water, than the average lysosomes. Actually, the density-distribution curve of uricase in deuterium oxide-sucrose differs to such an extent from that of acid

phosphatase that there can be little doubt about the actual dissociation of the two enzymes. However, it differs less from that of acid deoxyribonuclease, and the problem of the relationship between uricase and lysosomes must therefore remain open.

Fig. 9. Correlation between relative content of fractions in glutamic dehydrogenase and cytochrome oxidase. ●, Fractionation experiments of Beaufay, Bendall, Baudhuin & de Duve (1959), percentage of activity of 'E + N'; ○, sedimentation-velocity measurements in density gradient, $(C \times 100)/C_i$; +, density equilibration in water-sucrose, $(C \times 100)/(C_i \times 6)$; ×, density equilibration in D_2O -sucrose, $(C \times 100)/(C_i \times 6)$.

Fig. 10. Correlation between relative content of fractions in malic dehydrogenase and cytochrome oxidase. ●, Fractionation experiments of Beaufay *et al.* (1959), percentage of activity of homogenate ('E + N'); ×, density equilibration in D_2O -sucrose, $(C \times 100)/(C_i \times 6)$.

Fig. 11. Correlation between relative content of fractions in alkaline deoxyribonuclease and cytochrome oxidase. ●, Fractionation experiment of Beaufay *et al.* (1959), percentage of activity of homogenate ('E + N'); ○, sedimentation-velocity measurements in density gradient, $(C \times 100)/C_i$; +, density equilibration in water-sucrose, $(C \times 100)/(C_i \times 6)$; ×, density equilibration in D_2O -sucrose, $(C \times 100)/(C_i \times 6)$.

Fig. 12. Correlation between content of fractions in protein and cytochrome oxidase. ○, Sedimentation-velocity measurements in density gradient, $(C \times 100)/C_i$; ×, density equilibration in D_2O -sucrose, $(C \times 100)/(C_i \times 6)$. Regression lines calculated by the method of mean squares were practically the same whether one or the other variable was taken as independent (correlation coefficients were 0.99 and 0.98). Lines drawn are means between the two calculated lines. Note that the upper line (○—○) has co-ordinate values six times larger than the lower line (×—×) and, if drawn on the same scale, would be indistinguishable from the first sixth of the latter.

SUMMARY

1. Mitochondrial fractions from rat liver have been analysed by means of density-gradient centrifuging. Two types of experiments were performed. In one, sedimentation boundaries and the derived sedimentation-coefficient distribution curves were determined after incomplete sedimentation in a stabilizing gradient. In the other, solutions of high density were used to make the gradient and the particles were allowed to approach density equilibrium with the medium by prolonged centrifuging at high speed. In each case, the distributions of up to nine different enzymes were studied simultaneously.

2. Cytochrome oxidase, glutamic dehydrogenase, malic dehydrogenase and alkaline deoxyribonuclease showed practically identical distributions under all conditions applied. The bulk of the proteins of the preparation showed the same distribution pattern, which undoubtedly was that of the true mitochondria. These particles appeared to be remarkably homogeneous with respect to their enzymic content.

3. The five lysosomal hydrolases, acid phosphatase, β -glucuronidase, cathepsin, acid ribonuclease and acid deoxyribonuclease, formed another distinct group, characterized by a lower range of sedimentation coefficients, a greater dispersion of density and a higher average density, especially in heavy water, than the mitochondria. The distribution curves of these enzymes were similar but not identical, indicating that if lysosomes form a single species, they are not enzymically homogeneous.

4. Uricase showed many properties in common with the lysosomal enzymes, but differed from them by having sharper distribution curves and a particularly large increase in equilibrium density in heavy water.

This work has been supported by grants from the Centre National de Recherches sur la Croissance normale et pathologique, the Centre National de Recherches Enzymologiques, the Rockefeller Foundation and the Lilly Research Laboratories. One of us (D.S.B.) is indebted to the Belgian Government for a scholarship and to the Administrators of the H.E. Durham Fund, King's College, Cambridge, for a grant. The authors also acknowledge gratefully the kind help and co-operation of Dr J. Berthet.

REFERENCES

- Anderson, N. G. (1955). *Exp. Cell Res.* **9**, 446.
 Appelmans, F. & de Duve, C. (1955). *Biochem. J.* **59**, 426.
 Appelmans, F., Wattiaux, R. & de Duve, C. (1955). *Biochem. J.* **59**, 438.
 Baudhuin, P. (1959). *Arch. int. Physiol. Biochim.* **67**, 106.
 Beaufay, H., Baudhuin, P. & Bendall, D. S. (1958). *Abstr. 4th int. Congr. Biochem., Vienna*, p. 71.
 Beaufay, H., Bendall, D. S., Baudhuin, P. & de Duve, C. (1959). *Biochem. J.* **73**, 623.
 Beaufay, H. & de Duve, C. (1959). *Biochem. J.* **73**, 604.
 de Duve, C. (1957). *Symp. Soc. exp. Biol.* **10**, 50.
 de Duve, C., Berthet, J. & Beaufay, H. (1959). *Progr. Biophys.* **9**, 325.
 de Duve, C., Pressman, B. C., Gianetto, R., Wattiaux, R. & Appelmans, F. (1955). *Biochem. J.* **60**, 604.
 Kuff, E. L., Hogeboom, G. H. & Dalton, A. J. (1956). *J. biophys. biochem. Cytol.* **2**, 33.
 Lowry, O. H., Rosebrough, N. J., Farr, A. L. & Randall, R. J. (1951). *J. biol. Chem.* **193**, 265.
 Straus, W. (1958). *J. biophys. biochem. Cytol.* **4**, 541.
 Tedeschi, H. & Harris, D. L. (1955). *Arch. Biochem. Biophys.* **58**, 52.
 Thomson, J. F. & Klipfel, F. L. (1957). *Arch. Biochem. Biophys.* **70**, 224.
 Thomson, J. F. & Klipfel, F. L. (1958). *Exp. Cell Res.* **14**, 612.
 Thomson, J. F. & Mikuta, E. T. (1954). *Arch. Biochem. Biophys.* **51**, 487.
 Thomson, J. F. & Moss, E. M. (1956). *Arch. Biochem. Biophys.* **61**, 456.
 Wattiaux, R. & de Duve, C. (1956). *Biochem. J.* **63**, 606.

Yield of Oxidative Phosphorylation Associated with the Oxidation of Succinate to Fumarate

By P. GREENGARD,* K. MINNAERT, E. C. SLATER AND I. BETEL
*Molteno Institute, University of Cambridge and Laboratory of Physiological Chemistry,
 University of Amsterdam*

(Received 4 May 1959)

Studies of oxidative phosphorylation with succinate as substrate have a special interest, since succinate enters the respiratory chain at a point closer to oxygen than other substrates, and pyridine nucleotide is not involved in the reaction. Unfortunately, since fumarate is readily oxidized by mitochondria

which carry out oxidative phosphorylation, it is often difficult to study phosphorylation coupled to the oxidation of succinate to fumarate without complications caused by the further oxidation of the fumarate.

Phosphorylation coupled with the oxidation of succinate to fumarate was first shown by Belitzer & Tsiabkova (1939), who added arsenite to prevent

* Present address: Geigy Research Laboratories, P.O. Box 430, Yonkers, New York, U.S.A.