molecular weight. The basic molecular structure of the substances is probably similar.

The consistent appearance of the 280–290 m μ . band of low intensity is of some interest, since a similar low intensity band is characteristic of kitol, a dimeric polymer of vitamin A, described by Embree & Shantz (1943*a*) and was also observed in the sub-vitamin A isolated from shark-liver oil by Embree & Shantz (1943*b*). A structural relationship between these substances is indicated.

The influence of prewashing with 85 % ethanol in producing large amounts of anhydro-vitamin A from the column is unexpected, since 'cyclization' of vitamin A is usually produced by the presence of acid in anhydrous solvents. After 85 % ethanol washing, however, the columns have a high moisture content, and it is evident that there is a water-acid relationship which profoundly affects the decomposition and which, under certain conditions, leads to 'cyclization' of vitamin A.

For analytical purposes, the 90 % ethanol washed column is the most useful. By its use, with one liquid chromatogram, vitamin D can be separated from tachysterol and vitamin A. β -Carotene is destroyed on the column and does not interfere.

SUMMARY

1. Methods for the quantitative separation of vitamins D from interfering sterols and vitamin A have been developed.

2. The methods involve removal of provitamins and precipitable sterols by differential solubility and precipitation with digitonin under controlled conditions and a single chromatographic analysis on flordin earth.

3. The destruction of vitamin A on flordin earth has been studied.

The author wishes to thank Mr D. J. Outred and Mr D. O. Singleton who have rendered valuable and patient assistance with the experimental work.

REFERENCES

- Awapara, J., Mattson, F. H., Mehl, J. W. & Deuel, H. J. (1946). *Science*, **104**, 602.
 Brockmann, H. (1936). *Hoppe-Seyl. Z.* **241**, 104.
 Dam, H. (1928). *Biochem. Z.* **194**, 777.
 De Witt, J. B. & Sullivan, M. X. (1946). *Industr. Engng Chem. (Anal. ed.)*, **18**, 117.
 Embree, N. D. & Shantz, E. M. (1943*a*). *J. Amer. chem. Soc.* **65**, 910.
 Embree, N. D. & Shantz, E. M. (1943*b*). *J. Amer. chem. Soc.* **65**, 906.
 Emmerie, A. & Engels, C. (1939). *Rec. Trav. chim. Pays-Bas*, **58**, 283.
 Ewing, D. T., Kingsley, G. V., Brown, R. A. & Emmett, A. D. (1943). *Industr. Engng Chem. (Anal. ed.)*, **15**, 301.
 Ewing, D. T., Powell, M. J., Brown, R. A. & Emmett, A. D. (1948). *Anal. Chem.* **20**, 317.
 Green, J. (1951). *Biochem. J.* **49**, 36.
 Gudlet, I. (1941). *Proc. Sci. Inst. Vitamins Research U.S.S.R.* **3**, no. 1, 35.
 Milas, N. A., Heggie, R. & Raynolds, J. A. (1941). *Industr. Engng Chem. (Anal. ed.)*, **13**, 227.
 Müller, P. B. (1947). *Helv. chim. Acta*, **30**, 1172.
 Nield, C. N., Russell, W. C. & Zimmerli, A. (1940). *J. biol. Chem.* **136**, 73.
 Raoul, X. & Meunier, P. (1939). *C.R. Acad. Sci., Paris*, **209**, 546.
 Thaysen, Th. E. H. (1914). *Biochem. Z.* **62**, 115.

Studies on the Analysis of Vitamins D

3. EXPERIMENTS ON THE SEPARATION OF VITAMIN D FROM VITAMIN A AND OTHER SUBSTANCES BY CONDENSATION TECHNIQUES AND SELECTIVE IRRADIATION

By J. GREEN

Walton Oaks Experimental Station, Vitamins Ltd., Tadworth, Surrey

(Received 18 September 1950)

Proposals for the analytical separation of vitamins D from interfering substances include the method of selective condensation with maleic or citraconic anhydrides and that of removing interference by selective destruction with filtered ultraviolet radiation. The first method has been recommended for the removal of vitamin A in fish-liver oils and also for the removal of tachysterol in synthetic irradiation

products. The second method has been recommended for the destruction of vitamin A.

Vitamin A, carotenoid substances in general, the vitamins D, tachysterol and some other sterols condense, under suitable conditions, with maleic and citraconic anhydrides to give inactive substances capable of easy elimination. Windaus & Lüttringhaus (1931) stated that vitamin D reacted very

slowly with maleic anhydride. Kawakami (1935) separated vitamin A from fish-liver oil concentrates by condensing acetylated material with maleic anhydride in benzene for 15 hr. Hamano (1935) used a similar method. The most important analytical proposal has been due to Milas, Heggie & Reynolds (1941), who put the method to quantitative use in a projected spectrophotometric assay for vitamin D in fish-liver oils. After removing sterols by freezing, the vitamin A-containing concentrate was condensed in dioxan (freed from peroxides) with maleic anhydride for 1 hr. On spectroscopic evidence, elimination of vitamin A was claimed, but no experimental work on the effect on pure vitamin D was described.

The removal of vitamin A in concentrates by selective irradiation has been suggested by several authors and has been used in the spectroscopic analysis of vitamin A, for example by Chevallier & Chabre (1934), Neal, Haurand & Luckmann (1941), Neal & Luckmann (1944) and Little (1944). Most workers have used unfiltered radiation, but Little showed that the effective wavelengths for vitamin A destruction were between 300 and 400 m μ ., and he stated that all other wavelengths should be filtered out. He claimed selective destruction of vitamin A under these conditions. The effect of these radiations on small amounts of vitamin D does not appear to have been adequately studied.

EXPERIMENTAL AND RESULTS

Materials and methods

(1) *Vitamin A, vitamins D, and sterols.* For specification, see Parts 1 and 2 (Green, 1951 *a, b*). (2) *1:4-Dioxan.* Freed from peroxides by standing for 24 hr. over sodium amalgam and then distilling from the same material, collecting the fraction boiling at 99–102°. (3) *Maleic anhydride.* Prepared freshly by distillation from P₂O₅. (4) *Citraconic anhydride.* Redistilled.

Vitamin A was determined by the SbCl₃ colour reaction and vitamins D by the SbCl₃-acetyl chloride reagent of Nield, Russell & Zimmerli (1940) and the ICl₃ titration method of Green (1951 *a*).

Condensation with maleic and citraconic anhydrides

The condensation method was that described by Milas *et al.* (1941). However, in addition to the maleic anhydride-dioxan condensation used by these authors, condensations were also investigated, using anaesthetic ether and benzene as media and substituting citraconic anhydride for maleic anhydride. The vitamin A or sterol-containing material was saponified where necessary, and then dissolved in 5 ml. of the condensation solvent. The condensing agent (0.2–0.5 g.) was added and the mixture heated under reflux on a water bath for 1 hr. After cooling, 15 ml. of 0.5N-ethanolic KOH and, after standing 5 min., 20 ml. of water was added. Uncondensed material was extracted several times into ether and the ethereal extract washed twice with water and dried with sodium sulphate. After evaporation of the solvent, the residue was analysed by the methods referred to above.

Vitamin A. The claims of Milas *et al.* (1941) for their method could not be substantiated. Experimental trials on the dried non-saponifiable fraction of a high potency fish-liver oil (140,000 i.u./g.) in which the dioxan-maleic anhydride condensation was used showed that the elimination of vitamin A was not complete. These results are shown in Table 1. When initial quantities of vitamin A of the

Table 1. *Condensation of vitamin A with maleic anhydride in dioxan*

(Starting material, fish-liver oil (140,000 i.u./g.) saponified.)

Original amount of vitamin A (i.u.)	Vitamin A recovered (i.u.)
10,000	200
14,000	350
140,000	7,000
200,000	16,000

order 10,000–15,000 i.u. were used, 200–400 i.u. of vitamin A were recovered. When larger amounts of starting material were used, elimination was even less satisfactory, up to 10 % of the original vitamin A

Table 2. *Condensation of vitamin A in ether and benzene*

Starting material	Original amount of vitamin A (i.u.)	Solvent	Condensation agent	Vitamin A recovered (i.u.)
Fish-liver oil saponified	10,000	Benzene	Maleic anhydride	350
Fish-liver oil saponified	10,000	Ether	Maleic anhydride	700
Molecularly distilled concentrate	10,000	Benzene	Maleic anhydride	250
Molecularly distilled concentrate	10,000	Ether	Maleic anhydride	400
Molecularly distilled concentrate	10,000	Benzene	Citraconic anhydride	400
Molecularly distilled concentrate	10,000	Ether	Citraconic anhydride	550
Fish-liver oil saponified	140,000	Benzene	Maleic anhydride	1,200
Fish-liver oil saponified	140,000	Ether	Maleic anhydride	20,000
Fish-liver oil saponified	100,000	Benzene	Maleic anhydride	950

being recovered unchanged. Similar unsatisfactory results were obtained, even after removal of sterols from the vitamin A concentrate, by freezing in methanol as recommended by Milas *et al.* (1941).

A series of condensations was carried out by refluxing in ether and benzene, and the results are expressed in Table 2. Maleic and citraconic anhydrides were employed, and, in addition to the non-saponifiable fraction of a high potency fish-liver oil, a molecularly distilled concentrate of vitamin A (1.015×10^6 i.u./g.) was used. When the original quantity of vitamin A was 10,000–15,000 i.u., maleic anhydride gave slightly better results than citraconic anhydride, benzene being a somewhat better medium than ether. In all these experiments, over 250 i.u. of unchanged vitamin A were recovered. When over 100,000 i.u. of vitamin A were taken, benzene was clearly the best medium for reaction, elimination of vitamin A in this solvent exceeding that obtained in dioxan or ether. Condensation of these large amounts of vitamin A in ether was ineffective, 10,000–20,000 i.u. of vitamin A being regularly recovered.

Increasing the amount of condensation agent and increasing the time of reaction to 1.5 hr. had very little effect on improving the elimination of vitamin A.

Vitamin D and other sterols. Milas *et al.* (1941) described no experiments on the effect, under their analytical conditions, of the maleic anhydride-dioxan condensation on pure vitamin D. Pirlot & Rougir (1947) described a method for the analysis of calciferol in ergosterol-irradiation resins, in which they claimed, on empirical grounds, removal of tachysterol by condensation with citraconic or maleic anhydrides under reflux in benzene for 30 min. They found that, under these conditions, calciferol was stable to the condensation reagent and gave unchanged values with the SbCl_3 -acetyl chloride reagent after treatment.

Table 3 shows the results obtained by effecting maleic anhydride condensation with pure calciferol and vitamin D_3 , using the same conditions as for vitamin A. There was considerable difference in the behaviour of calciferol and vitamin D_3 , the latter being more vigorously attacked by the reagent. From the reaction in ether, 85% of unchanged vitamin D_3 was recovered and over 90% of calciferol. When benzene was used as a condensation solvent, as described by Pirlot & Rougir, only about 80% of calciferol was recovered and 60–65% of vitamin D_3 . Shortening the reaction time in benzene to 30 min. did not increase the recoveries. Condensation with maleic anhydride in dioxan, contrary to the empirical findings of Milas *et al.* (1941), was found to destroy large amounts of calciferol, and vitamin D_3 was destroyed almost quantitatively by this method.

Dioxan-maleic anhydride condensations were conducted on lumisterol, tachysterol, pure suprasterol II, and a sample of mixed suprasterols. Recovered sterols in the end products were determined by iodine trichloride titration. These results are also shown in Table 3. Tachysterol was always completely destroyed. Lumisterol was found to resist attack to a greater extent than calciferol, 50% of unchanged lumisterol being recovered after 75 min. reaction time. The suprasterols, which have non-conjugated structures, were recovered quantitatively after reaction times up to 2 hr.

Separation of vitamins A and D by selective irradiation

Irradiation procedure. The source of ultraviolet radiation was a Hanovia S. 500 mercury lamp. The lamp was mounted in a well ventilated metal housing, fitted with an opening 3 in. in diameter, parallel to the long axis of the lamp. Behind the lamp was fitted a curved aluminium reflector. At a distance of 15 in. from the lamp, in a vertical

Table 3. *Condensation of vitamin D and sterols*

Sterol used	Sterol weight (mg.)	Solvent used	Reaction time (min.)	Sterol recovered (mg.)
Calciferol	5.00	Ether	60	4.60
Calciferol	5.00	Benzene	60	4.10
Calciferol	2.50	Benzene	60	2.06
Vitamin D_3	3.80	Ether	60	3.25
Vitamin D_3	3.80	Benzene	60	2.55
Vitamin D_3	3.80	Benzene	30	2.50
Vitamin D_3	4.00	Dioxan	60	Nil
Vitamin D_3	8.00	Dioxan	60	Nil
Calciferol	4.80	Dioxan	45	1.86
Calciferol	5.60	Dioxan	45	2.25
Calciferol	5.80	Dioxan	75	0.59
Tachysterol	5.20	Dioxan	60	Nil
Lumisterol	5.90	Dioxan	45	3.55
Lumisterol	3.90	Dioxan	75	1.95
Suprasterol II	5.40	Dioxan	75	5.35
Mixed suprasterols	10.50	Dioxan	75	10.60

direction, was a filter rack into which filters, 3×3 in. (7.5×7.5 cm.), could be placed for the selective elimination of desired wavelengths. It was not practical to use distances shorter than 15 in. from lamp to irradiation vessel when using filtered radiation, since the heat generated by the lamp had an adverse effect on the filters. A cabinet round the lamp-house screened off all light in a vertical direction except that which passed through the filters.

The irradiation vessel was a 250 ml. flat-bottomed fused-silica flask, which, in most experiments, stood directly on the surface of the filter. During irradiation, the flask and its contents warmed up to about 50° , no cooling being used.

stances in the oil, which absorb in the same range as vitamin A.

Selective ultraviolet irradiation was also efficacious in removing artifacts of vitamin A produced by cyclization and the destructive chromatographic adsorption on floridin earth (Green, 1951b). Indications were that the rates of destruction of these substances were approximately the same as the rate of destruction of vitamin A. The products after irradiation were, in all cases, inactive towards the SbCl_3 and the ICl_3 reagents.

Irradiation of calciferol and vitamin D₃. Solutions of calciferol and vitamin D₃ of varying concentration in CCl_4 were irradiated at a distance of 15 in. from the S. 500 lamp. Both types of filter were used.

Table 4. *Irradiation of vitamin A in CCl_4*

Vitamin A source	Amount of vitamin A irradiated (i.u.)	Filter	Time for total destruction (min.)
Fish-liver oil 140,000 i.u./g.	1000	Wratten 39	8.5
Fish-liver oil 140,000 i.u./g.	2000	Wratten 39	14.0
Fish-liver oil 140,000 i.u./g.	1000	Ilford 828	9.0
Fish-liver oil 140,000 i.u./g.	2000	Ilford 828	14.0
Fish-liver oil 140,000 i.u./g.	5000	Ilford 828	35.0
Molecularly distilled concentrate	1000	Ilford 828	5.5
Molecularly distilled concentrate	2000	Ilford 828	9.0

A study was made of the effect of filtered ultraviolet radiation on vitamin A and the vitamins D in dilute solution in an inert solvent. Each sample was irradiated in dilute CCl_4 solution for a standard time and the unchanged vitamins A or D determined, after evaporation of the solvent, by the appropriate method.

Irradiation of vitamin A. The volume of solution being irradiated was standardized at 20 ml. in all experiments. Amounts of vitamin A irradiated varied from 1000 to 5000 i.u. Each solution was irradiated until a test on a small portion showed that the vitamin A had been totally destroyed. Table 4 shows the results of irradiating vitamin A solutions under varying conditions. Two filters were used, an Ilford 828 filter, transmitting ultraviolet radiation only in the range 300–400 $\text{m}\mu$, and a Wratten 39 filter, transmitting in the range 300–500 $\text{m}\mu$.

No significant difference in the rates of destruction of vitamin A was found between the two filters. There was a slight increase in the rate of destruction when more concentrated solutions of vitamin A were irradiated. Rates of destruction increased as the purity of the vitamin A solution was increased. As Table 4 shows, when a molecularly distilled concentrate of vitamin A was used, shorter times were necessary for total destruction of an equivalent amount of vitamin A than with a high potency fish-liver oil; this difference is probably accounted for solely by the internal filtration effect of other sub-

Residual solutions were analysed for unchanged vitamin D by the SbCl_3 -acetyl chloride reagent. Table 5 shows the results obtained.

Table 5. *Irradiation of vitamins D*

Original wt. of vitamin D (mg.)	Filter used	Irradiation time (min.)	Vitamin D recovered (mg.)
1950 D ₃	Ilford 828	30	1040
1950 D ₃	Ilford 828	10	1930
1950 D ₃	Wratten 39	10	1960
2400 calciferol	Wratten 39	10	2360
2400 calciferol	Wratten 39	20	2010
600 D ₃	Ilford 828	10	370
350 D ₃	Ilford 828	5	120

Using about 2 mg. of either calciferol or vitamin D₃, good recoveries were obtained with both types of filter when irradiation times up to 10 min. were used. Longer irradiation times resulted in partial destruction, only 53 % of vitamin D₃ being recovered after 30 min. irradiation. Poor recoveries were obtained when smaller quantities of vitamin D were used, even when the irradiation time was as short as 5 min. Preliminary analytical treatment of vitamin D-containing materials appeared to increase the destruction of vitamin D by irradiation.

DISCUSSION

The use of selective condensation with maleic or citraconic anhydrides, which has been adopted as an analytical step in the assay of irradiation products

and fish-liver oils for vitamin D, must be treated with reserve.

The experiments which have been described have shown that the treatment is ineffective in quantitatively removing vitamin A or, alternatively, in removing enough vitamin A to eliminate interference in the vitamin D determination. The procedure of Milas *et al.* (1941) completely destroys large amounts of vitamin D₃ and over 60 % of calciferol, even when large quantities of the latter are used. The safest solvent in which to carry out the condensation is ether, but unfortunately, although good recoveries of calciferol and vitamin D₃ have been obtained with this solvent, complete elimination of vitamin A is not possible within reasonable times of reaction. Condensation in benzene, as used by Pirlot & Rougir (1947), results in up to 20 % loss of calciferol and about 40 % loss of vitamin D₃.

The method of selective ultraviolet irradiation to remove vitamin A from vitamin D does not appear to be suitable for routine use. It is possible that, under carefully controlled conditions, small amounts of vitamin A (the bulk of which has been removed by other methods) can be successfully eliminated from vitamin D-containing material. Experiments have

shown that although vitamin A can be smoothly destroyed by filtered radiation in the 300–400 m μ . range, the irradiation, under similar conditions, of small amounts of vitamin D results in inevitable losses.

SUMMARY

1. The analytical purification of the vitamins D by the methods of maleic and citraconic anhydride condensation and by selective ultraviolet irradiation has been investigated.

2. The condensation method is not suitable for quantitative removal of vitamin A, and losses of vitamins D always occur in its use. Vitamin D destruction by the maleic anhydride-dioxan method is particularly severe.

3. Selective ultraviolet irradiation successfully destroys vitamin A, but the method cannot at present be used to separate vitamin A from vitamin D.

4. Methods for the analysis of vitamin D which include either of the above methods of purification are of doubtful value.

The author wishes to thank Mr D. J. Outred for valuable assistance with the experimental work.

REFERENCES

- Chevallier, A. & Chabre, P. (1934). *Bull. Soc. Chim. biol., Paris*, **16**, 1451.
 Green, J. (1951*a*). *Biochem. J.* **49**, 36.
 Green, J. (1951*b*). *Biochem. J.* **49**, 45.
 Hamano, S. (1935). *Sci. Pap. Inst. phys. chem. Res., Tokyo*, **26**, 87.
 Kawakami, K. (1935). *Sci. Pap. Inst. phys. chem. Res., Tokyo*, **26**, 77.
 Little, R. W. (1944). *Industr. Engng Chem. (Anal. ed.)*, **16**, 288.
 Milas, N. A., Heggie, R. & Reynolds, J. A. (1941). *Industr. Engng Chem. (Anal. ed.)*, **13**, 227.
 Neal, R. H., Haurand, C. H. & Luckmann, F. H. (1941). *Industr. Engng Chem. (Anal. ed.)*, **13**, 150.
 Neal, R. H. & Luckmann, F. H. (1944). *Industr. Engng Chem. (Anal. ed.)*, **16**, 358.
 Nield, C. N., Russell, W. C. & Zimmerli, A. (1940). *J. biol. Chem.* **136**, 73.
 Pirlot, G. & Rougir, E. V. (1947). *Bull. Soc. chim. Belg.* **56**, 296.
 Windaus, A. & Luttringhaus, A. (1931). *Ber. dtsch. chem. Ges.* **64B**, 850.

Ester Formation by Yeasts

1. ETHYL ACETATE FORMATION BY *HANSENULA* SPECIES

By R. DAVIES, E. A. FALKINER, J. F. WILKINSON AND J. L. PEEL
*Medical Research Council Unit for Chemical Microbiology, Biochemical Laboratory,
 University of Cambridge*

(Received 3 October 1950)

The ability to form traces of esters is a property of many yeasts and is partly responsible for the bouquet of wines; however, comparatively few species are able to form esters in considerable amounts. Prominent amongst these are the two species of *Hansenula*, *H. anomala* and *H. saturnus*. Very little in the way of quantitative work appears to have been done on ester formation by yeasts. Gomes (1945), in a study

of the microflora of the Douro valley, reported that a strain of *H. anomala* var. *sphaerica* (Naegeli) Dekker produced 9.59 g. total esters (as ethyl acetate) from 145 g. of total reducing sugars and at the same time ethanol formation was diminished. Gray (1949) isolated derivatives of acetaldehyde, ethanol and acetic acid from the products of the fermentation of glucose by *H. anomala* (Hansen)