37.5°. In the sample of external tissue fat analysed, (+)-14-methylhexadecanoic acid constitutes not less than 0.2% of the total fatty acids. Examination of other fractions from mutton fat is in progress, and the evidence already accumulated suggests that appreciable quantities of branched-chain acids are present.

The work now reported establishes that certain branched-chain acids isolated from wool grease (Weitkamp, 1945) and presumably secreted by the sebaceous glands of the sheep are also present in the external tissue fat, and possibly in the other fatty tissues of this animal.

SUMMARY

(+)-14-Methylhexadecanoic acid has been shown to occur as a minor constituent of mutton fat.

We wish to acknowledge the assistance given in this work by Mr M. Fieldes, Soil Bureau, Department of Scientific and Industrial Research, Wellington, who kindly carried out all X-ray measurements.

To Mr M. S. Carrie, Chemist to The Hawkes Bay Farmers Freezing Company Ltd., and to the management of that Company, we are indebted for arranging supplies of suitable material for this investigation.

REFERENCES

- Anderson, R. J. (1927). *J. biol. Chem.* **74**, 525.
 Anderson, R. J. (1941). *Chem. Rev.* **29**, 225.
 Anderson, R. J. & Chargaff, E. (1929). *J. biol. Chem.* **85**, 77.
 Cason, J. & Winans, W. R. (1950). *J. org. Chem.* **15**, 139.
 Francis, F. & Piper, S. H. (1939). *J. Amer. chem. Soc.* **61**, 577.
 Ginger, L. G. (1944). *J. biol. Chem.* **156**, 453.
 Hansen, R. P. & Shorland, F. B. (1950). *J. N.Z. Inst. Chem.* **14**, 142.
 Hansen, R. P. & Shorland, F. B. (1951). *Biochem. J.* **50**, 207.
 Hansen, R. P. & Shorland, F. B. (1952). *Biochem. J.* **50**, 358.
 Hansen, R. P., Shorland, F. B. & Cooke, N. J. (1951a). *Chem. & Ind.* p. 839.
 Hansen, R. P., Shorland, F. B. & Cooke, N. J. (1951b). *J. N.Z. Inst. Chem.* (in the Press).
 Hansen, R. P., Shorland, F. B. & Cooke, N. J. (1952). *Biochem. J.* **50**, 581.
 Hilditch, T. P. (1947). *The Chemical Constitution of Natural Fats*, 2nd ed. revised, p. 392. London: Chapman and Hall.
 Hofmann, K. & Lucas, R. A. (1950). *J. Amer. chem. Soc.* **72**, 4328.
 Lecky, H. S. & Ewell, R. H. (1940). *Industr. Engng Chem.* (Anal. ed.), **12**, 544.
 Nunn, J. R. (1951). *J. chem. Soc.* p. 1740.
 Velick, S. F. (1944a). *J. biol. Chem.* **152**, 533.
 Velick, S. F. (1944b). *J. biol. Chem.* **156**, 101.
 Velick, S. F. (1947). *J. Amer. chem. Soc.* **69**, 2317.
 Velick, S. F. & English, J. (1945). *J. biol. Chem.* **160**, 473.
 Weitkamp, A. W. (1945). *J. Amer. chem. Soc.* **67**, 447.
 Weitzel, G. & Thomas, K. Private communication, citing *Hoppe-Seyl. Z.* (1951), **288**, 251.

Seasonal Variations in Fatty Acid Composition of New Zealand Butterfat

BY R. P. HANSEN AND F. B. SHORLAND

*Fats Research Laboratory, Department of Scientific and Industrial Research,
 Wellington, New Zealand*

(Received 6 February 1952)

New Zealand butterfat has been shown by Cox & McDowall (1948) to exhibit seasonal variations in softening points, iodine values, saponification values and Reichert values. The characteristic trend of variation of each of these properties was observed to be remarkably uniform in the four different years reviewed, and in butterfats from different localities. Interpretation of the precise nature of these changes, however, requires a knowledge of the fatty acid composition and the glyceride structure of the different butterfats.

Many workers have determined the fatty acid composition of butterfat to elucidate its nature (Hilditch & Jones, 1929; Jack, Henderson & Hinshaw, 1946), to study the effect of diet on the

milk fat (Hilditch & Sleightholme, 1930; Hilditch & Thompson, 1936; Hilditch & Jasperson, 1943) or to find the factors responsible for seasonal changes (Hilditch & Jasperson, 1941). A great deal of information has thus been accumulated, but so far as we are aware no systematic investigations have been made similar to the work now reported. It comprises both triplicate analyses of butterfat samples taken from the same locality at regular intervals over a whole year, and statistical evaluation of the results.

The purpose of this investigation was to measure statistically at regular intervals the fluctuations in the fatty acid composition of New Zealand butterfat and to attempt correlation of these changes in

chemical composition with the findings of Cox & McDowall (1948). Six butterfat samples, representing butter churned at two-monthly intervals, were analysed in triplicate to provide data for the comparison of fatty acid composition at regular intervals over the period of 1 year. For the comparison of fatty acid composition in corresponding months of different years, triplicate analyses of a further two samples are submitted. Still another analysis serves to indicate the composition of the fat in the relatively dry month of March. Of the ten butterfat samples described in this paper, nine were supplied by the Rangitaiki Plains Dairy Co. Ltd., Whakatane, and were representative of the bulk churning of cream from numerous outdoor pasture-fed herds in the district. (Jerseys constitute no less than 85% of New Zealand's dairy-cow population, and have resulted from crossing Jersey sires with Shorthorn, Friesian and Ayrshire cows (New Zealand Department of Agriculture, 1948).) The tenth sample (B/1) was supplied by the Dairy Research Institute and represented an April 1945 churning from a mixed herd of outdoor pasture-fed cattle at Massey Agricultural College, Palmerston North.

As reported in a paper on the accuracy of the ester fractionation analyses of butterfat (Hansen, 1951), the procedure adopted in these analyses does not presume to provide confirmation of the absolute accuracy of results. A measure of the degree of reproducibility of results, however, has been established by analysing all samples in triplicate and determining statistically the standard deviation for each constituent acid within each sample and over all samples. Statistical assessment has also been made of the significance of differences between butterfats churned in different months of the same year and between those churned at corresponding months of different years.

EXPERIMENTAL

All analyses recorded in this paper have been carried out by the ester-fractionation method, based on the conventional procedure of Hilditch (1947) and Hilditch & Jones (1929). Minor modifications adopted in this work have been reported by Hansen (1951). The method followed was as outlined below.

The fatty acids from 750 g. butterfat were steam-distilled and the steam-non-volatile acids separated into 'liquids' and 'solids' by crystallization three times from 15 vol. acetone at -30° . The steam-volatile acids were fractionated in a 60×1.7 cm. electrically heated and packed column of the type described by Longenecker (1937), while the 'liquids' and 'solids' respectively were converted to methyl esters and distilled *in vacuo* in a 90×3.5 cm. packed fractionating column, electrically heated in three sections, as described by Jack & Henderson (1945). The efficiency of this latter column was measured at equilibrium using a mixture of CCl_4 and benzene, and was estimated using the

data of Coulson & Herington (1948) to possess seventeen theoretical plates.

Calculations involved in the resolution of distilled fractions into components have been based on saponification equivalents and iodine values, and in general have followed the principles adopted by Hilditch (1947). Nevertheless, to clarify interpretation of the results certain assumptions used in this work are specified below.

It has been customary (cf. Hilditch, 1947) to calculate the proportions of oleic and octadecadienoic acids from the iodine value of the C_{18} unsaturated acid fractions on the assumption that only these two acids are present, but recently it has been shown (Shorland, 1950) that in addition to diene acids there is present in New Zealand butterfat linolenic acid in somewhat similar proportions. We have, therefore, made no attempt to resolve the C_{18} unsaturated acids into their components, but instead have followed the practice which Hilditch (1947) applied to complex mixtures of unsaturated acids of the same number of C atoms, and have expressed the mean unsaturation in terms of H.

In the usual ester-fractionation technique it is not feasible to resolve the trace amounts of higher saturated and unsaturated acids. It now appears that in butterfat there is present a complex mixture of C_{30} and C_{22} unsaturated acids (Shorland & Johannesson, 1951), while the saturated acids above C_{18} , according to Helz & Bosworth (1936), are mainly C_{26} , with traces of C_{20} , C_{22} and C_{24} acids.

In determining the proportions of unsaturated acids, of equivalent greater than that of the C_{18} unsaturated components, the error was minimized by refractionation of the residue, the final residual fraction (approx. 2% of total fatty acids) containing little other than C_{30} or higher unsaturated acids, together with polymerized material. Hilditch (1947) calculated the butterfat saturated acids above C_{18} as C_{20} , but for the unsaturated acids above C_{18} he assumes a mean equivalent of 330 and describes these acids as C_{30} – C_{22} . This description is no doubt correct, but as the proportions of C_{30} and C_{22} components are unknown, there is the difficulty of assigning an exact value for the saponification equivalent. Moreover, recent investigations have established the presence of trace amounts of substances hitherto unidentified in butterfat. In this category are included certain C_{30} unsaturated acids (Shorland & Johannesson, 1951) and the various branched-chain acids (Hansen & Shorland, 1951 *a, b*; Hansen, Shorland & Cooke, 1951). One of these side-chain acids, incidentally, is a multi-branched C_{20} saturated acid. The determination of all these minor constituents is beyond the scope of present-day techniques of ester fractionation. Accordingly, for simplicity in both the saturated and unsaturated series, the acids above C_{18} were assumed to be entirely C_{30} .

In calculating those 'solid' fractions with saponification equivalents between 268 and 296, it was assumed that any unsaturation present was due to traces of methyl oleate, and that separation into 'solids' and 'liquids' by crystallization from acetone at -30° had removed from the 'solids' all the C_{18} unsaturated acids (cf. Hilditch, 1947). This assumption was based on the following evidence:

(a) The solubility curves of Foreman & Brown (1944) exclude hexadecenoic acid from the 'solids'.

(b) Only minor proportions of hexadecenoic acid (about 2%) are present in butterfat.

(c) The crystallization was carried out with precision and the proportions of unsaturated acids in the 'solids' (iodine value approx. 3.0) was low and in the fractions of molecular

weight approximating 270, almost negligible (iodine value approx. 0.3).

In resolving the steam-volatile fractions into their component fatty acids it was assumed that unsaturation when present in the fractions of lower boiling point was due to decenoic acid (C_{10}) with an iodine value of 149.6. No unsaturated acid lower in the series than C_{10} has been isolated from butterfat. Investigation of the steam-volatile residues showed that they contained unsaturated components of molecular weight greater than that of decenoic acid. As the C_{18} unsaturated acid components constitute up to 85% of the total unsaturated acids of butterfat, the practice in these analyses was to assume this higher molecular weight unsaturated material in the residues from the steam-volatile acids to be C_{18} , its presence probably being due to the slight steam-volatility of oleic acid and possibly to 'splashing' during steam distillation.

The presence of oleic acid in the volatile residues was further indicated when a methyl ester sample of this material, on repeated crystallization from acetone and from methanol, yielded a fraction with saponification equivalent 291.2 and iodine value 79.2.

DISCUSSION

In Table 1 are presented the mean values for triplicate fatty acid composition analyses of six butterfat samples churned at approximately two-monthly intervals over the period of 1 year. Accompanying these results is the standard deviation of each constituent both within analyses and overall analyses. It will be noted from Table 1 that five of the samples with churning dates 15. ix. 47, 18. xi. 47, 28. i. 48, 6. iv. 48 and 18. v. 48 are representative of one dairying season, which in New Zealand commences about mid-July. The sixth sample was churned 28. vii. 48, which date, under normal conditions, corresponds approximately with the beginning of a new season. Evidence to be presented later in this paper indicates that only minor differences exist in the fatty acid composition of butterfats derived in the same months of different years. Therefore, in order to compare a series of analyses typical of any one complete season, the July sample of 28. vii. 48 (D/34) is assumed to correspond with the July sample of a year earlier. The six samples in Table 1 are then arranged in the order of a season which commences in July and ends in May.

Seasonal variations in the proportions of the fatty acid constituents

In comparing the analyses of six butterfat samples shown in Tables 1 and 2, it will be observed that the most characteristic feature of this series of analyses is the marked change in chemical composition which occurs when the season has advanced 3-4 months. This is well illustrated in the uniform pattern of changes followed by each member of the C_6 - C_{14} group of saturated acids, and also by the C_{18} group of

Table 1. Triplicate fatty acid composition analyses of six butterfats churned at approximately two-monthly intervals over the period of a year (Mean values (moles %) and standard deviation of each constituent within analyses and overall analyses are recorded.)

Sample	Date of churning	Saturated fatty acids										Unsaturated fatty acid						Total saturated acids	C_{18} - C_{22} saturated acids
		C_4	C_6	C_8	C_{10}	C_{12}	C_{14}	C_{16}	C_{18}	C_{20}	C_{22}	C_{10}	C_{12}	C_{14}	C_{16}	C_{18}	C_{20}		
D/34	28. vii. 48	12.0 ±0.23	3.7 ±0.06	0.9 ±0.21	2.7 ±0.17	2.9 ±0.25	8.4 ±0.21	23.6 ±0.15	11.3 ±0.25	0.4 ±0.20	0.4 ±0.20	0.2 ±0.06	0.2 ±0.00	0.2 ±0.00	0.9 ±0.12	2.4 ±0.12	29.5 ±0.25	65.9 ±0.23	22.2 ±0.12
C/38	15. ix. 47	11.5 ±0.21	4.5 ±0.06	1.5 ±0.30	3.5 ±0.20	3.5 ±0.17	9.6 ±0.12	21.6 ±0.12	11.3 ±0.06	0.7 ±0.10	0.7 ±0.10	0.2 ±0.00	0.2 ±0.00	0.8 ±0.10	0.8 ±0.10	1.8 ±0.12	28.1 ±0.31	67.6 ±0.17	24.4 ±0.29
C/51	18. xi. 47	10.8 ±0.17	4.7 ±0.15	1.9 ±0.25	4.4 ±0.20	4.4 ±0.20	11.5 ±0.21	22.9 ±0.31	11.1 ±0.10	0.6 ±0.12	0.6 ±0.12	0.4 ±0.00	0.3 ±0.06	1.0 ±0.00	1.7 ±0.06	23.4 ±0.12	1.2 ±0.12	72.1 ±0.20	26.1 ±0.42
C/56	28. i. 48	11.4 ±0.35	4.3 ±0.06	1.4 ±0.32	3.6 ±0.12	3.7 ±0.15	10.6 ±0.32	23.1 ±0.10	11.2 ±0.25	0.7 ±0.10	0.7 ±0.10	0.3 ±0.06	0.3 ±0.06	1.0 ±0.06	1.7 ±0.21	25.6 ±0.21	1.1 ±0.10	70.0 ±0.17	24.4 ±0.59
D/3	6. iv. 48	9.6 ±0.23	3.9 ±0.20	1.2 ±0.20	3.0 ±0.21	3.1 ±0.32	10.4 ±0.31	22.0 ±0.10	11.9 ±0.10	0.6 ±0.12	0.6 ±0.12	0.2 ±0.06	0.3 ±0.10	1.2 ±0.12	2.0 ±0.10	29.2 ±0.17	1.4 ±0.15	65.7 ±0.23	20.7 ±0.10
D/12	18. v. 48	9.5 ±0.21	3.5 ±0.06	1.2 ±0.17	2.8 ±0.29	2.9 ±0.15	9.7 ±0.05	22.7 ±0.17	12.2 ±0.12	0.8 ±0.12	0.8 ±0.12	0.2 ±0.00	0.2 ±0.06	1.1 ±0.06	2.0 ±0.12	29.8 ±0.00	1.4 ±0.15	65.3 ±0.15	19.8 ±0.21
Overall standard deviation		±0.24	±0.11	±0.25	±0.20	±0.22	±0.22	±0.17	±0.16	±0.13	±0.04	±0.04	±0.05	±0.08	±0.13	±0.25	±0.11	±0.20	±0.33

unsaturated acids. These two groups of constituents comprise approximately 50 moles % of the total fatty acids. At the beginning of lactation the C_6 - C_{14} acids comprise 18.6 %, and the C_{18} unsaturated acids 29.5 % of the total molar content, but these proportions change to 26.9 and 23.4 % respectively in November, and then gradually return, at the end of lactation, to proportions similar to those at the beginning of the season. Although these changes in fatty acid composition involve a general compensation between the C_6 - C_{14} acids and the C_{18} unsaturated acids, the balance is not precise. For example, in July the low value for the C_6 - C_{14} saturated acids (18.6 moles %) is compensated for by high values in the C_4 and C_{16} saturated acids respectively, as well as by high values in the C_{18} unsaturated acids. In May, however, the low value for the C_6 - C_{14} saturated acids (20.1 moles %) is compensated for by a high content of stearic acid and of C_{18} unsaturated acids, the butyric acid content being at a minimum. The amount of butyric acid tends to decrease with the advance of the season, but its variations at two-monthly intervals are irregular. Most marked of the differences in C_4 content is that between the July sample (12.0 moles %) and the May sample (9.5 moles %). When, however, the butyric acid content is expressed on a weight percentage basis (cf. Table 2) a more uniform trend is illustrated and the decline in content is shown to range from 4.5 % in July to 3.5 % in May.

Palmitic acid, the major saturated component, is shown in Table 1 to be relatively constant in amount throughout the season, the highest value of 23.6 moles % occurring at the beginning of the season in July, and the lowest value of 21.6 moles % appearing less than 2 months later in September. Whereas the C_4 - C_{14} saturated acid components are characterized by a decrease in content from November to the end of the season, stearic acid is unique in that it shows a small but regular increase during this period. The total saturated acids in each of the six butterfat samples follow the same pattern of variation as the individual C_6 - C_{14} saturated acids. Statistical comparison of these latter constituents reveals the differences to be significant at the 1 % level in twelve of the fifteen comparisons (cf. Table 4). When the component saturated acids are expressed as a percentage of the total saturated acids in each sample (see Table 3), they display, unlike the unsaturated acids when similarly expressed, appreciable variations. Palmitic acid, for example, ranges from 35.8 moles % in July to 31.7 moles % in November. Butyric acid changes erratically when expressed on this basis, while the C_6 - C_{14} saturated components show a regular movement throughout the season, the contents rising to a maximum in November, and then slowly declining

Table 2. *Physical and chemical data for six butterfats representing churnings made at approximately two-monthly intervals over the period of a year*

Butterfat sample	Date of churning	Iodine value (Wijs)	C_{18} unsaturated acids (moles %)	Saponification equiv.	Saponification value (calc.)	Reichert value*	C_6 - C_{14} saturated acids (moles %)	Softening point†	C_4 acids (wt. %)	Total acids liquid at room temp. (moles %)	Total acids liquid at room temp. (wt. %)
D/34	28. vii. 48	39.9	29.5	247.8	226.4	31.1	22.2	32.7	4.5	50.7	47.3
C/38	15. ix. 47	38.4	28.1	244.4	229.6	31.7	24.4	32.3	4.3	49.9	47.4
C/51	18. xi. 47	33.6	23.4	240.5	233.3	32.0	26.1	32.7	4.2	45.4	46.4
C/56	28. i. 48	36.8	25.6	244.0	229.9	30.1	24.4	33.4	4.4	47.1	43.6
D/3	6. iv. 48	39.9	29.2	249.1	225.3	27.3	20.7	33.7	3.5	49.0	41.4
D/12	18. v. 48	40.8	29.8	249.3	225.1	25.2	19.8	34.3	3.5	48.9	46.3

* Determinations made by Dairy Research Institute, Palmerston North, New Zealand. (In some cases Reichert values were made on samples churned within a day or two of those reported here.)

† Softening points determined by Dairy Research Institute, Palmerston North, New Zealand, using the method of Barnicoat (1944).

Table 3. *Saturated acid components as percentage of total saturated acids, and unsaturated acid components as percentage of total unsaturated acids*

Sample	Saturated acid components expressed as percentage of total saturated acids										Unsaturated acid components expressed as percentage of total unsaturated acids									
	C ₄	C ₆	C ₈	C ₁₀	C ₁₂	C ₁₄	C ₁₆	C ₁₈	C ₂₀		C ₁₀	C ₁₂	C ₁₄	C ₁₆	C ₁₈	C ₂₀				
D/34	18.2	5.6	1.4	4.1	4.4	12.7	35.8	17.2	0.6		0.6	0.6	2.6	7.0	86.6	2.6				
C/38	17.0	6.6	2.2	5.2	5.2	14.2	31.9	16.7	1.0		0.6	0.6	2.5	5.6	86.7	4.0				
C/51	14.9	6.5	2.6	6.1	6.1	15.9	31.7	15.4	0.8		1.4	1.0	3.6	6.0	83.8	4.2				
C/56	16.3	6.1	2.0	5.1	5.3	15.2	33.0	16.0	1.0		1.0	1.0	3.3	5.7	85.3	3.7				
D/3	14.6	5.9	1.8	4.6	4.7	15.9	33.5	18.1	0.9		0.6	0.9	3.5	5.8	85.1	4.1				
D/12	14.5	5.4	1.8	4.3	4.4	14.9	34.8	18.7	1.2		0.6	0.6	3.2	5.8	85.3	4.0				

Table 4. *Significant differences between sample means on the basis of the 't' test, and the classification of pairs of samples whose differences are significant at 5 and 1 % level*

(Designation of samples: a, C/38 (churned 15. ix. 47); b, C/51 (churned 18. xi. 47); c, C/56 (churned 28. i. 48); d, D/3 (churned 6. iv. 48); e, D/12 (churned 18. v. 48); f, D/34 (churned 28. vii. 48).)

	Saturated fatty acids										Unsaturated fatty acids										Total saturated
	C ₄	C ₆	C ₈	C ₁₀	C ₁₂	C ₁₄	C ₁₆	C ₁₈	C ₂₀		C ₁₀	C ₁₂	C ₁₄	C ₁₆	C ₁₈	C ₂₀					
5% significant difference	0.43	0.20	0.44	0.36	0.39	0.40	0.31	0.29	0.23		0.07	0.08	0.15	0.23	0.44	0.19					C ₄ -C ₁₂ 0.35
1% significant difference	0.60	0.28	0.62	0.51	0.54	0.56	0.43	0.41	0.32		0.10	0.12	0.20	0.32	0.61	0.27					0.49
Not significant	ac de	ab df	ab de ac df ad ef ae bc cd ee	ac de df ef	ac de df ef	ae cd	bc be ce	ab ac af bc bf cf	ab bf ac cd ad ee bc bd		af de bc ef bd cd oe cf ce	af de bc ef bd cd oe cf ce	af cf bc de be ef bf cd ce	ab ac ae bc de	df ef ae bc de	ab ad ae bc de					ac
Significant at 5% but not at 1%	af cf	ac	cf	—	ad	—	ad	de	af df be cf		ae be df	ae be df	ab bd de	ad ee be od	de	ac bf ef					—
Significant at 1%	ab od ad ee ae df bc ef bd be bf	ad od ae ee af cf bc de bd ef be bf	af bd be bf	ab bf ad od ae ee af cf bc bd be	ab od ae ee af cf bc bd be bf be	ab bf ac ee ad cf af de bc df bd cf be	ab cf ac de ae df af ef bd bf cd	ad ef ae bd be cd ce df	ef		ab cf ac bd be bf ce	ab ac ad bf	ac ad ae df	af bd bf cf df ef ae od af ee be cf	ab bd ac be ad bf ae od af ee be cf	af ef bd be cd ee df					ab be ad bf ae cd ad ee af cf bc bd df ef

to a minimum in May. The C_6 - C_{14} saturated acids follow the same trends as they do when expressed as percentages of the total fatty acids.

Of the six butterfat analyses shown in Table 1, those which exhibit extreme differences are the November (C/51) and the May (D/12) samples. Statistical interpretation of the variations between these samples demonstrates significant differences between all components, except the C_{14} unsaturated acids which are present in minor amounts. Of these differences, ten of the thirteen are significant at the 1% level (cf. Table 4). The C_{18} unsaturated acids (the major constituents of butterfat) are at a minimum in November (23.4 moles %) and rise to a maximum in May (29.6 moles %), a seasonal variation of 6.2 moles %. The accentuated difference between these two samples does, however, conform to the findings of Cox & McDowall (1948), who observed that seasonal changes in New Zealand butterfats were characterized by low average iodine values in November-December (about 34.0) which rose to relatively high values in May-June (about 40.0).

While the content of C_{18} unsaturated acids in New Zealand butterfats is limited to the range of 23.4-29.8 moles % the figures presented by Hilditch & Jasperson (1941) for English butterfats vary from 25.5 to 37.7 moles %. The average of five values cited by Hilditch (1947) was 34.8 moles %. For a sample of Californian butterfat, Jack & Henderson (1945) recorded the surprisingly low figure of 17.0 moles %.

The palmitic acid content of New Zealand butterfat is relatively constant (21.6-23.6 moles %) while that of English butterfat is shown to range between 21.1 moles % (Hilditch & Paul, 1940) and 26.7 moles % (Hilditch & Jasperson, 1941). Jack & Henderson (1945) report 27.2 moles % of palmitic acid in their Californian sample, and Achaya & Banerjee (1945) show samples of Indian buffalo ghees as fluctuating from 19.0 to 31.9 moles % of this constituent. From the evidence available, it would appear that New Zealand butterfats, by comparison with English butterfats, are characterized generally by a lower content of C_{18} unsaturated acids and of palmitic acid.

Annual variations in fatty acid composition

In Table 5 are recorded the results of triplicate analyses of a further four butterfat samples. For the purpose of determining annual variations in fatty acid composition, data are shown in Table 6 representing the differences between means of analyses of butterfats churned in November 1946 (B/36) and November 1947 (C/51), and between May 1946 (C/18) and May 1947 (D/12) respectively, together with the levels at which these differences are significant.

Table 5. *Triplicate fatty acid composition analyses of four butterfat samples*

Sample	Date of churning	Saturated fatty acid										Unsaturated fatty acid						Total satur- ated	Total unsatur- ated
		C ₄	C ₆	C ₈	C ₁₀	C ₁₂	C ₁₄	C ₁₆	C ₁₈	C ₂₀	C ₁₀	C ₁₂	C ₁₄	C ₁₆	C ₁₈	C ₂₀			
B/1, 9	About 5. iv. 45	10.0	4.2	1.7	3.3	3.6	10.3	23.7	10.7	0.6	0.3	0.3	1.4	2.3	26.1	1.5	68.1	31.9	
B/1, 12		10.8	3.6	1.8	3.3	4.0	9.5	22.6	11.4	0.5	0.3	0.3	1.2	1.7	27.2	1.8	67.5	32.5	
B/1, 25		10.8	3.8	2.0	3.3	3.7	9.5	23.6	10.9	0.6	0.3	0.3	1.1	1.7	27.0	1.4	68.2	31.8	
	Mean	10.5	3.9	1.8	3.3	3.8	9.8	23.3	11.0	0.6	0.3	0.3	1.2	1.9	26.7	1.6	68.0	32.0	
B/36, 1	About 18. xi. 46	10.7	5.0	2.2	4.2	5.4	12.3	23.0	9.9	0.6	0.3	0.3	1.0	1.5	21.7	1.9	73.3	26.7	
B/36, 6		11.3	4.5	2.2	4.6	4.8	11.8	22.9	10.7	0.6	0.3	0.3	1.0	1.5	22.2	1.3	73.4	26.6	
B/36, 10		10.6	4.8	2.2	4.2	5.2	12.6	22.6	10.3	0.6	0.3	0.3	1.1	1.8	22.2	1.2	73.1	26.9	
	Mean	10.9	4.8	2.2	4.3	5.1	12.2	22.8	10.3	0.6	0.3	0.3	1.0	1.6	22.1	1.5	73.2	26.8	
C/18, 1	About 13. v. 47	10.3	3.6	1.4	2.8	3.3	9.3	21.9	11.8	0.6	0.3	0.3	0.9	1.9	29.5	2.1	65.0	35.0	
C/18, 2		10.3	3.8	1.4	2.6	3.6	9.3	21.4	11.7	0.9	0.3	0.3	0.9	1.8	29.6	2.1	65.0	35.0	
C/18, 3		10.0	4.1	1.1	2.7	3.2	9.6	21.6	11.8	0.5	0.2	0.2	1.0	2.1	29.8	2.1	64.6	35.4	
	Mean	10.2	3.8	1.3	2.7	3.4	9.4	21.6	11.7	0.7	0.3	0.3	1.0	1.9	29.6	2.1	64.8	35.2	
D/86, 1	14. iii. 49	10.5	3.9	1.1	3.4	3.8	9.9	23.2	11.9	0.4	0.3	0.3	1.0	1.6	27.0	1.7	68.1	31.9	
D/86, 3		10.6	4.2	1.2	3.0	3.5	9.4	23.7	11.6	0.6	0.3	0.3	1.2	1.8	27.1	1.5	67.8	32.2	
D/86, 4		11.0	4.2	1.2	2.8	3.5	9.7	23.3	11.5	0.4	0.3	0.3	1.1	1.9	26.9	1.9	67.6	32.4	
	Mean	10.7	4.1	1.2	3.0	3.6	9.6	23.4	11.7	0.5	0.3	0.3	1.1	1.8	27.0	1.7	67.8	32.2	

Statistical comparison of the two November samples B/36 and C/51 (cf. Table 6), shows that the variables of greatest magnitude are the C_{12} , C_{14} and C_{18} saturated acids, and the C_{18} unsaturated acids. All of these differences are significant at the 1% level. In the two May samples C/18 and D/12, the variables significant at the 1% level are the saturated acids C_4 and C_{18} , and the C_{20} unsaturated acids, which latter are present in minor amounts. It is not surprising that the C_{18} unsaturated acids show significant differences in the November samples, for these acids undergo rapid changes in content at this time of the year (cf. Table 1) and it would be considered unlikely that the period of minimum content of this constituent would fall at precisely the same time each year. With this reservation it is considered that the statistical evaluation of the differences in fatty acid composition of butterfats churned in corresponding months of different years indicates that most of the components do not fluctuate appreciably in amount, and that where significant differences do occur the percentage variations are relatively small.

Although the two comparisons referred to distinguish between fatty acid composition of two examples only, both of which represent samples churned in corresponding months of consecutive years, nevertheless, the results serve as a guide to the probable limits of annual variation.

Relationship between chemical and physical constants

As shown in Table 2, both the July and the September butterfats (D/34 and C/38) possess high iodine values of 39.9 and 38.4 respectively, while the November sample displays a sudden drop to a minimum value of 33.6, which slowly rises to give a maximum value of 40.8 in May (D/12). The corresponding C_{18} unsaturated acids follow the same trend. Although the iodine values are not directly proportional to the C_{18} unsaturated acid contents, the figures suggest that the iodine values are an indication of the amount of these acids present (cf. Bartley *et al.* 1951).

For the six butterfats considered in Table 1, the sum of the C_{18} unsaturated acids and the total saturated acids in each sample is constant within very narrow limits (95.6, 95.4, 94.9, 95.7, 95.6 and 95.5%) and constitutes an average of 95.5% of the total molar weight. Accordingly, values for the total saturated acids shown in Table 1 are as would be predicted from a knowledge of the C_{18} unsaturated content of corresponding samples. As the iodine value is an index of the C_{18} unsaturated acid content of butterfat, so also is it an index of the total content of saturated acids present. When the unsaturated acid components are expressed as a percentage of the total unsaturated acids, it is seen from Table 3

Table 6. Differences between means of samples taken at corresponding months of different years, and the level at which they are significant

Samples	Saturated fatty acids (moles %)								Unsaturated fatty acids (moles %)							
	C_4	C_6	C_8	C_{10}	C_{12}	C_{14}	C_{16}	C_{18}	C_{20}	C_{10}	C_{12}	C_{14}	C_{16}	C_{18}	C_{20}	
B/36 (churned 18. xi. 46) - C/51 (churned 18. xi. 47)	+0.07	+0.10	+0.33	-0.07	+0.73	+0.76	0.00	-0.80	+0.03	-0.10	-0.03	+0.03	-0.07	-1.34	+0.34	
Level of significance	—	—	—	—	1%	1%	—	1%	—	1%	—	—	—	1%	5%	
C/18 (churned 13. v. 47) - D/12 (churned 18. v. 48)	+0.73	+0.36	+0.10	-0.07	+0.44	-0.33	-1.07	-0.46	-0.16	+0.07	+0.04	-0.10	-0.04	-0.17	+0.67	
Level of significance	1%	5%	—	—	5%	—	1%	5%	—	5%	—	—	—	—	1%	

that the content of C_{18} unsaturated acids remains approximately constant over the period of a year, the values being 86.6 (D/34), 86.7 (C/38), 83.8 (C/51), 85.3 (C/56), 85.1 (D/3) and 85.8% (D/12). Thus, although the total amount of unsaturated acids will vary at intervals, the C_{18} unsaturated acid components constitute an approximately fixed proportion of those totals.

The saponification equivalents and iodine values, without bearing any direct mathematical relationship with each other, follow the same trends. In conformity with the graphs of Cox & McDowall (1948) saponification equivalents when converted to saponification values exhibit an interrelationship with the Reichert values, an increase in one corresponding with an increase in the other. As might be anticipated, the Reichert values are closely correlated with the proportions of C_4 – C_{12} saturated acids found to be present in the samples analysed, and are at a maximum of 32.0 in November and a minimum of 25.5 in May. In Table 2 it is shown that minimum values for both the saponification equivalents and the Reichert values occur in the May sample in which the content of C_{18} unsaturated acids is highest, and maximum values occur in the November sample in which the C_{18} unsaturated acid content is lowest. Thus changes in content of C_4 – C_{12} saturated acids are reflected in compensating changes in the content of C_{18} unsaturated acids.

No correlation can be established between the softening points and iodine values of the butterfat samples reported in this paper. It appears, however, that softening points are influenced more by the content of lower saturated acids (C_4 – C_{12}) than by the unsaturated components; high softening points corresponding with low contents of C_4 – C_{12} saturated acids and also with low Reichert values (cf. Eckles & Palmer, 1916). Attempts to correlate softening points with the total content of acids liquid at room temperature (C_4 , C_6 and C_8 saturated acids, together with all the unsaturated acids) have not established any definite relationship. From data in Table 2 it might be expected that the November sample C/51 with a content of 45.4 moles % of acids liquid at room temperature (minimum value recorded for the six samples) would possess a softening point higher than that of the other butterfats: actually its softening point is only 32.7°, which is identical with that of sample D/34 (July) with 50.7 moles % (maximum value recorded for these six samples). Conversely, if a definite relationship existed it would be expected that the May sample (D/12) with the highest softening point of 34.3° would possess the minimum value for the total acids liquid at room temperature. However, its content of liquid acids is 48.9 moles %, only 1 mole % lower than that for the September sample (C/38), which, of

the six butterfats analysed, has the lowest softening point of 32.3°. A similar lack of interrelationship exists when the acids liquid at room temperature are expressed in terms of weight per cent (cf. Table 2). The absence of correlation between the content of acids liquid at room temperature and the softening points of corresponding samples, suggests that softening points depend on the mode of fatty acid distribution throughout the glyceride molecules.

Factors influencing seasonal variations in fatty acid composition of New Zealand butterfat

Among the many factors which have been suggested as being responsible for seasonal variations in the fatty acid composition of butterfat are (a) diet, (b) plane of nutrition, (c) stage of lactation, (d) temperature of environment, (e) degree of freedom of movement, (f) age of cow, and (g) breed of cow.

Diet. The influence of diet on the fatty acid composition of butterfat has been established by a number of experiments. Hilditch & Thompson (1936) fed linseed oil, which is rich in diene and triene C_{18} acids, and observed that there was no perceptible increase in the amounts of these substances present in the milk fats. When, however, cows were fed liberal amounts of palm-kernel oil (Hilditch & Jasperson, 1943), or coconut oil (Hilditch & Sleightholme, 1930), both of which substances are rich in lauric and myristic acids, the proportions of lauric and particularly of myristic acids in the butterfat were increased. Similarly, the feeding of rape-seed oil (Hilditch & Thompson, 1936), which contains erucic acid, led to the appearance of this acid in butterfat. The fats present in grass do not amount to more than 2–4% of the dry matter (cf. Shorland, 1944), and their main constituents, linolenic and octadecadienoic acids, are found in not more than minor amounts (about 1%) in butterfat (cf. Shorland, 1950). From the ample evidence available it appears that under normal grazing conditions the influence of diet would remain substantially the same throughout the season and would not be responsible for variations in fatty acid composition of butterfat.

Of the other factors referred to, those considered to have exerted greatest influence on the samples described in this paper are the plane of nutrition and the stage of lactation.

Plane of nutrition. Investigations by Eckles & Palmer (1916), Riddet (1938), Riddet, Campbell, McDowall & Cox (1941) and by Smith & Dastur (1938) have all established that underfeeding has marked effects on the physical and chemical constants of butterfat. Smith & Dastur (1938), who supported their evidence with fatty acid composi-

tion analyses, showed that inanition led to an increased oleic acid content accompanied by a reduction in the lower saturated acids. Their fatty acid composition analyses were consistent with the results of the other investigators who observed a rise in iodine value and a decrease in Reichert value with a lowering of the plane of nutrition.

In New Zealand, cows are pasture-grazed throughout the year, the diet normally being supplemented with hay or silage during winter. Growth of pasture reaches its maximum within the months of September to January and declines slowly towards the end of the season (May-June), the amount of feed available to cows in New Zealand diminishing accordingly. If the plane of nutrition were to influence the fatty acid composition of the butterfat samples shown in Table 1 it would be anticipated that when pasture was abundant, as it normally is in November, analyses would reveal a low content of C_{18} unsaturated acids, with a corresponding high content of lower volatile acids, and that as the season advanced and food became increasingly scarce, the content of C_{18} unsaturated acids would rise. This pattern has been followed, the minimum content of C_{18} unsaturated acids appearing in November (C/51) and the peak value in May (D/34). Similarly, the lower volatile acids (C_4 - C_{12}) suffer a decline from 26.1 moles % in November to 19.8 moles % in June.

In Great Britain when the cows change over from stall-feeding to pasture-feeding in the summer the animals suffer a decline in the plane of nutrition and the iodine value rises to a maximum. It would appear that this factor may be responsible for the differences in composition observed between New Zealand and English summer butterfat. The analyses by Hilditch & Jaspersen (1941) for English summer butterfat from pasture-fed animals are in close agreement with our analyses for New Zealand winter butterfat, except that the English butterfat characteristically has a slightly higher content of C_{18} unsaturated acids. Similarly, Hilditch & Sleightholme (1930) observed a close relationship between New Zealand summer butterfat and an English (stall-fed) winter butterfat.

Stage of lactation. That the stage of lactation exerts an effect upon the composition of butterfat was reported by Eckles & Shaw (1913), Holland & Buckley (1918), and by Riddet *et al.* (1941). The former observed that the volatile acids showed a uniform decline from the beginning to the end of the lactation period, and that the iodine value underwent a gradual increase, the highest point being reached at the end of the season. Similar trends in iodine values which appeared, regardless of seasonal changes in temperature, and in which the variations in iodine value were associated largely with changes in the estimated oleic acid content of

the fat, have been reported by Bartley *et al.* (1951). Hilditch & Sleightholme (1930), when commenting on the milk-fat analyses published by Holland & Buckley (1918), also considered that the amount of unsaturated acids in butter increased with the time the cows had been in lactation.

In this present study of New Zealand butterfats, the contents of C_{18} unsaturated acids (which have been shown to be closely correlated with iodine values) do not gradually increase throughout the season as anticipated from the investigations of Eckles & Shaw (1913), but rather do they follow the trends of Bartley *et al.* (1951), who observed that milk fats possessed a high iodine value during the early stages of lactation followed by a decline (about 8 units of iodine value) to the fourth or fifth month of lactation, and then a slight increase (about 3 units of iodine value) to the end of lactation. Samples analysed in this present survey decreased in content of C_{18} unsaturated acids from 29.5 moles % at the commencement of lactation in July, to 23.4 moles % in November, and only after that date did the content of these acids increase as the season advanced, reaching a peak (29.8 moles %) at the end of lactation. Whereas Bartley *et al.* (1951) found that the iodine value of the butterfat was 5 units lower at the end of the lactation period than at the beginning, our results show that the New Zealand butterfats produced at the beginning and at the end of the lactation period have approximately the same iodine values.

The figures reported in Tables 1 and 2 are consistent with the changes in fatty acid composition resulting from the varying planes of nutrition throughout the season. Furthermore, they conform to those seasonal trends in content of C_{18} unsaturated acids observed by Bartley *et al.* (1951) to accompany the advance of lactation.

From the data now presented for New Zealand butterfat, it is felt that it is not possible to differentiate between the effects of the plane of nutrition and the stage of lactation. It is anticipated, however, that this problem will be elucidated by experiments now being conducted by the Dairy Research Institute, Palmerston North, New Zealand.

In conclusion, it is considered that the results presented in this paper illustrate the trends of the variations in content of each of the major fatty acid constituents of New Zealand butterfat during one season. The analyses further indicate that only minor changes occur in the composition of butterfat samples in the same month of different years. This information, when supported by the comprehensive data relating to the constants of butterfat over a 4-year period (Cox & McDowall, 1948), affords a guide to the seasonal changes that can be predicted in the fatty acid composition of New Zealand butterfat.

SUMMARY

1. Six butterfats taken at approximately two-monthly intervals throughout the season have been subjected to ester-fractionation analyses in triplicate and the standard deviation for each constituent has been calculated.

2. Throughout the season, beginning in July, there was found to be a regular increase in the content of C_6 - C_{14} saturated acids (and total saturated acids) up to November, and thereafter a slow decline to the end of the season. The C_{18} unsaturated acids varied in the reverse direction to that of the C_6 - C_{14} constituents.

The butyric acid content changed irregularly, but tended to decline throughout the season. Palmitic acid showed little change, but when expressed as a percentage of the total saturated acids it showed a tendency to increase throughout the season.

3. In comparing butterfats sampled at two-monthly intervals during 1 year, statistical evaluation showed that significant differences existed between many of the fatty acid constituents. Extreme differences in composition were exhibited by the May and November samples, where, of the thirteen differences, ten were assessed to be significant at the 1% level.

4. Comparing the butterfats taken during the

same month, but in different years, significant differences were observed between certain of the component fatty acids. However, the differences were small, and this fact, together with Cox & McDowall's (1948) results showing the similarity between the constants of butterfats sampled during the same months of different years, is sufficient to indicate that the present study would serve as a useful guide for predicting the fatty acid composition of New Zealand butterfat for any given month.

5. The variations in content of C_{18} unsaturated acids reflect the influence of the changing plane of nutrition throughout the year and at the same time are consistent with the changes observed by Bartley *et al.* (1951) to accompany the progression of lactation.

The authors wish to acknowledge with appreciation the assistance in experimental work provided by Miss M. D. L. White.

To Dr F. H. McDowall, Chief Chemist, Dairy Research Institute, Palmerston North, New Zealand, we are indebted for arranging all butterfat samples used in this survey and for helpful discussions.

Statistical interpretations of the results reported in this paper were made by the Applied Mathematics Laboratory, Department of Scientific and Industrial Research, Wellington.

REFERENCES

- Achaya, K. T. & Banerjee B. N. (1945). *Indian J. vet. Sci.* **15**, 271.
- Barnicoat, C. R. (1944). *Analyst*, **69**, 176.
- Bartley, E. E., Zaletel, J. H., Bird, E. W., Cannon, C. Y., Wise, G. H. & Kempthorne, O. (1951). *J. Dairy Sci.* **34**, 536.
- Coulson, E. A. & Herington, E. F. G. (1948). *Trans. Faraday Soc.* **44**, 629.
- Cox, G. A. & McDowall, F. H. (1948). *J. Dairy Res.* **15**, 377.
- Eckles, C. H. & Palmer, L. S. (1916). *Res. Bull. Mo. agric. Exp. Sta.* no. 25.
- Eckles, C. H. & Shaw, R. H. (1913). *The Influence of the Stage of Lactation on the Composition and Properties of Milk*. U.S. Dep. of Agriculture.
- Foreman, H. D. & Brown, J. B. (1944). *Oil & Soap*, **21**, 183.
- Hansen, R. P. (1951). *J. Amer. Oil Chem. Soc.* **28**, 375.
- Hansen, R. P. & Shorland, F. B. (1951). *Biochem. J.* **50**, 207.
- Hansen, R. P. & Shorland, F. B. (1952). *Biochem. J.* **50**, 358.
- Hansen, R. P., Shorland, F. B. & Cooke, N. J. (1951). *Chem. & Ind.* p. 839.
- Helz, G. E. & Bosworth, A. W. (1936). *J. biol. Chem.* **116**, 203.
- Hilditch, T. P. (1947). *The Chemical Constitution of Natural Fats*, 2nd ed. revised. London: Chapman and Hall.
- Hilditch, T. P. & Jasperson, H. (1941). *J. Soc. chem. Ind., Lond.*, **60**, 305.
- Hilditch, T. P. & Jasperson, H. (1943). *Biochem. J.* **37**, 238.
- Hilditch, T. P. & Jones, E. E. (1929). *Analyst*, **54**, 75.
- Hilditch, T. P. & Paul, S. (1940). *J. Soc. chem. Ind.* **59**, 138.
- Hilditch, T. P. & Sleightholme, J. J. (1930). *Biochem. J.* **24**, 1098.
- Hilditch, T. P. & Thompson, H. M. (1936). *Biochem. J.* **30**, 677.
- Holland, E. B. & Buckley, J. P. (1918). *J. agric. Res.* **12**, 719.
- Jack, E. L. & Henderson, J. L. (1945). *J. Dairy Sci.* **28**, 65.
- Jack, E. L., Henderson, J. L. & Hinshaw, E. B. (1946). *J. biol. Chem.* **162**, 119.
- Longenecker, H. E. (1937). *J. Soc. chem. Ind.* **56**, 199T.
- New Zealand Department of Agriculture (1948). *Primary Production in New Zealand*.
- Riddet, W. (1938). *Proc. 7th. Ann. Conf. N.Z. Grasslands Ass.*
- Riddet, W., Campbell, I. L., McDowall, F. H. & Cox, G. A. (1941). *N.Z. J. Sci. Tech.* **23**, 99A.
- Shorland, F. B. (1944). *Nature, Lond.*, **153**, 168.
- Shorland, F. B. (1950). *Nature, Lond.*, **166**, 745.
- Shorland, F. B. & Johannesson, D. L. (1951). *Nature, Lond.*, **168**, 75.
- Smith, J. A. B. & Dastur, N. N. (1938). *Biochem. J.* **32**, 1868.