REFERENCES

- Bertrand, G. (1906). *Bull. Soc. chim. Fr.* (iii), **35**, 1285.
- Birkinshaw, J. H., Charles, J. H. V., Hetherington, A. C. & Raistrick, H. (1931). *Phil. Trans. B*, **220**, 55.
- Friedel, R. A. & Orchin, M. (1951). *Ultraviolet Spectra of Aromatic Compounds*. New York: John Wiley and Sons, Inc.
- Koehler, B. (1938). *J. agric. Res.* **56**, 291.
- Lehmstedt, K. (1935). *Ber. dtsch. chem. Ges.* **68**, 1455.
- Marchionatto, J. B. (1942). *Rev. Fac. Agron. B. Aires*, **9**, 159.
- Munsen, L. S. & Walker, P. H. (1906). *J. Amer. chem. Soc.* **28**, 663.
- Munsen, L. S. & Walker, P. H. (1912). *Bull. U.S. Dep. Agric.* **107**, 240.
- Raistrick, H. & Smith, G. (1933). *Biochem. J.* **27**, 1814.
- Randall, H. M., Fowler, R. G., Fuson, N. & Dangel, J. R. (1949). *Infrared Determination of Organic Structures*. New York: D. Van Nostrand Co. Inc.
- Raper, K. B. & Thom, C. (1949). *A Manual of the Penicillia*. Baltimore: The Williams and Wilkins Co.
- Semeniuk, G. & Barre, H. J. (1944). *Rep. Ia agric. Exp. Sta.* Pt. II, 52-7.
- Smith, G. (1942). *An Introduction to Industrial Mycology*, 2nd ed. London: Edward Arnold and Co.
- Westling, R. (1911). *Ark. Bot.* **11**, 53, 88.

Separation of Tea Polyphenols on Paper Chromatograms

BY E. A. H. ROBERTS* AND D. J. WOOD

Tocklai Experimental Station, Indian Tea Association, Cinnamara, Assam, India

(Received 28 June 1952)

Previous work on the chromatography of tea polyphenols (Roberts & Wood, 1951a) was carried out with phenol and *n*-butanol-acetic acid as the solvents. It has now been shown that water is also a useful solvent for the separation of polyphenols on a paper chromatogram. In particular, results have been obtained which are most simply interpreted by assuming that the water effects a separation of optical antipodes of catechins and gallo catechins, thus enabling a provisional decision to be made as to which of the various optically active forms of these substances occur naturally in the tea leaf. The use of water as a solvent in paper chromatography also differentiates between aglycones and glycosides in the flavonol group.

A preliminary account of this work has already been presented (Roberts & Wood, 1951b).

METHODS AND MATERIALS

Chromatographic technique. This is as described by Roberts & Wood (1951a). Two-way chromatograms are carried out using water as the first solvent, and the *n*-butanol-acetic acid mixture, previously described, as the second solvent. Such a chromatogram will be referred to as a two-way water chromatogram. Ferric alum (0.2%, w/v) and ammoniacal AgNO₃ (equal volumes of 0.1 N-AgNO₃ and 5 N-NH₃ aq.) are used as sprays for the detection of polyphenols. Chlorogenic acid is detected by spraying first with 1% (w/v) NaNO₂ dissolved in 10% acetic acid (v/v), and then with N-NaOH (Roberts & Wood, 1951c). Exposure of the dry paper, before spraying, to NH₃ vapour shows up anthoxanthins and chlorogenic acid as yellow patches.

* Present address: Indian Tea Association, Chemical Laboratory, Butler's Wharf, Lafone Street, London, S.E. 1.

Polyphenols. (±)-Galocatechin, (-)-epigallocatechin and (-)-epicatechin were samples obtained from Dr A. E. Bradfield. Quercitrin was obtained from Prof. W. Baker, rutin from Dr J. F. Crouch and myricetin from Prof. T. R. Seshadri.

The (+)-catechin was Merck's commercial product. Its aqueous solution was rather dark in colour which made polarimetry difficult, but $[\alpha]_D^{26}$ was found to be +16.1° in acetone-water (50%, v/v), which leaves little doubt that it is (+)-catechin.

Quercetin was prepared from rutin by hydrolysis with 0.1% (v/v) HCl for 4 hr., and butin from the blooms of *Butea frondosa* by the method of Rupe & Schaerer (1932).

A dilute solution of (+)-gallo catechin was obtained from tea-leaf juice by the following procedure. A close row of spots of tea-leaf juice was applied along the starting line of a chromatogram. After development with *n*-butanol-acetic acid, the (+)-gallo catechin band was located by spraying guide strips. Transverse strips of the paper, containing the (+)-gallo catechin, were then eluted with ethyl acetate saturated with water. The solvent was removed *in vacuo* and the residue taken up in water.

A dilute solution containing gallo catechin-*a* gallate and catechin-*a* gallate was obtained by a similar procedure from reconstituted juice autoclaved at 130° for 1 hr. (Roberts & Wood, 1951a). The solvent used for development in this case was water.

Tannase. This was prepared from *Aspergillus niger* according to Freudenberg (1933). In later experiments a culture of *A. niger* M, obtained from Long Ashton Research Station, was used.

RESULTS

A typical two-way water chromatogram is illustrated in Fig. 1, which shows the pattern of spots for polyphenols from clone 14/3/32. This source is chosen as it contains naturally occurring gallo catechin-*a* gallate. Catechin-*a* gallate, if present,

would have occupied a position immediately below the spot for gallo catechin-*a* gallate.

Identifications of the spots on the chromatogram were carried out as in our earlier communication (Roberts & Wood, 1951*a*).

Gallo catechin-*a* gallate and catechin-*a* gallate have R_F values in water very little greater than zero, and are readily separated from the corresponding galloyl esters of (–)-epigallocatechin and (–)-epicatechin. Spot 1, provisionally, but probably erroneously, associated with *m*-digallic acid, has a much higher R_F in water than (–)-epigallocatechin so that overlapping of these two substances does not occur, as when phenol is the first

resulted in partial epimerization of catechins and gallo catechins. These changes may also be demonstrated on two-way water chromatograms. On autoclaving fresh leaf juice for 1 hr. at 130°, or refluxing for 2 hr., there is a marked decrease in the intensities of the spots associated with (–)-epigallocatechin gallate and (–)-epicatechin gallate, while spots corresponding with gallo catechin-*a* gallate and catechin-*a* gallate make their appearance. This confirms earlier findings. Spots associated with (–)-epigallocatechin and (–)-epicatechin

Fig. 1. Chromatogram of tea-leaf juice (clone 14/3/32), showing position of major polyphenolic components. The chromatogram was run first from left to right with water as solvent, and then downwards with *n*-butanol-acetic acid. R_F values with water are approximate only. Anthoxanthins detected by NH_3 vapour, other polyphenols by ammoniacal AgNO_3 . Key to polyphenols (provisional identifications): (1) unknown, previously provisionally identified as *m*-digallic acid, (2) (–)-epigallocatechin, (3) (+)-gallo catechin, (4) gallic acid, (5) gallo catechin-*a* gallate ((+)-gallo catechin gallate in this source?), (6) (–)-epigallocatechin gallate, (8) (–)-epicatechin gallate, (9) (+)-catechin, (10) (–)-epicatechin. Anthoxanthins etc. shown by dotted lines, (A4) rutin, (A11) quercitrin, (A8) chlorogenic acid, other spots unidentified as yet.

solvent. Gallic acid also has a high R_F in water which separates it well from (–)-epigallocatechin gallate, although with one batch of paper it overlapped (+)-catechin. Apart from this one inconsistency, the positions assumed by polyphenols on two-way water chromatograms have proved to be reproducible.

Effect of heat on polyphenol solutions

It was shown by Roberts & Wood (1951*a*) that heating solutions containing tea polyphenols

Fig. 2. Portion of chromatogram showing positions of flavanols not occurring naturally but produced on heating tea-leaf juice. Conditions of chromatography as in Fig. 1. Key to new flavanols produced on epimerization, (2A) (+)-epigallocatechin, (3A) (–)-gallo catechin, (9A) (–)-catechin, (10A) (+)-epicatechin (provisional identifications). The arrows indicate the direction of epimeric changes which take place on heating the solutions.

also decrease in intensity, but instead of a corresponding increase in the intensity of spots 3 and 9, two new spots (3A and 9A on Fig. 2) appear. Spots 3 and 9 also show a slight decrease, and if the autoclaving is carried out at 140° these decreases become more marked and two further new spots, 2A and 10A, make their appearance. These four new spots are never found in freshly expressed juice, nor can they be detected in autoclaved juice when phenol takes the place of water as the first solvent for chromatography.

Following the above observations it was noted that (±)-gallo catechin gives two spots, 3 and 3A, on a two-way water chromatogram, but gives a single spot only when phenol is the first solvent. This not only indicates that the spots 3 and 3A are to be identified as (+)- and (–)-gallo catechin, but also suggests that the other pairs of spots, 2 and 2A, etc., represent pairs of optical antipodes.

Epimerization of individual catechins

A solution of (-)-epicatechin (1 mg./ml.) was autoclaved at 130° for 1 hr. Its two-way water chromatogram then showed the new spot 9A in addition to spot 10, corresponding with unchanged (-)-epicatechin; spot 9 was not observed. According to Freudenberg & Purmann (1924) the main transformation product obtained from (-)-epicatechin under these conditions is (-)-catechin, although a certain amount of racemization also occurs. As the identity of (-)-epicatechin is beyond reasonable doubt (Bradfield & Penney, 1948), it follows that spot 9A is probably to be identified with (-)-catechin.

In order to epimerize (+)-catechin a procedure similar to that described by Freudenberg (1933) was carried out. (+)-Catechin (5 g.) and potassium bicarbonate (15 mg.) were dissolved in water (15 ml.) and autoclaved at 130° for 6 hr. The mass of crystals which separated on cooling was dissolved in hot water (12.5 ml.). On cooling, this solution commenced to crystallize after standing 30 min.; 5 min. later it was filtered. The filtrate, which continued to crystallize, was again filtered after 1 hr., and the second filtrate allowed to stand overnight when a third crop of crystals was obtained. In confirmation of Freudenberg's findings, chromatograms of the first two fractions showed them to consist essentially of unchanged (+)-catechin. The chromatogram of the third fraction showed the new spot 10A in addition to unchanged (+)-catechin. Addition of (-)-epicatechin as a marker gave a well-defined spot (10), which was clearly separated from spot 10A. As the main transformation product of (+)-catechin under the above conditions is (+)-epicatechin (Freudenberg & Purmann, 1924), it seems probable that this substance is responsible for spot 10A.

Epimeric changes in the gallo catechins do not appear to have been investigated. The results presented below suggest that the gallo catechins behave in a similar manner to the corresponding catechins.

After autoclaving a solution in water of (-)-epigallocatechin (1 mg./ml.) at 130° for 1 hr., the two-way water chromatogram shows two spots, one of which is associated with unchanged material, while the other, 3A, by analogy with the transformation of (-)-epicatechin, is probably due to (-)-gallo catechin.

The dilute solution of gallo catechin, isolated chromatographically from reconstituted juice, gives a single spot (3) on a two-way water chromatogram. As (-)-gallo catechin appears to be responsible for spot 3A, and (±)-gallo catechin gives spots 3 and 3A, it follows that spot 3 is probably due to (+)-gallo catechin. If the gallo catechin solution is autoclaved at 130° for 2 hr. its two-way water

chromatogram shows spots 3 and 2A. This may be taken as evidence for the identification of spot 2A with (+)-epigallocatechin.

Gallocatechin-a gallate

A solution containing gallocatechin-a gallate and small amounts of catechin-a gallate was incubated with tannase for 12 days. A two-way water chromatogram then showed a spot corresponding with gallic acid, a fairly strong spot 3A, and a much weaker spot 9A. The results of hydrolysis with tannase therefore suggest the identification of these substances as the galloyl esters of (-)-gallocatechin and (-)-catechin respectively.

Anthoxanthins and chlorogenic acid

Two-way water chromatograms of tea-leaf juice reveal a number of yellow spots when the dried paper is exposed to ammonia vapour. Of these A4 and A11 are probably due to rutin and quercitrin respectively, as addition of these flavonols to the juice intensifies these spots. Spots A1, A2, A6, A7 and A8 also correspond with spots of the same number on chromatograms run first with phenol (Roberts & Wood, 1951a). These spots are all shown on Fig. 1; other spots are also observed, but, as they have not been identified with the corresponding spots on the phenol and *n*-butanol-acetic acid chromatograms, are omitted from Fig. 1.

Spots A6, A7 and A8 differ from the others in having higher R_f values in water. They all give a pink colour reaction on spraying with nitrous acid followed by sodium hydroxide, and spot A8 is intensified if chlorogenic acid is added to the juice. Spot A8 is therefore probably due to chlorogenic acid (Roberts & Wood, 1951c), and spots A6 and A7 may be related substances.

Butin, myricetin and quercetin all have zero R_f in water. If this behaviour of aglycones is general, it appears likely that none of the anthoxanthins of tea leaf is an aglycone, as all spots giving a yellow colour with ammonia vapour on tea-leaf juice chromatograms so far observed, have R_f values in water significantly greater than zero. Confirming this it is found that all anthoxanthin spots on the chromatogram disappear if the juice is first hydrolysed with *N*-hydrochloric acid for 2 hr. at 100°. As a result of this hydrolysis other anthoxanthin spots appear on the chromatogram, one of which appears to be quercetin and possibly originates from rutin.

The zero R_f values for aglycones afford a ready method of distinguishing them from their glycosides. A rapid way of carrying out such a test is to deliver a few μ l. of an anthoxanthin solution on to filter paper so as to give a spot about 1.5 cm. in diameter. On drying the paper and exposing to ammonia vapour, the whole spot assumes a yellow

colour if the anthoxanthin is a glycoside. With an aglycone the yellow colour is much more intense and confined to a small circle a few mm. in diameter.

DISCUSSION

The use of water as a solvent in paper chromatography is of some theoretical interest, as the separations achieved are apparently due to adsorption rather than to partition effects. Many polyphenols are quite strongly adsorbed on filter paper, and such adsorption probably affects the R_f values when solvents other than water are used for chromatography.

It has not yet been possible to study the behaviour of a large number of polyphenols on paper chromatograms with water as a solvent, so that conclusions must remain tentative, but certain tendencies seem to emerge. A low R_f , that is a ready adsorption by filter paper, is most marked in substances with a flavonoid structure. Flavanols (catechins) are somewhat less strongly adsorbed than the flavonols and the one flavanone investigated, provided that these latter are not glycosidic.

Spots 10 and 10A have been provisionally identified as (-) and (+)-epicatechin respectively. The other pairs, 9 and 9A, 2 and 2A, and 3 and 3A, have also been identified provisionally as pairs of optical antipodes. If these identifications are confirmed, it follows that a separation of optical antipodes on a paper chromatogram has been achieved. The cellulose may be considered as an optically active adsorbent and, in adsorption chromatography, such separations are theoretically possible.

Table 1. *Provisional identification of polyphenols on chromatograms*

Spot no.	Polyphenol
1	Unknown, previously provisionally identified as <i>m</i> -digallic acid
2	(-)-Epigallocatechin
2A	(+)-Epigallocatechin
3	(+)-Galocatechin
3A	(-)-Galocatechin
4	Gallic acid
5	(+) or (-)-Galocatechin gallate
6	(-)-Epigallocatechin gallate
7	(+) or (-)-Catechin gallate
8	(-)-Epicatechin gallate
9	(+)-Catechin
9A	(-)-Catechin
10	(-)-Epicatechin
10A	(+)-Epicatechin
4A	Rutin
4B	Chlorogenic acid
4C	Quercitrin

The naturally occurring flavanols in tea leaf are associated with spots 2, 3, 9 and 10. Accepting it as established that (-)-epigallocatechin and (-)-

epicatechin occur naturally (Bradfield, Penney & Wright, 1947; Bradfield & Penney, 1948; Bradfield & Bate-Smith, 1950), it appears probable that (+)-galocatechin and (+)-catechin also occur naturally. The isolation of (±)-galocatechin and (±)-catechin from green tea extracts (Bradfield *et al.* 1947; Bradfield & Penney, 1948) is probably due to the relative ease of the transformations

which may be expected to take place during the working up of the extracts. Admixture of the naturally occurring (+)-catechin and (+)-galocatechin with the corresponding (-)-isomers, produced as a result of these transformations, is likely to lead to solutions from which the racemic forms would separate.

It is much more difficult to effect the transformations

and if such transformations do occur during the working up of the extracts, the resultant products are present in amounts too small to be detected chromatographically.

Owing to the low R_f in water of galocatechin-*a* gallate resolution of its optical antipodes on a paper chromatogram is not to be expected. The effect of hydrolysis with tannase, however, suggests that the galocatechin-*a* gallate, produced from (-)-epigallocatechin gallate on heating a tea infusion, is (-)-galocatechin gallate. It probably therefore originates by simple epimeric change. Although the artifact is probably (-)-galocatechin gallate, the naturally occurring galocatechin-*a* gallate found in leaf of clone 14/3/32 may differ in structure, and there is some evidence (unpublished observations) which suggests that it may be (+)-galocatechin gallate.

Identifications of spots 2, 4, 6, 8, 9 and 10 on the chromatograms have been carried out with duly authenticated specimens of the corresponding polyphenols. The pair of spots 3 and 3A has been similarly identified with (±)-galocatechin. Such identifications may be accepted with the usual reservations made in such cases. So far we have not had access to such flavanols as (-)-catechin and (+)-epicatechin which do not occur naturally, or their galocatechin analogues, so that identifications of spots 2A, 3A, 9A and 10A must be considered tentative only. The identification of spots 2A and 3A also rests on the assumption that galocatechins epimerize in the same way as the catechins. Finally, the suggestions made as to the identity of spots 5 and 7 are provisional only and require much further investigation.

SUMMARY

1. Water may be used as the mobile solvent in the paper chromatography of tea polyphenols. With water a separation of the optical antipodes of catechin, epicatechin, galocatechin and epigallocatechin appears to be achieved.

2. It is provisionally concluded that (+)-catechin and (+)-galocatechin occur naturally in the tea leaf, and that isolation of the corresponding ($\pm$) compounds is due to partial epimerizations of (–)-epicatechin and (–)-epigallocatechin during extraction.

3. Gallocatechin-*a* gallate is provisionally identified as (–)-galocatechin gallate, originating from (–)-epigallocatechin gallate by epimerization.

4. Chlorogenic acid occurs in tea leaf.

5. Aglycones of flavonols have zero R_f in water, and may thus be distinguished from their glycosides.

The authors wish to express their thanks to Dr A. E. Bradfield for useful criticisms of the manuscript, and for determining the optical activity of the (+)-catechin. Thanks are also due to Mr E. Hainsworth for assistance in growing the *A. niger*, to Mr C. J. Harrison, Chief Scientific Officer, for his continued interest, and to the Indian Tea Association for permission to publish this work.

REFERENCES

- Bradfield, A. E. & Bate-Smith, E. C. (1950). *Biochim. biophys. Acta*, **4**, 441.
Bradfield, A. E. & Penney, M. (1948). *J. chem. Soc.* p. 2249.
Bradfield, A. E., Penney, M. & Wright, W. B. (1947). *J. chem. Soc.* p. 32.
Freudenberg, K. (1933). *Tannin, Cellulose, Lignin*. Berlin: Julius Springer.
Freudenberg, K. & Purrmann, L. (1924). *Liebigs Ann.* **437**, 274.
Roberts, E. A. H. & Wood, D. J. (1951*a*). *Biochem. J.* **49**, 414.
Roberts, E. A. H. & Wood, D. J. (1951*b*). *Biochem. J.* **49**, xxxiii.
Roberts, E. A. H. & Wood, D. J. (1951*c*). *Arch. Biochem. Biophys.* **33**, 299.
Rupe, H. & Schaerer, M. (1932). In *Handbuch der Pflanzen-analyse*, by G. Klein. Wien: Julius Springer.