REFERENCES

- Spiegel-Adolf, M., Hamilton, R. H. jun., & Henny, G. C. (1942). *Biochem. J.* **36**, 825.
- Weisz, O., Cochran, W. & Cole, W. F. (1948). *Acta cryst., Camb.*, **1**, 83.

EXPLANATION OF PLATE

Powder X-ray diffraction photographs of: (a) synthetic L-triiodothyronine hydrochloride, (b) hydrochloride of pure compound isolated from thyroid gland, (c) L-thyroxine, (d) L-triiodothyronine, (e) L-diiodothyronine. Note the correspondence between patterns of (a) and (b); this is also shown quantitatively in Table 1. (f) Crystals of L-triiodothyronine hydrochloride photographed with ordinary light; (g) polarized light; (h) crystals of hydro-

chloride of compound isolated from the thyroid gland photographed with ordinary light; (i) polarized light. The direction of extinction in both the natural and synthetic crystals is parallel to the needle axis. (j) Single crystal X-ray diffraction photograph of synthetic triiodothyronine hydrochloride. Crystal oscillated 15° round the *b*-axis. Filtered Cu radiation.

3:5:3'-Triiodothyronine

2. PHYSIOLOGICAL ACTIVITY

By J. GROSS* AND ROSALIND PITT-RIVERS

National Institute for Medical Research, Mill Hill, London, N.W. 7

(Received 5 August 1952)

On establishing the identity of 3:5:3'-triiodothyronine as a normal body constituent, it was decided to investigate its biological activity. Preliminary experiments on its effect in preventing thiouracil-induced goitre (Gross & Pitt-Rivers, 1952*b*) and on human myxoedema (Gross, Pitt-Rivers & Trotter, 1952), indicated that triiodothyronine possessed marked thyroid-hormone like properties.

The present paper describes a more comprehensive survey of the biological activity of 3:5:3'-triiodo-L-thyronine both in the goitre prevention test and in the growth and organ weights of thyroidectomized rats.

Dempsey & Astwood (1943) showed that the thiouracil-induced hyperplasia of the thyroid gland was quantitatively inhibited by graded doses of thyroxine; subsequently, Reineke, Mixner & Turner (1945) demonstrated that this response could be closely correlated with the standard metabolism tests. For this reason, and also because of its simplicity, this test was chosen to determine the activity of triiodothyronine relative to thyroxine as well as to examine the activities of the DL and D forms of the new amino-acid.

* Merck Fellow in Natural Sciences, National Research Council of Canada. Present address: State University of New York Medical Center, 350 Henry Street, Brooklyn, N.Y.

METHODS

Goitre prevention assay

Male hooded rats, bred in the National Institute for Medical Research, weighing 100 ± 15 g. were housed two in a cage and fed the stock diet 41 for rats and mice (Bruce & Parkes, 1949) *ad lib.*; this diet contains approximately 1 μ g. I/g. Thiouracil was given in 0.1% solution in the drinking water. Groups of six to eight animals were used to test each dose. The test compounds (thyroxine and triiodothyronine) were dissolved in 0.1 N-NaOH and diluted with 0.9% NaCl to give the desired dose in 0.1 ml. solution; the final concentration of alkali was of the order of 0.001 N. Similar solutions omitting the test substance were used for the control groups. The doses were injected subcutaneously in the scruff of the neck. In the assay comparing the relative effectiveness of injected and oral doses the latter were diluted with 15% sucrose solution, the dose being contained in 0.2 ml. which was fed to the animals from a graduated syringe; the doses injected for comparison were given in 0.1 ml. solution.

After ten daily doses, the animals were killed with CHCl_3 , weighed, their thyroids dissected from the trachea and weighed on a torsion balance to the nearest 0.1 mg. In the first experiments the pituitaries were also removed, fixed in Susa fluid, sectioned and stained with Masson's trichrome stain.

Using the procedure described by Finney (1951), the slopes of the curves and the potencies were calculated. The potencies were expressed as percentages of the activity of equimolar amounts of L-triiodothyronine. The limits of error at the 5% level are also given for each result, for

J. GROSS AND ROSALIND PITT-RIVERS—3:5:3'-TRIODOTHYRONINE
1. ISOLATION FROM THYROID GLAND AND SYNTHESIS

example the potency derived from Fig. 1 is expressed as $20 \pm 3.5\%$, which indicates that there is less than a 5% chance that L-thyroxine has a potency greater than 23.5% or less than 16.5% that of L-triiodothyronine.

In addition to the goitre-prevention assays the following experiment was done to compare the relative effects of thyroxine and triiodothyronine in restoring the growth rate of young thyroidectomized rats and also to discover whether relatively large doses of triiodothyronine would produce signs of hyperthyroidism. Fifty male hooded rats weighing 50–60 g. were divided into five groups each containing ten animals. Four of these groups were thyroidectomized surgically as follows: the animals were anaesthetized by intraperitoneal injection of 0.1 ml. pentobarbital solution (veterinary pentobarbital diluted 10 times with saline) and anaesthesia was maintained with ether. The skin over the anterior neck region was incised in the mid-line, the thyroids were exposed by blunt separation of the subcutaneous tissues and strap muscles of the neck. The caudal pole of each thyroid lobe was grasped with a pair of fine forceps and removed by gentle traction toward the head. After bleeding had been controlled by small pledgets of cotton wool, the skin was closed with interrupted silk sutures. The fifth group of animals was treated in the same manner, except that the thyroid glands were not removed; these animals formed the group of sham-operated controls. The animals were given diet 41 *ad lib.* and for the first 2 weeks calcium gluconate (1%, w/v) in the drinking water; thereafter plain tap water.

The sham-operated group and one thyroidectomized group received the alkaline-saline control solution. The three remaining groups of thyroidectomized animals received 1 and 20 μg . L-triiodothyronine and 5 μg . L-thyroxine respectively. All solutions were made up as before and were injected subcutaneously each day for 73 days. The animals were weighed at frequent intervals. At the end of the experiment the animals were all injected intraperitoneally with 5 μg . tracer Na^{131}I and killed with CHCl_3 24 hr. later. Pituitary and adrenal glands, heart and kidneys were removed from each animal, fixed for 24 hr. in Susa's fluid and then weighed.

As a check on the completeness of the thyroidectomy, the tissues of the neck region were ground finely with powdered glass in 5 ml. water and left overnight. The suspension was then treated with an equal volume of 20% (w/v) trichloroacetic acid, and after centrifugation the precipitate was washed with 5 ml. 10% (w/v) trichloroacetic acid containing inactive iodide. The residue was dissolved in N-NaOH (0.1 ml.), diluted to 10 ml. with water and counted in a liquid counter. This procedure has been shown to eliminate almost all of the iodide present in plasma (Leblond & Gross, 1948), and on this basis any animals which showed appreciable amounts of radioactivity in the precipitate were eliminated from the series as being incompletely thyroidectomized.

RESULTS

Goitre prevention assay

Comparison of the relative potencies of L-thyroxine and L-triiodothyronine. In preliminary assays (Gross & Pitt-Rivers, 1952b) it appeared that L-triiodothyronine was 3–4 times as active as L-thyroxine on an equimolar dosage. In order to

obtain a better estimate of the relative potencies, the effect of three dose levels of each compound was determined. The doses were 1, 2 and 4 μg . of L-thyroxine and 0.16, 0.32 and 0.64 μg . of L-triiodothyronine; the molar ratio of each corresponding dose is 5:1. The results obtained are shown in Fig. 1. The relative potency calculated statistically was $20.4 \pm 3.5\%$; i.e. L-triiodothyronine possessed 5 times the activity of L-thyroxine in inhibiting the release of thyrotropin from the pituitary.

Fig. 1. Effect of graded subcutaneous doses of L-thyroxine and L-triiodothyronine on the thyroid weights of rats given 0.1% thiouracil in the drinking water. Each point represents the average of eight animals. The curves were fitted mathematically and showed that L-thyroxine was 20% as potent as equimolar amounts of L-triiodothyronine.

The effect of L-triiodothyronine on the pituitary was directly demonstrable in sections of glands taken from the rats used in this experiment (Pl. 4a–d). This plate shows that 0.64 μg . L-triiodothyronine was more effective than 1 μg . L-thyroxine in preventing the degranulation of the acidophils and the hypertrophy and vacuolation of the basophils which follows thyroid hormone deficiency (cf. Halmi, 1952).

Table 1. *Relative activities of 3:5:3'-triiodo-L-thyronine and L-thyroxine after oral and subcutaneous administration to thiouracil-fed rats*

	Thyroid wt. (mg./100 g. body wt.)
Controls	$21.8 \pm 1.9^*$
Triiodothyronine (0.64 μg . oral)	9.9 ± 2.8
Triiodothyronine (0.64 μg . subcutaneous)	7.9 ± 1.1
Thyroxine (4.0 μg . oral)	17.2 ± 3.3
Thyroxine (4.0 μg . subcutaneous)	10.1 ± 2.9

* Standard deviation about the mean.

Comparison of the oral and subcutaneous activities of L-thyroxine and L-triiodothyronine. From the data given in Table 1, it is evident (a) that both thyroxine and triiodothyronine are more effective

after subcutaneous injection than oral administration, and (b) that triiodothyronine is about twice as effective as thyroxine when given orally. Statistical evaluation of the data gives the following values: oral thyroxine has $39 \pm 26.1\%$ of the activity as the same parenteral dose; oral triiodothyronine has $86 \pm 17\%$ of the activity as the same parenteral dose.

(3) *Activities of DL- and D-triiodothyronine.* The results (Fig. 2) show that D-triiodothyronine possesses only slight activity ($7.4 \pm 2.4\%$ of the activity of L-triiodothyronine). DL-Triiodothyronine (Fig. 3) possessed $59 \pm 24\%$ of the activity of L-triiodothyronine.

Peripheral action of L-triiodothyronine

Using the potency values obtained from Fig. 1 as a basis for dosage, the relative effects of L-thyroxine and L-triiodothyronine on the growth of thyro-

parathyroidectomized rats was investigated (Fig. 4). It was considered unpractical to adjust the dose to the weight of the growing rat, so fixed daily doses of $5 \mu\text{g}$. L-thyroxine and $1 \mu\text{g}$. L-triiodothyronine were given. The rates of growth in these two groups (Fig. 4, C and E) were significantly greater ($P = < 0.001$) than the thyroidectomized controls (B), and did not differ significantly from each other. Similarly, both substances prevented the marked enlargement of the pituitary which occurred in the thyroidectomized animals (Table 2).

The administration of $20 \mu\text{g}$. triiodothyronine induced a degree of hyperthyroidism (Fig. 4, D) which inhibited growth; it also caused a significant increase in the heart and kidney weights (Table 2). Histological sections (Pl. 4e-h) of the kidney showed that the hypertrophy was due to a relative increase in the thickness of the cortex, and that of

Fig. 2.

Fig. 2. The relative effects of L- and D-triiodothyronine on the thyroid weights of thiouracil-treated rats. Each point represents the average gland weight from six animals. The calculated potency of D-triiodothyronine is 7% that of the L isomer.

Fig. 3.

Fig. 3. The relative effects of L- and DL-triiodothyronine on the thyroid weights of thiouracil-treated rats. Each point represents the average gland weight from six animals. The calculated potency of DL-triiodothyronine is 59% that of the L isomer.

Table 2. *Effect of 3:5:3'-triiodo-L-thyronine on organ weights of thyroidectomized rats*

(All weights expressed as mean $\pm$ s.d.)

Treatment	No. of rats	Pituitary (mg./100 g. body wt.)	Adrenals (mg./100 g. body wt.)	Heart (g./100 g. body wt.)	Kidneys (g./100 g. body wt.)
Sham-operated controls	8	3.6 ± 0.4	15.0 ± 0.7	0.45 ± 0.05	0.87 ± 0.06
Thyro-parathyroidectomized controls	6	$7.4 \pm 0.9^*$	22.5 ± 5.3	0.47 ± 0.06	0.85 ± 0.14
Thyro-parathyroidectomized + $1 \mu\text{g}$. triiodothyronine	7	3.9 ± 0.6	19.0 ± 3.0	0.46 ± 0.04	0.82 ± 0.05
Thyro-parathyroidectomized + $20 \mu\text{g}$. triiodothyronine	5	$4.1 \pm 0.3^\dagger$	35.9 ± 12.8	0.84 ± 0.30	1.34 ± 0.26
Thyro-parathyroidectomized + $5 \mu\text{g}$. thyroxine	6	4.2 ± 0.6	21.9 ± 5.2	0.56 ± 0.07	0.96 ± 0.06

* Mean obtained from five values.

† Mean obtained from four values.

the heart to an increase in the thickness of the ventricular wall. The adrenal glands in this group also appeared to be enlarged (Table 2), but because of the variation between animals, the difference is beyond the limit of statistical significance.

Fig. 4. Growth curves. (A) Sham-operated controls (eight animals); (B) thyroidectomized controls (six animals); (C) thyroidectomized rats (seven animals) + 1 μ g. L-triiodothyronine/day; (D) thyroidectomized rats (five animals) + 20 μ g. L-triiodothyronine/day; (E) thyroidectomized rats (seven animals) + 5 μ g. L-thyroxine/day. All doses were given subcutaneously, following thyroidectomy on the first day of the experiment.

DISCUSSION

From these investigations on the biological activity of triiodothyronine it is apparent that it can duplicate the effects of thyroid hormone on body growth, kidney and heart weight and morphology, and pituitary weight and function; it also acts as a thyroid hormone replacement in returning the respiratory and lipid metabolism of hypothyroid patients to normal (Gross *et al.* 1952). In its effect on pituitary function (i.e. the release of thyrotrophin) triiodothyronine is clearly several times as active as thyroxine (about 5 times in the rat). With regard to its peripheral action on growth and metabolism, the data obtained are at present insufficient to establish clearly a greater activity than thyroxine. There are, however, indications from the work of Tomich (personal communication) and from experiments now in progress with Thibault on the effect of triiodothyronine on basal metabolism of intact and thyro-parathyroidectomized rats, that it is significantly more active in raising oxygen consumption.

It would appear, therefore, that triiodothyronine possesses a greater amount of thyroidal activity than thyroxine; if at the same time one considers that it is a normal constituent of the thyroid gland and of extrathyroidal tissues (Gross & Leblond, 1951*a, b*; Gross & Pitt-Rivers, 1951, 1952*a*; Roche, Lissitzky & Michel, 1952) and that it is derived from thyroxine in the animal body (Gross & Leblond, 1951*a*), then it is reasonable to suppose that triiodothyronine, either alone or in combination with some non-iodinated molecule, is the form of the thyroid hormone which acts at the periphery, and that thyroxine is its precursor. The necessity for this conversion in order to obtain a peripherally active substance may also account for the failure of thyroxine itself to produce any effect on isolated tissues *in vitro*, but it would not be profitable to discuss this point in detail until an *in vitro* system has been found in which either triiodothyronine or thyroxine, or both, can exert an effect at physiological levels of dosage.

If such a conversion does in fact occur, then the daily body requirement of triiodothyronine by the rat can be calculated from the curves in Fig. 1, using the method proposed by Dempsey & Astwood (1943) for thyroxine. These authors suggested that the amount of thyroxine administered daily which could maintain a normal thyroid weight in a thiouracil-treated animal represented the normal secretion rate. In our animals the normal thyroid weight was 9.2 ± 1.3 mg./100 g. body weight, indicating that the daily body requirement of triiodothyronine would be about 0.6 μ g./100 g./day, and for thyroxine about 3.75 μ g./100 g./day. This suggests that only about one-fifth of the molecules of thyroxine produced by the rat thyroid are converted into triiodothyronine, and would account to some extent for the excretion of thyroxine in the faeces of normal rodents (Gross & Leblond, 1951*a*). It may be that the rate of the reaction thyroxine $\rightarrow$ triiodothyronine is the major factor in determining the varying levels of thyroid activity found in different species (Albert, 1949; Monroe & Turner, 1949).

The presence in thyroid tissue of a substance more potent than thyroxine would also explain some of the results obtained when the biological activity of desiccated thyroid (on the basis of the thyroxine content) is compared with the biological activity of pure thyroxine. Frieden & Winzler (1948) found that parenterally administered desiccated thyroid was about twice as active as thyroxine when assayed by the goitre-prevention test. These authors estimated the thyroxine content of the desiccated thyroid by a butanol-extraction method very similar to the one used to isolate triiodothyronine by Gross & Pitt-Rivers (1952*c*), and would therefore probably have included triiodo-

thyronine and other iodine-containing compounds in their 'thyroxine' estimate. A relatively small triiodothyronine content in the desiccated thyroid could well account for its increased potency. Further, these authors (Frieden, Tuckich & Winzler, 1949) showed that natural thyroid preparations given orally retained 70–98% of their potency by injection, while thyroxine retained only half its parenteral potency when given by mouth. This difference could again be explained, at least in part, by the finding (Table 1) that oral triiodothyronine exhibits 85% of its parenteral activity, while oral thyroxine retains only about 40% of its parenteral activity. These considerations would obviate the necessity of assuming that the greater biological activity of thyroid gland compared to thyroxine is due to the presence of peptides containing thyroxine and diiodotyrosine, as has been suggested by many authors.

It is likely from the relative effectiveness of the L and D forms of triiodothyronine (Figs. 2 and 3) that the L form occurs naturally. Preliminary experiments on the potency of the material isolated from the thyroid gland supports this view (Gross & Pitt-Rivers, unpublished).

As regards the biosynthesis of triiodothyronine, it is clear that reductive deiodination of thyroxine can occur in the body (Gross & Leblond, 1951a). In the thyroid, however, it might be suggested that triiodothyronine arose from the coupling of one molecule of diiodotyrosine and one molecule of monoiodotyrosine. Some experiments have been done to see whether such a coupling could be achieved *in vitro*, by incubating the acetyl derivatives of mono- and di-iodotyrosine together at a slightly alkaline pH (cf. Pitt-Rivers, 1948); at no time was any triiodothyronine chromatographically detectable. These experiments are in no way conclusive, but they support the suggestion originally put forward by Harington (Harington & Barger, 1927), and later elaborated (Harington, 1944), that the active principle of the thyroid results from the coupling of two molecules of diiodotyrosine; it is only then necessary to add one further step to the reaction, the deiodination of thyroxine to triiodothyronine. Roche, Michel, Michel & Lissitzky

(1951), who demonstrated the deiodination of diiodotyrosine by thyroid slices *in vitro* were unable to demonstrate any deiodination of thyroxine (Roche *et al.* 1952); nevertheless, it seems probable that a deiodinating enzyme does occur in the thyroid as well as in other tissues, which is able to effect this reduction, a reaction which we now believe is the last stage in thyroid hormone biosynthesis.

SUMMARY

1. 3:5:3'-Triiodo-L-thyronine was found to possess 5 times the activity of L-thyroxine when assayed by the goitre prevention test in rats.

2. Orally administered L-triiodothyronine possessed about 86% of the activity of the same dose given subcutaneously; oral L-thyroxine only had about 40% of the activity of the same dose given subcutaneously.

3. D-Triiodothyronine had about 7% of the activity of L-triiodothyronine after subcutaneous administration in the goitre prevention assay; DL-triiodothyronine had 59% of the activity of the L form.

4. L-Triiodothyronine (1 µg. daily) maintained a rate of growth in the thyroidectomized rat equivalent to that obtained with L-thyroxine (5 µg. daily). A large dose of triiodothyronine (20 µg. daily) reduced the rate of growth and resulted in enlargement of the kidney cortex, heart and possibly the adrenal glands.

5. L-Triiodothyronine (0.64 µg. daily) was more effective than L-thyroxine (1 µg. daily) in preventing the enlargement of the basophil cells and degranulation of the acidophil cells in the pituitary gland of the thiouracil-treated rat. L-Triiodothyronine (1 µg. daily) also prevented an enlargement of the pituitary in the thyro-parathyroidectomized rat.

6. From the above evidence it is suggested that triiodothyronine is the peripheral thyroid hormone and that thyroxine is its precursor.

We wish to thank Miss M. V. Mussett for the statistical analyses; we also thank Mr P. Bovill for help with animal experiments, Mr F. Higginson for the histological sections, Mr C. Sutton and Mr E. Hitchcock for the photographs and Mr M. R. Young for the photomicrographs.

REFERENCES

- Albert, A. (1949). *Ann. N.Y. Acad. Sci.* **50**, 466.
 Bruce, H. M. & Parkes, A. S. (1949). *J. Hyg., Camb.*, **47**, 202.
 Dempsey, E. W. & Astwood, E. B. (1943). *Endocrinology*, **32**, 509.
 Finney, D. J. (1951). *J. gen. Microbiol.* **5**, 223.
 Frieden, E., Tuckich, E. B. & Winzler, R. J. (1949). *Endocrinology*, **45**, 82.
 Frieden, E. & Winzler, R. J. (1948). *Endocrinology*, **43**, 40.
 Gross, J. & Leblond, C. P. (1951a). *Proc. Soc. exp. Biol., N.Y.*, **76**, 686.
 Gross, J. & Leblond, C. P. (1951b). *Endocrinology*, **48**, 714.
 Gross, J. & Pitt-Rivers, R. (1951). *Lancet*, **2**, 766.
 Gross, J. & Pitt-Rivers, R. (1952a). *Lancet*, **1**, 439.
 Gross, J. & Pitt-Rivers, R. (1952b). *Lancet*, **1**, 593.
 Gross, J. & Pitt-Rivers, R. (1952c). *Biochem. J.* **53**, 645.
 Gross, J., Pitt-Rivers, R. & Trotter, W. R. (1952). *Lancet*, **1**, 1044.
 Halmi, N. S. (1952). *Endocrinology*, **50**, 140.
 Harington, C. R. (1944). *Proc. Roy. Soc. B*, **132**, 223.

J. GROSS AND ROSALIND PITT-RIVERS—3:5:3'-TRIODOTHYRONINE
2. PHYSIOLOGICAL ACTIVITY

- Harington, C. R. & Barger, G. (1927). *Biochem. J.* **21**, 169.
- Leblond, C. P. & Gross, J. (1948). *Endocrinology*, **43**, 306.
- Monroe, R. A. & Turner, C. W. (1949). *Res. Bull. Mo. agric. Exp. Sta.* no. 446.
- Pitt-Rivers, R. (1948). *Biochem. J.* **43**, 223.
- Reineke, E. P., Mixner, J. P. & Turner, C. W. (1945). *Endocrinology*, **36**, 64.
- Roche, J., Lissitsky, S. & Michel, R. (1952). *C.R. Acad. Sci., Paris*, **234**, 1228.
- Roche, J., Michel, R., Michel, O. & Lissitsky, S. (1951). *C.R. Soc. Biol., Paris*, **145**, 288.

EXPLANATION OF PLATE

Representative sections from the central portions of the pituitaries of rats used in the experiment shown in Fig. 1. (Masson Trichrome $\times 950$ green filter.) The sections show the appearance of the pituitary of the thiouracil-treated animal after (a) no further treatment, (b) L-thyroxine (1 $\mu\text{g./day}$) and (d) L-triiodothyronine (0.64 $\mu\text{g./day}$) compared with the gland of the normal animal (c). The appearance of enlarged basophils (pale cells) and decreased number of darker appearing acidophils resulting from thiouracil treatment is partially prevented by 1 $\mu\text{g. L-thyroxine}$ (b) and completely prevented by 0.64 $\mu\text{g. L-}$

triiodothyronine (d). Representative sections of the kidneys (upper) and hearts (lower) of the animals whose growth curves are given in Fig. 5 (haematoxylin and eosin $\times 3$ green filter). (e) Sham-operated controls; (f) thyroidectomized controls; (g) thyroidectomized + 1 $\mu\text{g. L-triiodothyronine/day}$; (h) thyroidectomized + 20 $\mu\text{g. triiodothyronine/day}$. The sections have been taken from approximately the same level in all cases. The animals receiving the large dose (h) show a relative increase in the size of the kidney cortex and the ventricles of the heart.

Studies on the Determination of the Sequence of Amino-Acids in Peptides and Proteins

1. THE PREPARATION, PROPERTIES AND CHROMATOGRAPHIC ADSORPTION OF THE AZOBENZENE-*p*-SULPHONYL DERIVATIVES OF VARIOUS AMINO-ACIDS

By H. M. FLOWERS AND W. S. REITH
Organic Chemistry Department, The University, Leeds

(Received 19 July 1952)

Almost simultaneously with the conception of the polypeptide structure of proteins, attempts were made to devise methods for the determination of the amino-acids situated at that end of the polypeptide chain which bears the free α -amino group. Fischer & Abderhalden (1907) were the first to introduce such a method of end-group determination, and clearly to state their aims. By reaction of the free amino group with naphthalene-2-sulphonyl chloride and subsequent hydrolysis of the peptide bonds, they obtained the amino-acid bearing the end group in form of the *N*-naphthalene-2-sulphonyl derivative. However, further work showed that the method was inapplicable in certain cases, and the yields were often very poor. Later, Gurin & Clarke (1934) used benzenesulphonyl chloride as a 'marker' in an investigation on the nature of the free amino groups in gelatin.

In these methods the *N*-substituted ('marked') amino-acids were isolated from the hydrolysate by methods depending ultimately on differences in solubility. The difficulties were considerable and losses occurred during the isolation.

Sanger (1945), in a brilliant investigation of the free amino groups of insulin, used 1-fluoro-2:4-dinitrobenzene as 'marker'. The 2:4-dinitrophenyl-amino acids were isolated from the hydrolysate by partition chromatography on silica columns.

We resolved to investigate the possibility of the application of adsorption-chromatographic methods to the problem of the separation of 'marked' amino-acids, and to use as 'markers' other derivatives more appropriate for the particular problems we proposed to investigate.

Karrer, Keller & Szönyi (1943) applied to the resolution of amino-acid mixtures the method first introduced for the separation of sugars by one of us (Reich, 1939) using the coloured *N-p*-phenylazobenzoyl derivatives. As the *N*-acyl derivatives prepared by reaction of amino-acids with carboxylic acid chlorides are very unstable to acid hydrolysis, we prepared the present series of derivatives with sulphonyl chlorides to investigate their value for end-group determinations.

Part of this work has been communicated to the Biochemical Society (Reich, 1948).