2. One method is based on condensation with ethyl acetoacetate in 2N-NaOH to form a product which gives a red colour with Ehrlich's reagent (*p*-dimethylaminobenzaldehyde).

3. The other method is a modification of the Jaffé picrate reaction and depends on the reaction of an alkaline picrate complex of ALA with HCl to give an orange-red colour.

4. The effects of varying different factors in the determination, specificity and application to biological materials are described for both methods.

I wish to thank Professor A. Neuberger, F.R.S., for his valuable advice and interest in this work, and for the hospitality of his laboratory; Dr J. J. Scott for providing many of the compounds tested in these assays; Mr B. L. Wright, Mr F. Grover and Mr C. Hill for technical assistance.

REFERENCES

- Berlin, N. I., Gray, C. H., Neuberger, A. & Scott, J. J. (1954). *Biochem. J.* **58**, xxx.
- Berlin, N. I., Neuberger, A. & Scott, J. J. (1956). *Biochem. J.* **64**, 80.
- Bonsnes, R. W. & Taussky, H. H. (1945). *J. biol. Chem.* **158**, 581.
- Cookson, G. H. & Rimington, C. (1954). *Biochem. J.* **57**, 476.
- Eisner, V., Elvidge, J. A. & Linstead, R. P. (1950). *J. chem. Soc.* p. 2223.
- Elson, L. A. & Morgan, W. T. J. (1933). *Biochem. J.* **27**, 1824.
- Gabriel, S. & Pinkus, G. (1893). *Ber. dtsch. chem. Ges.* **26**, 2197.
- Gibson, K. D., Neuberger, A. & Scott, J. J. (1955). *Biochem. J.* **61**, 618.
- Granick, S. & Vanden Schrieck, H. G. (1955). *Proc. Soc. exp. biol., N.Y.*, **88**, 270.
- Hervey, G. R. (1953). *Nature, Lond.*, **171**, 1125.
- Jaffé, M. (1886). *Hoppe-Seyl. Z.* **10**, 391.
- Knorr, L. (1886). *Liebigs Ann.* **236**, 290.
- Knorr, L. & Lange, H. (1902). *Ber. dtsch. Chem. Ges.* **35**, 2998.
- Moore, S. & Stein, W. H. (1948). *J. biol. Chem.* **176**, 367.
- Neuberger, A. & Scott, J. J. (1953). *Nature, Lond.*, **172**, 1093.
- Neuberger, A. & Scott, J. J. (1954). *J. chem. Soc.* p. 1820.
- Neuberger, A., Scott, J. J. & Gray, C. H. (1954). *Biochem. J.* **58**, xli.
- Neuberger, A., Scott, J. J. & Shuster, L. J. (1956). *Biochem. J.* **64**, 137.
- Owen, J. A., Iggo, B., Scandrett, F. J. & Stewart, C. P. (1954). *Biochem. J.* **58**, 426.
- Schloss, B. (1951). *Analyt. Chem.* **23**, 1321.
- Selah, M. & Berger, A. (1955). *J. Amer. chem. Soc.* **77**, 1893.
- Shemin, D. & Russell, C. S. (1953). *J. Amer. chem. Soc.* **76**, 4873.

The Penetration of Ornithine and Citrulline into Liver Slices

BY J. R. BRONK* AND R. B. FISHER
Department of Biochemistry, University of Oxford

(Received 3 October 1955)

The observations of Gornall & Hunter (1943) show that, at concentrations below 100 mg./100 ml., ornithine is uniformly a more effective catalyst of urea synthesis from ammonia in rat-liver slices than is citrulline. This difference in effectiveness appears to be at variance with what is to be expected of the cycle proposed by Krebs & Henseleit (1932), in which ornithine is transformed through citrulline into arginine, which, in turn, is hydrolysed to yield urea and more ornithine.

The increase in the steady-state rate of urea formation would be expected to be the same following equimolar additions of either ornithine or citrulline to systems catalysing this cycle, and during the approach to steady-state conditions the rate of increase in velocity of urea formation would be expected to be greater when citrulline is added than

when ornithine is added. Therefore, the mean rate of urea formation over any period following the addition of a given amount of citrulline would be expected to be not less than the rate over the same period following the addition of an equimolar amount of ornithine.

It has been suggested by Krebs (1952) that the discrepancy between observation and this expectation is due to differences in accessibility of liver cells to the two amino acids. This suggestion appeared to receive some support from the observation of Christensen, Riggs, Fischer & Palatine (1952) that α -diaminobutyric acid, the next lower homologue of ornithine, is powerfully concentrated by mouse liver.

The present investigation was undertaken to determine directly whether the greater effectiveness of ornithine could be accounted for in terms of differences of ease of penetration of ornithine and citrulline into liver cells.

* Present address: Laboratory of Cellular Physiology, National Heart Institute, Bethesda, Maryland, U.S.A.

METHODS

Materials

Male and female rats of about 250 g. body wt. were used. All animals were fed on stock diets until required.

Liver slices were cut with the apparatus of McIlwain & Buddle (1953) at a nominal thickness of 0.225 mm. Immediately after cutting, the slices were washed with about 250 ml. of warm oxygenated Krebs bicarbonate-lactate medium (Krebs & Henseleit, 1932), the cutting and washing being complete within 10 min.

L-Ornithine hydrochloride and L-citrulline were obtained from Roche Products Ltd.

Incubation procedure

Approx. 100 mg. of slices was incubated at 37° in 10 ml. of the bicarbonate-lactate medium. The slices were kept suspended by a stream of gas mixture ($O_2 + CO_2$; 95:5) introduced at the bottom of the tube. The incubation tubes were fitted with air condensers. A trace of tributyl citrate was used to prevent frothing. Control experiments showed that this had no effect on the ability of liver slices to synthesize urea.

Fifteen minutes after adding ornithine or citrulline the slices were removed from the medium, blotted briefly on Whatman no. 54 filter paper, and placed in weighed bottles containing 1 ml. of 50% (w/v) trichloroacetic acid (TCA). A 5 ml. sample of medium was also pipetted into 0.65 ml. of 50% TCA. The slices were weighed and were then quantitatively transferred to a glass mortar with water, and ground up finely. The resulting suspension was quantitatively transferred to a weighed centrifuge tube with more water, and the solid separated by centrifuging. The whole of the supernatant (7–10 ml.) was taken for analysis. The dry weight of the slices was determined by reweighing the centrifuge tubes after they had been dried at 105° overnight. The trichloroacetic acid sublimes during the heating at 105°. Liver water was taken to be equal to the difference between fresh weight and dry weight of the slices.

Analytical methods

Direct colorimetric estimations. Ornithine was estimated according to the method of Chinard (1952) applied directly to TCA extracts, with a Spekker absorptiometer (Ilford spectrum filter no. 603). Citrulline was estimated by the method of Fearon (1939), as modified by Gornall & Hunter (1941), with a Pulfrich step photometer modified in a manner similar to that described by Stämpfli (1945).

Separation of ornithine and citrulline from interfering substances. Columns (5 cm. $\times$ 0.5 cm.) of Dowex 50 resin (2% cross-linkage, 200–400 mesh) were washed with 10 ml. of N-HCl followed by 10 ml. of water. The TCA extract to be analysed was transferred quantitatively to the column and allowed to run through, followed by 10 ml. of water. The column was washed with 5 ml. of 0.1 N-NH₃ soln. This did not displace ornithine or citrulline from a column of these dimensions, and it was thought that it might displace weakly held contaminant cations. The column was then washed with 10 ml. of water. All eluates to this point were discarded. The column was then washed with 10 ml. of N-NH₃ soln. followed by 10 ml. of water, and the eluates were collected. Tests showed that up to 1 mg. of ornithine or citrulline was quantitatively recovered from the column

in these eluates. Finally, the eluates were evaporated to dryness at low pressure and temperature (not over 40°), and the residue was taken up in 0.5 ml. of water.

Paper electrophoresis. The apparatus used was similar to that described by Markham & Smith (1952). The addition of coils for the circulation of tap water through the CCl₄ bath in which the paper was immersed kept the temperature of the bath constant and made prolonged electrophoresis more satisfactory.

Rat liver contains a major unidentified ninhydrin-reacting component which runs close to citrulline at most pH values and is most readily separated from it at pH 3.2. Since ornithine is easily separated from other amino acids at this pH we have used electrophoresis for 3 hr. at 20 v/cm. in 0.045 M phthalate buffer of pH 3.2 as a routine procedure. Whatman no. 3 paper (18 cm. $\times$ 50 cm.) was used and 100 μ l. spots of concentrated column eluate were applied. Table 1 gives the rates of migration of a number of amino acids in these conditions.

Colour development and measurement on paper. The method used was essentially that of Harris, Mittwoch, Robson & Warren (1954). We are grateful to Professor Warren for supplying the details before publication. The paper was dipped in 0.4% ninhydrin in 95% aqueous acetone (v/v) and dried. The colour was developed by heating at 60° for 30 min. This was most satisfactorily accomplished by wrapping the paper round a 10 l. bottle filled with water thermostatically maintained at 60°. After heating, the paper was dipped into a solution made by diluting 2 ml. of saturated aqueous Cu(NO₃)₂ and 0.4 ml. of 10% (v/v) HNO₃ to 100 ml. with acetone, and allowed to dry. The colours are intensified and made orange-red by this treatment. The coloured spots were cut out and the colour was eluted with methanol (5 ml. or more according to the intensity of the colour). The extinctions of the orange-red methanol solutions were measured at 510 m μ . in a Unicam spectrophotometer.

Calibration curves were prepared for solutions of the amino acids in the Krebs lactate medium and carried through the whole procedure, including addition of TCA, the resin stage, concentration, and separation by electrophoresis. Linear relations were obtained for both amino acids up to concentrations of 100 mg./100 ml. in the original solution. The same linear relations between concentration and extinction were found for ornithine and citrulline whether they were placed directly on paper or taken through

Table 1. *Migration of amino acids during electrophoresis*

Aminoacid	Migration (cm.)
Aspartic acid	5.4
Glutamic acid	7.1
Citrulline	11.4
Glycine	16.5
Histidine	27.9
Arginine	30.0
Lysine	30.9
Ornithine	33.6

Electrophoresis was carried out on a 50 cm. strip of Whatman no. 3 paper at pH 3.2, with a potential gradient of 20 v/cm. The values represent migration towards the cathode, uncorrected for endosmotic flow.

the above complex procedure. However, since the slope of the lines varied somewhat with different ninhydrin solutions, we always included standard solutions of ornithine and citrulline in every set of estimations to correct for this variable.

RESULTS

Measurement of the penetration of ornithine into liver slices

Chinard's method. Replicate sets of liver slices were incubated for 15 min. with 0, 11.8, and 23.6 μ moles of ornithine in the 10 ml. of medium, and ornithine concentrations were measured in the slices and medium by Chinard's (1952) method.

Fig. 1. Relation between liver and medium water concentrations of ornithine after incubation for 15 min., ornithine being estimated by Chinard's method. Slope = 0.704 ± 0.116 ; intercept = 4.84.

Fig. 2. Relation between liver and medium water concentrations of ornithine after incubation for 15 min., paper electrophoresis being used to estimate ornithine. Slope = 0.928 ± 0.028 ; intercept = 1.33.

Concentrations in the medium water and liver water (wet wt. — dry wt.) are compared in Fig. 1.

The linearity of the relation between the concentrations in liver water and in medium water suggests that ornithine enters by simple diffusion, but the large apparent ornithine content of the liver in the absence of ornithine in the medium, together with the known lack of specificity of the method (see Chinard, 1952), makes the interpretation of the data uncertain.

Paper electrophoresis. This more specific method was used in experiments in which the incubation period was again 15 min. and the initial ornithine concentrations were 0, 5.9, 11.8 and 17.7 μ moles in the 10 ml. of medium. Fig. 2 shows the results. These experiments give regular results, pointing unequivocally to a linear relation between concentrations in liver water and in medium water, and the apparent ornithine in the absence of added ornithine is low.

Measurement of the penetration of citrulline into liver slices

Fearon's method. The same general procedure of incubation with amino acid for 15 min. as was used in the ornithine experiments was followed in all the experiments with citrulline. Initial citrulline concentrations of 0, 11.4 and 22.8 μ moles/10 ml. were used. Fig. 3 gives the results of five experiments in which citrulline was estimated by Fearon's (1939) method. They show a linear relation between concentrations in medium and liver water, but the apparent citrulline in the liver water in the absence of added citrulline is rather high.

Fig. 3. Relation between liver and medium water concentrations of citrulline after incubation for 15 min., Fearon's method being used to estimate citrulline. Slope = 1.146 ± 0.092 ; intercept = 9.31.

Paper electrophoresis. The results are given in Fig. 4. There is no doubt that they are best described by a linear relation, and that the apparent citrulline in the absence of added citrulline is much lower than that found with Fearon's method of estimation.

Simultaneous penetration of ornithine and citrulline

We have shown (Bronk & Fisher, 1955) that ornithine and citrulline compete with one another in their catalytic effects on urea synthesis. Although the data for penetration are consistent with the hypothesis that both amino acids enter liver cells

rapidly by simple diffusion, it was thought advisable, in view of the competition in catalysis of urea synthesis, to make sure that they did not compete for a pathway of entry into liver cells. Experiments were therefore made in which liver slices were incubated for 15 min. in medium containing both ornithine and citrulline. The results are presented in Table 2, in which the ratio of concentration in liver water to concentration in medium water is given for these experiments and for the earlier ones in which the amino acids were incubated singly with liver at the same concentrations. It is clear that the presence of the second amino acid does not affect penetration.

Effect of time of incubation on penetration

Comparison of the results obtained with the direct colorimetric methods for the two amino acids with those obtained from paper-electrophoretic separation makes it clear that a large part of the apparent ornithine and citrulline found by the direct colorimetric methods in liver slices incubated in the absence of added amino acid must be some other contaminant material. The estimate of citrulline obtained by Fearon's method in these circumstances is three times that obtained by electrophoretic separation. The estimate of ornithine obtained by Chinard's method is between four and five times that obtained by the electrophoretic method.

However, in liver incubated in the absence of added amino acids the paper-electrophoretic technique still indicates appreciable amounts of apparent ornithine and citrulline, and the apparent concentrations of these two amino acids in liver water are much higher than their concentrations in the medium. This raises the question of whether there may be processes by which ornithine and citrulline can be concentrated in the liver cells. If such processes were occurring it would be expected that, at all concentrations of these amino acids in the medium, their concentration in liver water would rise in time above that in the medium. We have therefore made experiments identical with the 15 min. experiments (Table 2), except that the incubations were continued for 30 min. The data are given in Table 3. It is seen that the internal concentration of amino acid is not increased significantly in any instance above that to be expected in comparable 15 min. experiments. The failure of either amino acid to undergo further apparent accumulation in the liver in the experiments in which amino acid was added cannot be ascribed to lack of material in the medium: the ratio of volume of medium water to liver water in the experiments is about 20:1, so that enough apparent ornithine and citrulline were present to permit large increases in concentration ratio.

Fig. 4. Relation between liver and medium water concentrations of citrulline after incubation for 15 min., paper electrophoresis being used to estimate citrulline. Slope = 1.051 ± 0.087 ; intercept = 3.01.

Table 2. *Absence of competition between ornithine and citrulline for access to liver water*

All analyses were made by paper electrophoresis. Experiments with both amino acids were made with 5.9 μ moles of ornithine and 22.8 μ moles of citrulline added to 10 ml. of medium. Experiments with one amino acid added constitute all data from earlier experiments in which the final concentrations in liver water were comparable with those found in the experiments with two amino acids.

Additions to medium	Concn. in liver water/concn. in medium water	
	Ornithine	Citrulline
One amino acid		1.18
	1.01	1.05
	0.86	1.13
		0.99
Both amino acids	1.45	1.06
	1.05	1.01

Recovery experiments

In order to interpret the penetration data it was necessary to know the completeness of recovery of the amino acids from liver slices. Recovery experiments were made by adding liver slices, incubated in the absence of added amino acid, to TCA in the usual fashion, and then adding a known amount of amino acid and carrying out the appropriate method of estimation. The results of these experiments are given in Table 4.

DISCUSSION

The uniformity with which the results of the penetration experiments can be described in terms of a linear relation means that one can say that every increment of concentration in the medium water produces a proportionate increment of the concentration in liver water. If it were possible to establish that all the apparent ornithine or citrulline in liver water in the absence of added amino acid was not ornithine or citrulline but a contaminant, then one could say that the concentration of the amino acid in liver water was directly proportional to that in the medium, the constant of proportionality being the slope of the regression line. This proviso is not fulfilled. But it is clear, from comparison of the estimates obtained by the Chinard and Fearon methods with those obtained by paper electrophoresis, that most of the apparent ornithine and citrulline found by the direct colorimetric procedures when no amino acid has been added to the medium must be contaminant material. Also (see Table 3) prolonging the period of incubation of the

slices from 15 to 30 min. does not increase the concentration of apparent ornithine or citrulline in the liver water. Thus it is difficult to account for the apparent ornithine and citrulline on any hypothesis of active accumulation, and it seems most probable that the apparent residual material is a contaminant. The data of Table 3 are most simply explained on the assumption that the method of paper electrophoresis, though far more specific than the Chinard and Fearon methods, is still not perfectly specific for ornithine and citrulline, and that little or none of the amino acid estimated in the liver slices incubated in the absence of added ornithine or citrulline is in the form of ornithine or citrulline.

This is not a very important question in relation to the main purpose of this work since, even if active accumulation of the amino acids can occur to some extent, the data of Table 3 indicate that it must be occurring maximally at very low internal amino acid concentration. At the concentrations of ornithine and citrulline which are usually used in the study of urea synthesis the liver cells behave almost exactly as if the sole process governing entry was passive diffusion of the amino acids.

The slopes of the regression lines in Figs. 2-4 should therefore represent ratios of increments in amino acid concentration in liver water to increments in amino acid concentration in medium water, provided that there is nothing in liver which interferes with the estimation of added amino acid. Table 4 compares the regression coefficients (multiplied by 100 so that they give concentration in liver water as percentage of that in the medium) with percentage recoveries. It appears that liver contains something which interferes with Chinard's

Table 3. *Effect of incubation time on the penetration of ornithine and citrulline*

Amino acid	Amount added initially (μ moles)	No. of experiments	Concn. after 30 min. incubation (μ moles/10 ml.)		Concn. expected in liver water* (μ moles/10 ml.)
			Medium water	Liver water	
Citrulline	0	6	0.72 $\pm$ 0.051	4.55 $\pm$ 0.59	3.77 $\pm$ 1.73
	22.8	4	20.50 $\pm$ 0.52	19.14 $\pm$ 0.74	24.55 $\pm$ 1.16
Ornithine	0	6	0.10 $\pm$ 0.017	1.36 $\pm$ 0.06	1.42 $\pm$ 0.24
	5.9	6	4.29 $\pm$ 0.055	5.19 $\pm$ 0.46	5.31 $\pm$ 0.17

* These figures are calculated from the regression equations of Figs. 2 and 4 on the assumption that no further amino acid has entered after 15 min.

Table 4. *Comparison of the coefficients of regression of concentrations in liver water on concentrations in medium with the recoveries of ornithine and citrulline from rat-liver slices*

The numbers in parentheses denote the numbers of observations.

Amino acid	Analytical method	Regression coefficient ($\times 100$)	Percentage recovery
Ornithine	Chinard (1952)	70.4 $\pm$ 11.7 (12)	74.5 $\pm$ 7.6 (4)
	Paper electrophoresis	92.8 $\pm$ 2.8 (7)	96.9 (2)
Citrulline	Fearon (1939)	114.6 $\pm$ 9.3 (15)	94.5 $\pm$ 3.7 (5)
	Paper electrophoresis	105.1 $\pm$ 8.7 (8)	96.8 (2)

method for the estimation of ornithine. There is practically quantitative recovery of added ornithine by the paper-electrophoretic method, so that the low recovery by Chinard's method cannot be ascribed to binding of amino acid by liver solids.

Comparison between the regression coefficients and the percentage recoveries shows that in all instances the two agree within the limits of experimental error. Little or no weight can be placed on the differences which are shown, but it is of interest that the trend is for regression coefficients to be less than recoveries for ornithine and greater than recoveries for citrulline, so that there is a slight suggestion that citrulline penetrates into liver water more rapidly than ornithine.

We may conclude from this work that, at the concentrations of ornithine and citrulline ordinarily used in the study of urea synthesis, the intracellular concentration of both amino acids becomes indistinguishable from the external concentration within 15 min. of addition of the amino acid. In experiments such as those of Gornall & Hunter (1943) involving incubations of 2 hr. there can be no question of differences of cell permeability contributing appreciably to the differences in response to ornithine and citrulline.

It might, however, appear to be possible to invoke a permeability factor in connexion with penetration of the amino acids into intracellular particles, but such an explanation of the lower effectiveness of citrulline has to contend with the demonstration by Cohen & Hayano (1948) that the enzymes converting citrulline into arginine occur in the non-particulate fraction of the cytoplasm. That is, any intracellular permeability barrier would be expected to operate to the disadvantage of ornithine.

SUMMARY

1. Methods are described for concentration and separation by paper electrophoresis of ornithine and citrulline, and for their subsequent determination by the ninhydrin procedure of Harris *et al.* (1954).

2. These methods, and those of Chinard (1952) and Fearon (1939), have been applied to the study of the penetration of ornithine and citrulline into rat-liver slices.

3. Citrulline appears to penetrate a little more readily than ornithine. Both amino acids have almost the same concentration in liver water as in medium water after incubation for 15 min. at 37°.

4. No significant changes in the concentrations of the amino acids in liver water occur when the period of incubation is prolonged to 30 min.

5. The data do not support the view that ornithine is a more effective catalyst of urea synthesis than citrulline because it has easier access to liver cells.

REFERENCES

- Bronk, J. R. & Fisher, R. B. (1955). *Biochem. J.* **60**, xxix.
Chinard, F. P. (1952). *J. biol. Chem.* **199**, 91.
Christensen, H. N., Riggs, T. R., Fischer, H. & Palatine, H. M. (1952). *J. biol. Chem.* **198**, 17.
Cohen, P. P. & Hayano, M. (1948). *J. biol. Chem.* **142**, 405.
Fearon, W. R. (1939). *Biochem. J.* **33**, 902.
Gornall, A. G. & Hunter, A. (1941). *Biochem. J.* **35**, 650.
Gornall, A. G. & Hunter, A. (1943). *J. biol. Chem.* **147**, 593.
Harris, H., Mittwoch, U., Robson, E. R. & Warren, F. L. (1954). *Ann. hum. Genet., Lond.*, **19**, 196.
Krebs, H. A. (1952). *The Enzymes, Chemistry and Mechanism of Action*, vol. 2, part 2, p. 866. Ed. by Sumner, J. B. & Myrback, K. New York: Academic Press Inc.
Krebs, H. A. & Henseleit, K. (1932). *Hoppe-Seyl. Z.* **210**, 33.
McIlwain, H. & Buddle, H. L. (1953). *Biochem. J.* **53**, 412.
Markham, R. & Smith, J. D. (1952). *Biochem. J.* **52**, 552.
Stämpfli, R. (1945). *Helv. physiol. Acta*, **3**, 628.

The Role of Ornithine and Citrulline in Urea Synthesis

By J. R. BRONK* AND R. B. FISHER

Department of Biochemistry, University of Oxford

(Received 3 October 1955)

We have shown (Bronk & Fisher, 1956) that the anomalously greater effectiveness of ornithine, compared with that of citrulline, in catalysing synthesis of urea in rat-liver slices is not due to greater ease of access of ornithine to liver cells, as suggested by Krebs (1952). A possible alternative

explanation is that ornithine and citrulline catalyse separate processes producing urea. This paper describes experiments made to test this possibility.

METHODS

Biological material. Liver slices from rats and guinea pigs (average weights were 250 and 600 g. respectively) were used. The animals were fed on laboratory stock diet until

* Present address: Laboratory of Cellular Physiology, National Heart Institute, Bethesda, Maryland, U.S.A.