REFERENCES

- Barker, S. B. (1955). *Endocrinology*, **57**, 414.
- Bartlett, P. D., Grimmett, P., Beers, L. & Shelata, S. (1956). *J. biol. Chem.* **218**, 419.
- Baruch, H. & Chaikoff, I. L. (1955). *Endocrinology*, **56**, 609.
- Chernick, S. S. & Moe, J. G. (1956). *Endocrinology*, **58**, 280.
- Duve, C. de, Pressman, B. C., Gianetto, R., Wattiaux, R. & Appelmans, F. (1955). *Biochem. J.* **60**, 604.
- Engel, F. L. (1954). *Progr. Allergy*, **4**, 227.
- Gianetto, R. & Duve, C. de (1955). *Biochem. J.* **59**, 433.
- Greenbaum, A. L. (1955). *The Hypophyseal Growth Hormone, Nature and Actions*, p. 330. Ed. by Smith, R. W., Gaebler, O. H. & Long, C. N. H. New York: McGraw-Hill Publ. Co.
- Greenbaum, A. L. & Greenwood, F. C. (1954). *Biochem. J.* **56**, 625.
- Henderson, R. W. & Rawlinson, W. A. (1956). *Biochem. J.* **62**, 21.
- Heppel, L. A., Whitfeld, P. R. & Markham, R. (1955). *Biochem. J.* **60**, 8.
- Hokin, E. & Hokin, M. R. (1954). *J. Histochem. Cytochem.* **2**, 395.
- Kaplan, H. L. & Heppel, L. A. (1956). *J. biol. Chem.* (in the Press).
- Kuff, E. L. & Schneider, W. C. (1954). *J. biol. Chem.* **206**, 677.
- Kuriaki, K. & Kensler, C. J. (1954). *J. Biochem., Tokyo*, **41**, 409.
- Lamirande, G. de, Allard, C., Costa, H. C. da, & Cantero, A. (1954). *Science*, **119**, 351.
- Lowe, C. U. (1955). *J. nat. Cancer Inst.* **15**, 1619.
- MacLeod, L. D. & Reiss, M. (1940). *Biochem. J.* **34**, 820.
- Perry, W. F. & Bowen, H. F. (1956). *Amer. J. Physiol.* **184**, 59.
- Phillips, A. H. & Langdon, R. G. (1956). *Biochim. biophys. Acta*, **19**, 380.
- Pressman, B. C. & Lardy, H. A. (1952). *J. biol. Chem.* **197**, 547.
- Reid, E. (1955). *Nature, Lond.*, **175**, 461.
- Reid, E. (1956). *J. Endocrin.* **13**, 319.
- Reid, E., O'Neal, M. A. & Lewin, I. (1956). *Biochem. J.* **64**, 730.
- Reid, E., O'Neal, M. A., Stevens, B. M. & Burnop, V. C. E. (1956). *Biochem. J.* **64**, 33.
- Ringler, I. & Leonard, S. L. (1954). *Endocrinology*, **55**, 363.
- Roth, J. S. (1954a). *J. biol. Chem.* **208**, 181.
- Roth, J. S. (1954b). *Nature, Lond.*, **174**, 129.
- Roth, J. S. (1956). *Fed. Proc.* **15**, 341.
- Schneider, W. C. & Hogeboom, G. H. (1950). *J. nat. Cancer Inst.* **10**, 969.
- Schneider, W. C. & Hogeboom, G. H. (1952). *J. biol. Chem.* **198**, 155.
- Schneider, W. C. & Potter, V. R. (1943). *J. biol. Chem.* **149**, 217.
- Tipton, S. R. (1944). *Endocrinology*, **34**, 181.
- Tipton, S. R., Leath, N. J., Tipton, I. H. & Nixon, W. L. (1946). *Amer. J. Physiol.* **145**, 693.
- Victor, J. & Andersen, D. H. (1938). *Amer. J. Physiol.* **122**, 296.
- Weiss, P. (1955). *The Hypophyseal Growth Hormone, Nature and Actions*, p. 265. Ed. by Smith, R. W., Gaebler O. H. & Long, C. N. H. New York: McGraw-Hill Publ. Co.
- Wollman, S. H. & Scow, R. O. (1951). *Endocrinology*, **49**, 105.

Studies on Ali-esterases

4. DIFFERENTIATION OF ESTERASES IN PANCREAS AND BRAIN BY THE USE OF ORGANOPHOSPHORUS INHIBITORS*

By D. K. MYERS†

Pharmaco-therapeutic Laboratory, University of Amsterdam, Amsterdam, Holland

(Received 15 March 1956)

The results obtained in previous investigations have demonstrated the value of the organophosphorus compounds as experimental tools for the differentiation of the various types of enzymes with esterase activity. By means of these inhibitors it is possible to distinguish clearly between certain of the esterases which are present in the same tissue, e.g. between the A-esterases, ali-esterases (B-type esterase) and a resistant tributyrinase which may all be present in plasma (Aldridge, 1953, 1954; Mounter & Whittaker, 1953; Myers, Schotte & Mendel, 1955a).

* Part 3: Myers, Schotte, Boer & Borsje-Bakker (1955b).

† Present address: Suffield Experimental Station, Ralston, Alberta, Canada.

Later it was also shown that at least three types of enzymes with esterase activity can be distinguished in pancreas by the use of these organophosphorus inhibitors (Myers, Schotte, Boer & Borsje-Bakker, 1955b). The lipase, which is responsible for the hydrolysis of tributyrin by pancreas suspension, is relatively resistant to inhibition by most of the organophosphorus derivatives except diethyl *p*-nitrophenyl phosphate (E 600) (cf. Aldridge, 1954); while the cholesteroesterase, which is probably responsible for the hydrolysis of methyl and phenyl butyrates by pancreas (cf. Fodor, 1950), is readily inhibited by all the organophosphorus compounds tested. The slow hydrolysis of salicyl butyrate by pancreas is due largely to a third enzyme (cf.

Hofstee, 1952) which appeared to be resistant to inhibition by E 600 as well as by the other organophosphorus compounds. Further experiments on the differentiation of the enzymes and their specificities will be presented below.

These investigations with organophosphorus compounds have been extended to brain in an effort to define more clearly the esterases present in this tissue, in the same way as reported previously for the esterases of blood and pancreas (Aldridge, 1953, 1954; Mounter & Whittaker, 1953; Myers *et al.* 1955*a, b*). The results obtained indicate that brain contains at least three esterases capable of hydrolysing fatty acid esters such as tributyrin, phenyl butyrate or ethyl butyrate. These esterases could be clearly distinguished from each other and from the other esterases present in brain, i.e. the cholinesterases and a proteolytic enzyme capable of hydrolysing amino acid esters.

METHODS

Esterase activity was measured manometrically by the Warburg method at 37.5° and pH 7.4 in a medium containing 0.025 M-NaHCO₃ and saturated with CO₂ + N₂ (5:95, by vol.). Aqueous homogenates of rat brain and rat pancreas were prepared with a glass Potter-Elvehjem homogenizer. Unless otherwise indicated in the text, the reaction mixture containing tissue suspension was incubated for 30 min. at 37.5° in the Warburg vessels with or without inhibitor before a small volume of substrate was added to determine the residual esterase activity.

For the experiments with pancreas, which are reported in Table 1, a concentrated suspension containing 10 mg. of fresh pancreas/ml. was incubated with or without inhibitor in bicarbonate buffer for 2 hr. at 37.5° and pH 7.4; subsequently a suitable amount of this suspension was pipetted into the Warburg vessels and the esterase activity determined in the usual way after a further equilibration period of 10 min. at 37.5°. A portion (5 ml.) of the concentrated suspension could be pipetted directly into the Warburg vessels when the ethyl esters of tyrosine or leucine were used as substrates, since these esters gave a low activity in the control vessels without inhibitor. However, the activity towards tributyrin was very high, and consequently it was necessary to dilute the pancreas suspension 1000-fold for the experiments with tributyrin as substrate; for this purpose, the concentrated suspension was first diluted 100-fold in bicarbonate buffer and 0.5 ml. of this diluted suspension was pipetted into the Warburg vessels which contained a total final volume of 5.0 ml. Intermediate dilutions were employed with the esters which were hydrolysed at an intermediate rate, the dilution being adjusted so that the control vessels with each substrate showed an activity of 15–35 μ l. of CO₂/5 min. These dilutions could produce a return of the enzyme activity in the vessels with inhibitor if the inhibition was rapidly reversible; however, evidence was presented in the previous paper (Myers *et al.* 1955*b*) to show that the lipase and esterase of rat pancreas are inhibited irreversibly by organophosphorus compounds such as E 600. Moreover, the enzyme activity was always measured at 5 min. intervals for 30–60 min. and no gradual

return of the activity in the vessels containing inhibited enzyme was observed during this period.

The substrates were prepared as an aqueous emulsion or solution containing 5% (w/v) gum acacia and adjusted to pH 7.4. Esters such as tributyrin were emulsified in the gum-acacia solution with the aid of a Potter-Elvehjem homogenizer. The resultant emulsion is finely divided and moderately stable; when stored at 0°, the tributyrin does tend to settle out but is readily resuspended on shaking the emulsion.

In most cases, the final concentrations of ester and gum acacia in the reaction mixture were 0.2 and 0.1% (w/v) respectively; the final concentration of salicyl butyrate was increased to 1% (w/v) in order to increase the rate of enzymic hydrolysis by the pancreas suspensions. In all experiments the esterase activity was calculated from the initial rate of hydrolysis of the ester after the appropriate corrections had been made for the slow non-enzymic hydrolysis of this substrate.

In addition to the esters described previously (Myers *et al.* 1955*b*), ethyl butyrate, ethyl β -phenylpropionate (Erlenmeyer, 1866) and the ethyl ester of β -phenylalanine (Curtius & Muller, 1904) were prepared by the methods described in the literature. The esterase inhibitors used were E 600, diisopropyl *p*-nitrophenyl phosphate (DINP), tetraethyl pyrophosphate (TEPP) and eserine. E 600 and DINP were kindly supplied by Albright and Wilson, Ltd., London, and TEPP by Dr H. R. Gersmann, Laboratory for General and Inorganic Chemistry, University of Amsterdam.

RESULTS

Pancreas

Table 1 shows the effect of various concentrations of E 600 and DINP on the activity of fresh pancreas homogenates towards a series of esters. The degree of inhibition of the esterases by a given concentration of E 600 or DINP is dependent on the concentration of pancreas in the reaction mixture (Myers *et al.* 1955*b*). In these experiments, therefore, a concentrated enzyme preparation was incubated with the inhibitors in order to produce an irreversible inhibition of the various esterases under identical experimental conditions (cf. Myers *et al.* 1955*b*); subsequently a suitable amount of the suspension was pipetted into the Warburg vessels and the residual esterase activities were measured.

The results obtained for the hydrolysis of tributyrin by the pancreatic lipase and for the hydrolysis of phenyl butyrate by the pancreatic esterase have been described previously (Myers *et al.* 1955*b*) but are included in Table 1 for comparative purposes. It can be seen that the lipase is scarcely affected by DINP but is almost completely inhibited by 5×10^{-4} M E 600 under the experimental conditions (cf. Aldridge, 1954; Myers *et al.* 1955*b*); on the basis of these criteria, it appears that the lipase is responsible for 1% of the total activity towards phenyl butyrate and for 17% of the total activity towards ethyl butyrate, as well as for all the activity of the pancreas homogenate towards

tributyrin (Table 1). The esterase responsible for the remaining 99 % of the total activity towards phenyl butyrate and the remaining 83 % of the total activity towards ethyl butyrate is completely inhibited by 5×10^{-5} M DINP (Table 1). On the basis of sensitivity to inhibition by DINP, it appears that the activity of the pancreas homogenates towards ethyl β -phenylpropionate and the amino acid esters is due mainly to the same esterase which hydrolyses phenyl butyrate.

A slight activity towards the amino acid esters remains which is not inhibited by high concentrations of E 600 or DINP under the experimental conditions (Table 1) and which is therefore probably due to a third enzyme. The enzyme responsible for the E 600-resistant activity towards the amino acid esters could not be distinguished from the enzyme responsible for the E 600-resistant activity towards salicyl butyrate by the type of experiment shown in Table 1 (cf. Myers *et al.* 1955*b*). However, Table 2 shows that these enzymes must be different. The enzyme responsible for the E 600-resistant activity towards salicyl butyrate is strongly inhibited by cysteine and activated by MgCl_2 and NaCl ; in this respect it appears to resemble carboxypeptidase (cf. Smith, 1951*a, b*). None of the other enzymes studied exhibits a similar pattern of results

(Table 2). It might be noted that carboxypeptidase, in contrast to the other proteolytic enzymes in pancreas, does not appear to be sensitive to inhibition by the organophosphorus compounds (Balls & Jansen, 1952). Since this enzyme hydrolyses α -carboxyphenethyl chloroacetate and other esters which possess a free carboxyl group in the appropriate position on the alcohol residue of the ester, it is conceivable that carboxypeptidase might also hydrolyse salicyl butyrate. The inhibitory effects of β -phenylpropionic acid and phenylacetic acid on the hydrolysis of salicyl butyrate by the pancreas homogenates (Table 2) offer further evidence in support of this suggestion (cf. Smith, 1951*a, b*).

The enzyme responsible for the E 600-resistant activity of rat pancreas towards ethyltyrosine is probably chymotrypsin. Both trypsin and chymotrypsin are capable of hydrolysing amino acid esters such as ethyltyrosine and ethylphenylalanine (Neurath & Schwert, 1950; Neurath, 1952), and both enzymes are relatively resistant to inhibition by E 600 (Hartley & Kilby, 1952; Kilby & Youatt, 1954). However, trypsin is not stable under the experimental conditions used in this investigation (cf. Kunitz, Northrop & Herriot, 1948) and does not hydrolyse these esters rapidly, in contrast to chymotrypsin. Although the E 600-resistant

Table 1. *Effect of DINP and E 600 on the esterase activities of rat pancreas*

Homogenates containing 10 mg. of fresh rat pancreas/ml. were incubated at pH 7.4 and 37.5° for 2 hr. with various concentrations of DINP or E 600. The esterase activities were subsequently measured after suitable dilution of the pancreas suspensions as described in the text.

Inhibitor	Concn. of inhibitor (M)	Esterase activity ($\mu\text{l. of CO}_2/\text{mg. of pancreas/hr.}$)						
		Tributyrin	Phenyl butyrate	Ethyl butyrate	Ethyl β -phenylpropionate	Ethyl- β -phenyl-alanine	Ethyl-tyrosine	Ethyl-leucine
None	—	6400	1200	94	210	19.0	5.1	8.0
DINP	5×10^{-8}	6400	1025	84	180	16.8	4.6	7.2
	5×10^{-7}	6400	310	36	54	4.5	2.2	2.6
	5×10^{-6}	6400	30	18	4	1.7	1.5	0.8
	5×10^{-5}	6200	12	16	3	1.5	1.5	0.8
	5×10^{-4}	5100	10	14	2	1.5	1.5	0.8
E 600	5×10^{-5}	1560	3	4	1.4	—	1.5	0.8
	5×10^{-4}	180	0.2	0.2	1.2	—	1.2	0.8

Table 2. *Effect of several inhibitors and activators on the esterases of rat pancreas*

The activities towards ethyltyrosine and salicyl butyrate were measured in the presence of 5×10^{-4} M E 600 in order to inhibit the hydrolysis of these esters by E 600-sensitive enzymes (see Table 1 and Myers *et al.* 1955*b*).

Compound added to the reaction mixture	Esterase activity as percentage of the normal activity in 0.025 M bicarbonate buffer			
	Salicyl butyrate (+ E 600)	Ethyl-tyrosine (+ E 600)	Phenyl butyrate	Tributyrin
MgCl_2 (0.1 M)	230	70	30	74
NaCl (0.5 M)	180	75	66	76
Cysteine (0.01 M)	10	45	90	87
β -Phenylacetic acid (0.001 M)	79	—	—	—
β -Phenylpropionic acid (0.001 M)	26	—	—	—

activity of pancreas towards ethyltyrosine is not significantly inhibited by E 600 under the conditions used in Table 1, it could be inhibited by very high concentrations of this compound. The activity of this enzyme was, in fact, 80 % inhibited after incubation with 5×10^{-3} M E 600 for 2 hr. at 37.5° ; this value is of the same order of magnitude as that found by Hartley & Kilby (1952) for the inhibition of purified chymotrypsin by E 600.

Brain

In the previous investigations on selective esterase inhibitors (cf. Mendel, Myers, Uyldert, Ruys & Bruyn, 1953; Myers, 1954), it was assumed for practical purposes that the eserine-resistant activity of rat brain towards tributyrin is due to a single enzyme. This assumption is not strictly correct, although the major portion of this activity is due to a single esterase. The esterases of brain were studied in more detail by Aldridge (1954), who obtained evidence to show that brain must contain more than one ali-esterase, although his data do not define the substrate specificity of these enzymes clearly. These ali-esterases cannot be clearly distinguished when E 600, TEPP or *NN*-dimethylcarbamoyl fluoride is employed to inhibit their activities (Myers, 1954; cf. Myers, 1956); apparently the various ali-esterases of brain do not differ markedly in their sensitivity to these inhibitors. However, the results obtained with DINP clearly show that the hydrolysis of tributyrin by rat brain must be due to two esterases, both of which are resistant to inhibition by 2×10^{-6} M eserine and therefore distinct from the cholinesterases (Table 3). Similar results were obtained with tributyrin when certain other inhibitors were used in place of DINP, e.g. *p*-nitrophenyl *NN'*-dimethylphosphorodiamidate, and bis-*p*-nitrophenyl methyl- or phenyl-

phosphoroamidate (Myers, 1954); further investigations have been carried out only with DINP.

The inhibition of the activity of brain homogenates towards tributyrin or triacetin increases steadily as the concentration of DINP is increased, until a steady value is attained when the concentration of DINP reaches 5×10^{-6} M (Table 3). A further increase in the concentration of DINP up to 5×10^{-5} M has little or no effect on the degree of inhibition, while an increase to 5×10^{-4} M DINP causes only a small depression of the residual esterase activity towards these substrates. Apparently the brain contains one esterase which is completely inhibited by 5×10^{-6} M DINP and a second esterase which is resistant to inhibition by 5×10^{-6} or 5×10^{-5} M DINP and only partially inhibited by 5×10^{-4} M DINP. This conclusion is supported by the finding that the low activity of the brain towards ethyl butyrate is completely inhibited by 5×10^{-6} M DINP (Table 3); the hydrolysis of this ester can be ascribed entirely to the DINP-sensitive ali-esterase.

A comparison of the data obtained with tributyrin and triacetin as substrates (Table 3) suggested that a third esterase might be present. The DINP-resistant activity towards tributyrin is slightly more sensitive to inhibition by high concentrations of DINP than is the DINP-resistant activity towards triacetin. It was also observed that the DINP-resistant activity towards a series of esters did not show a regular pattern; the rate of hydrolysis of phenyl esters decreased regularly in the order acetate, propionate, butyrate, but the rate of hydrolysis of the triglycerides decreased in the order butyrate, acetate, propionate.

In order to investigate the possibility that the DINP-resistant esterase activity might be due to more than one enzyme, the sensitivity of the

Table 3. *Effect of DINP and E 600 on the esterase activities of rat brain*

Homogenates containing 20 mg. of fresh rat brain/ml. were incubated at pH 7.4 and 37.5° for 30 min. with various concentrations of DINP or E 600 in the Warburg vessels. The esterase activities were subsequently measured without further dilution of the enzyme preparation. All measurements were carried out in the presence of 2×10^{-6} M eserine in order to inhibit selectively any cholinesterase activity.

Inhibitor	Concn. of inhibitor (M)	Esterase activity (μ l. of CO_2 /100 mg. of brain/hr.)			
		Triacetin	Tributyrin	Ethyl butyrate	Ethylleucine
None	—	210	548	40	38
DINP	5×10^{-9}	202	495	34	—
	5×10^{-8}	166	348	20	—
	5×10^{-7}	127	212	5	38
	5×10^{-6}	114	158	0	38
	5×10^{-5}	114	156	0	38
	5×10^{-4}	100	88	0	—
E 600	5×10^{-8}	—	334	—	—
	5×10^{-7}	—	192	—	—
	5×10^{-6}	108	154	0	38
	5×10^{-5}	60	102	0	38
	5×10^{-4}	15	30	0	36

esterase activities to destruction at an alkaline pH has been studied. The results of a previous investigation had demonstrated the existence of an eserine-resistant triacetinase in rat brain which, unlike the other esterases, was not inactivated at pH 10.0–10.5 (Myers, 1949). A homogenate containing 200 mg. of brain (wet weight)/ml. was, therefore, exposed to pH 10.5 for 30 min. at room temperature and subsequently neutralized. The DINP-sensitive ali-esterase was almost completely inactivated, while the cholinesterase activity towards acetyl- β -methylcholine was partially destroyed. The DINP-resistant activity towards tributyrin was reduced from 156 to 20 μ l. of CO₂/100 mg. of brain/hr., while the DINP-resistant activity towards triacetin was only slightly affected, decreasing from 114 to 105 μ l. of CO₂/100 mg. of brain/hr. Apparently it is possible to distinguish between a DINP-resistant tributyrinase and a DINP-resistant triacetinase in brain by this method. The DINP-resistant tributyrinase, like the DINP-sensitive enzyme, appears to be inactivated by exposure to alkali, while the DINP-resistant triacetinase retains the major portion of its activity under these conditions. The small

activity towards tributyrin which remains after exposure to pH 10.5 cannot be further reduced by a more drastic treatment without inactivating the enzyme responsible for the hydrolysis of triacetin (cf. Myers, 1949); it seems probable, therefore, that the residual activities towards tributyrin and triacetin are due to the same enzyme, i.e. the DINP-resistant triacetinase also exhibits a small activity towards tributyrin.

The differences between the brain esterases in their sensitivity to various inhibitors and to destruction by alkali are summarized in Table 4. On the basis of these differences, it is possible to distinguish between the activities of each of the esterases of rat brain and consequently to study the substrate specificities of each of these enzymes. The results obtained in this way are given in Table 5.

It can be seen that each of the esterase activities which have been separated exhibit a characteristic substrate-specificity pattern; moreover, the substrate-specificity patterns do not exhibit any marked irregularities such as were observed with the total DINP-resistant esterase activity (Table 5). These facts support the conclusion that each of the

Table 4. *Characteristics of the esterases of rat brain*

Enzyme	Inhibition by 2×10^{-6} M eserine	Inhibition by 5×10^{-6} M DINP	Destruction at pH 10.5 for 30 min.	Inhibition by 5×10^{-4} M E 600
Cholinesterase	+	+	(+)	+
DINP-sensitive ali-esterase	—	+	+	+
DINP-resistant tributyrinase	—	—	+	+
DINP-resistant triacetinase	—	—	—	+

Table 5. *Rate of hydrolysis of various esters by the esterases of rat brain*

Esterase activities were measured at pH 7.4 and 37.5° as described in the text. The total activity of a homogenate of rat brain towards various esters is given in the second col.; this activity was divided into its component parts on the basis of the characteristics described in Table 4. The portion of the activity which was inhibited by 2×10^{-6} M eserine or 10^{-7} M TEPP was ascribed to the cholinesterases and is given in the third col. The values given for the DINP-sensitive ali-esterase represent that portion of the activity which was inhibited by 5×10^{-6} M DINP but not by 2×10^{-6} M eserine. The activity of the DINP-resistant triacetinase was measured in the presence of 5×10^{-6} M DINP after the brain homogenate had been allowed to stand at pH 10.5 for 30 min. at room temperature. The activity of the DINP-resistant tributyrinase was estimated from the difference between the activity of a fresh brain homogenate in the presence of 5×10^{-6} M DINP and the residual activity of the DINP-resistant triacetinase which remained after the homogenate had been treated with alkali as described above in order to inactivate the tributyrinase. In the cases where no value is given, the activity of the esterases in question was too low to be measured accurately.

Substrate	Esterase activity (μ l. of CO ₂ /mg. of brain/hr.)				
	Total activity	Cholin- esterases	DINP- sensitive ali-esterase	DINP- resistant triacetinase	DINP- resistant tributyrinase
Tributyrin	5.5	—	3.9	0.2	1.4
Tripropionin	8.1	0.6	6.5	0.6	0.4
Triacetin	3.3	1.3	0.9	1.0	—
Phenyl butyrate	11.6	1.3	8.8	1.3	—
Phenyl propionate	21.0	7.4	10.8	2.2	—
Phenyl acetate	22.6	8.4	7.7	5.9	—
Benzyl acetate	15.3	6.6	7.5	1.0	—
Ethyl butyrate	0.38	—	0.38	—	—
Acetyl- β -methylcholine	3.25	3.25	—	—	—

esterase fractions distinguished by the methods given in Table 4 does, in fact, represent a distinct esterase. The results do not prove, of course, that all of the esterases in the brain homogenate have been identified; on the other hand, there is no clear evidence which would suggest the presence of other esterases capable of hydrolysing the substrates in question at an appreciable rate.

The data given in Table 5 show that both the DINP-sensitive esterase and the DINP-resistant triacetinase are capable of hydrolysing phenyl esters as well as triglycerides; in this respect these enzymes differ from the esterases in pancreas described previously (Myers *et al.* 1955*b*). Both of these enzymes appear to hydrolyse phenyl esters more efficiently than triglycerides and to act most efficiently on the esters of short-chain fatty acids. The rate of hydrolysis by the DINP-sensitive esterase decreases in the order propionate, butyrate, acetate, whereas the rate of hydrolysis by the DINP-resistant triacetinase decreases in the order acetate, propionate, butyrate (Table 5).

Owing to the experimental methods used to distinguish the DINP-resistant tributyrinase (cf. Table 4), reliable values for the activity of this enzyme could only be obtained with tributyrin and tripropionin as substrates (Table 5). However, the DINP-resistant tributyrinase does not hydrolyse phenyl esters rapidly and is the only one of the esterases in rat brain which appears to act preferentially on tributyrin or possibly on the triglycerides of higher fatty acids. As judged from the data given by Aldridge (1954) and in Table 5, it is highly probable that this enzyme hydrolyses tricaproin even more rapidly than tributyrin. In many respects this enzyme appears to resemble the lipase of pancreas (cf. Myers *et al.* 1955*b*), although the activity of pancreatic lipase towards tributyrin is about 5000 times greater per mg. of fresh tissue (Tables 1 and 5).

Data on the activities of the cholinesterases present in the rat-brain homogenates are also included in Table 5. These data are of minor interest since no attempt was made to distinguish between true cholinesterase and pseudocholinesterase, and since the specificities of the cholinesterases have been studied extensively by other investigators (cf. Whittaker, 1951). However, the results given in Table 5 show that the cholinesterases may be responsible for a considerable portion of the total activity of rat brain towards some of the non-choline esters used as substrates. In the present investigation, all of the experiments on the aliesterases of rat brain were carried out in the presence of 2×10^{-6} M eserine in order to inhibit the cholinesterases completely and selectively (cf. Easson & Stedman, 1937; Richter & Croft, 1942; Mendel & Rudney, 1943; Myers & Mendel, 1949).

As an additional check on this method, some experiments were also carried out in the presence of 10^{-7} M TEPP; this compound inhibits the true cholinesterase of rat brain almost completely at a concentration of 10^{-6} M under the experimental conditions but does not have an appreciable effect on the esterase activity towards tributyrin until the concentration approaches 10^{-6} M (Myers, 1954). The results obtained with 2×10^{-6} M eserine and 10^{-7} M TEPP proved to be identical.

In extension of the investigations on the esterases of pancreas, we also studied the hydrolysis of the ethyl esters of leucine and β -phenylalanine by rat-brain homogenates. The ethyl ester of phenylalanine was not hydrolysed at an appreciable rate; in confirmation of the data given by Krinsky & Racker (1949), a low activity towards the ethyl ester of leucine could be demonstrated (Table 3). This activity is due to a proteolytic enzyme which hydrolyses various amino acid amides and esters (Krinsky & Racker, 1949) as well as peptides (Krinsky & Racker, 1952; cf. Zeller, 1945; Racker & Krinsky, 1948). However, this enzyme is scarcely affected by either E 600 or DINP under the experimental conditions (Table 3). The cathepsin responsible for the hydrolysis of ethylleucine by brain suspensions is therefore not identical with any of the other esterases described above.

DISCUSSION

The organophosphorus compounds have proved to be very useful for differentiating the complicated mixture of esterases present in pancreas and brain as well as those present in other tissues, e.g. blood. This task would be considerably more difficult and more laborious by conventional methods of purification of the enzymes concerned; the purification would be especially difficult for those esterases in the brain which seem to be attached to insoluble lipid material (cf. Myers, 1949; Ord & Thompson, 1951). These selective inhibitors make it possible to divide the total activity of a crude tissue preparation into its component parts, each of which can be assigned to a different enzyme.

At the same time, the results indicate some of the limitations of this method of distinguishing between various enzymes with esterase activity. Since pancreas contains more than one enzyme which is relatively resistant to inhibition by E 600, it was incorrectly suggested that the E 600-resistant activities towards salicyl butyrate and ethyltyrosine might be due to a single enzyme (Myers *et al.* 1955*b*); other methods of analysis showed that this suggestion was not justified (Table 2). Similarly, the two DINP-resistant esterases in brain cannot be clearly distinguished by this method, although they do differ in their sensitivity to destruction in alkaline solutions (Table 4).

It may also be difficult to distinguish between various esterases when a single, reactive inhibitor such as E 600 is used. The activities of the ali-esterase and A-esterase in plasma (Aldridge, 1953) and the activities of the cholesteroesterase and lipase in pancreas (Myers *et al.* 1955*b*; cf. Aldridge, 1954) can be separated by this means. However, a less reactive inhibitor such as DINP is necessary in order to distinguish certain of the other esterases present in normal and post-heparin plasma (Myers *et al.* 1955*a*) and to distinguish between the eserine-resistant esterases in brain (Table 3); in both instances, the DINP-sensitive and DINP-resistant esterases do not differ greatly in their sensitivity to E 600. Evidently the best results will be obtained by the use of both types of inhibitor in turn, since some of the enzymes which are resistant to DINP exhibit different sensitivities to inhibition by E 600. In fact, it seems probable that a wider variety of inhibitors would increase the range of applicability of this method for the differentiation of enzymes with esterase activity.

The organophosphorus compounds used in the present investigation represent an unusual type of inhibitor, since they react with the esterases in the same way as do the normal substrates and yet produce an inhibition which is only slowly reversible—in some cases practically irreversible. Nevertheless, it seems probable that the same general principles would be involved as with selective inhibitors of other enzymes. As long as the selectivity of the inhibitor is open to question, the data obtained for the substrate-specificity pattern of a crude enzyme by the use of a selective inhibitor will be more equivocal than those obtained with the purified enzyme. However, these inhibitors seem almost indispensable for investigations on the number of enzymes present in crude tissue preparations and for the preliminary identification of the reactions catalysed by these enzymes.

The results obtained in this investigation emphasize the fact that a single tissue may contain a complex mixture of enzymes with esterase activity towards the same or similar substrates. Rat brain, for example, contains at least five enzymes capable of hydrolysing triacetin or tributyrin or both, namely the true cholinesterase, the pseudocholinesterase (cf. Whittaker, 1951) and three other enzymes (see Tables 4 and 5) with the characteristics of ali-esterases as defined by Richter & Croft (1942; cf. Mounter & Whittaker, 1953). Further it appears to contain a cathepsin capable of hydrolysing ethylleucine. The DINP-resistant tributyrinase seems to have the characteristics of a lipolytic enzyme, as noted above; physiological substrates could, therefore, be assigned to this enzyme as well as to the true cholinesterase and the cathepsin, even

though further investigations are required in order to define the exact physiological functions of the DINP-resistant tributyrinase and the cathepsin. However, we still have little or no information concerning the physiological substrates of the other esterases.

Except in cases where the rate of hydrolysis of the substrate is limited by the rate of decomposition of the intermediary acyl-enzyme derivative (cf. Wilson, 1952), the enzymes with esterase activity generally hydrolyse a variety of substrates at a rate which parallels the inherent lability of the susceptible bond, i.e. thio esters > esters > amides > peptides (Neurath & Schwert, 1950; Smith, 1951*b*; Suzuoki-Ziro & Suzuoki-Tuneko, 1954; Johnston, 1955). It is possible, therefore, that the physiological substrates of the esterases mentioned above might not be esters but belong, rather, to another class of physiological compound, e.g. amides and peptides of the amino acids, or thio esters such as the acyl derivatives of coenzyme A. The esterases in question can be clearly distinguished from the cathepsin in brain described by Krinsky & Racker (1949); however, this does not completely exclude the possibility that at least one of these esterases might have a function in protein metabolism, since the results described by Zeller (1945) indicate that the brain may contain more than one proteolytic enzyme.

Each of the enzymes studied in rat pancreas could be identified with some degree of certainty, i.e. the lipase, the cholesteroesterase, chymotrypsin and carboxypeptidase or a closely related enzyme. The identification of the esterase responsible for the hydrolysis of phenyl, methyl and ethyl butyrates by pancreas homogenates is complicated by the finding that this enzyme exhibits a relatively low activity towards the ethyl esters of amino acids (Table 1). However, the only proteolytic enzymes in pancreas reported to hydrolyse these esters are trypsin and chymotrypsin (Neurath & Schwert, 1950) and both are approximately 100 000 times less sensitive than the pancreatic esterase to inhibition by E 600 (cf. Hartley & Kilby, 1952; Kilby & Youatt, 1954; Myers *et al.* 1955*b*). It seems fairly certain, therefore, that this esterase is not a proteolytic enzyme; moreover, its characteristics are similar to those of the enzyme in pancreas which is responsible for the hydrolysis of cholesterol esters (Fodor, 1950; Myers *et al.* 1955*b*). This finding suggests that the pancreatic esterase is identical with the cholesteroesterase of pancreas and that its low activity towards amino acid esters is due simply to a lack of absolute substrate specificity; the rate of hydrolysis of ethyl β -phenylalanine, it might be noted, is much lower than the rate of hydrolysis of ethyl β -phenylpropionate, the analogous ester without an amino group (Table 1).

SUMMARY

1. The substrate specificities of the esterases in rat pancreas and brain have been studied with organophosphorus compounds as selective esterase inhibitors.

2. The pancreas contains at least two enzymes with esterase activity which are relatively resistant to inhibition by diethyl *p*-nitrophenyl phosphate (E 600). The data obtained suggest that the E 600-resistant activity towards salicyl butyrate is due to carboxypeptidase or an enzyme with similar properties, whereas the E 600-resistant activity towards ethyltyrosine is probably due to chymotrypsin.

3. The enzyme in pancreas identified previously as cholesterol esterase (Fodor, 1950; Myers *et al.* 1955*b*) appears to be capable of hydrolysing a variety of esters including the ethyl esters of tyrosine, phenylalanine and leucine. The low activity towards these amino acid esters probably reflects the broad substrate specificity of this enzyme.

4. Three ali-esterases with distinct substrate-specificity patterns could be distinguished in brain homogenates. These enzymes are distinct from the cholinesterases and from the cathepsin in brain, described by Krimsky & Racker (1949).

5. The possible physiological functions of these enzymes and the differentiation of esterases by means of selective inhibitors are discussed.

The author is much indebted to Miss J. W. Tol and Miss M. H. T. de Jonge for invaluable technical assistance, and to Professor E. C. Slater for his suggestions concerning the preparation of this manuscript.

REFERENCES

- Aldridge, W. N. (1953). *Biochem. J.* **53**, 110.
 Aldridge, W. N. (1954). *Biochem. J.* **57**, 692.
 Balls, A. K. & Jansen, E. F. (1952). *Advanc. Enzymol.* **13**, 321.
 Curtius, T. & Muller, E. (1904). *Ber. deutsch. chem. Ges.* **37**, 1261.
 Easson, E. H. & Stedman, E. (1937). *Biochem. J.* **31**, 1723.
 Erlenmeyer, E. (1866). *Liebigs Ann.* **47**, 250.
 Fodor, P. J. (1950). *Arch. Biochem.* **26**, 331.
 Hartley, B. S. & Kilby, B. A. (1952). *Biochem. J.* **50**, 672.
 Hofstee, B. H. J. (1952). *J. biol. Chem.* **199**, 365.
 Johnston, R. B. (1955). *Fed. Proc.* **14**, 232.
 Kilby, B. A. & Youatt, G. (1954). *Biochem. J.* **57**, 303.
 Krimsky, I. & Racker, E. (1949). *J. biol. Chem.* **179**, 903.
 Krimsky, I. & Racker, E. (1952). *J. biol. Chem.* **198**, 721.
 Kunitz, M., Northrop, J. H. & Herriot, R. M. (1948). *Crystalline Enzymes*. New York: Columbia University Press.
 Mendel, B., Myers, D. K., Uyldert, I. E., Ruys, A. C. & Bruyn, W. M. de (1953). *Brit. J. Pharmacol.* **8**, 217.
 Mendel, B. & Rudney, H. (1943). *Biochem. J.* **37**, 59.
 Mounter, L. A. & Whittaker, V. P. (1953). *Biochem. J.* **54**, 551.
 Myers, D. K. (1949). M.A. Thesis: University of Toronto.
 Myers, D. K. (1954). Ph.D. Thesis: University of Amsterdam.
 Myers, D. K. (1956). *Biochem. J.* **62**, 556.
 Myers, D. K. & Mendel, B. (1949). *Proc. Soc. exp. Biol., N.Y.*, **71**, 357.
 Myers, D. K., Schotte, A. & Mendel, B. (1955*a*). *Biochem. J.* **60**, 481.
 Myers, D. K., Schotte, A., Boer, H. & Borsje-Bakker, H. (1955*b*). *Biochem. J.* **61**, 521.
 Neurath, H. (1952). In *Modern Trends in Physiology and Biochemistry*, p. 453. New York: Academic Press.
 Neurath, H. & Schwert, G. W. (1950). *Chem. Rev.* **46**, 69.
 Ord, M. G. & Thompson, R. H. S. (1951). *Biochem. J.* **49**, 191.
 Racker, E. & Krimsky, I. (1948). *J. biol. Chem.* **173**, 519.
 Richter, D. & Croft, P. G. (1942). *Biochem. J.* **36**, 746.
 Smith, E. L. (1951*a*). In *Enzymes and Enzyme Systems*, p. 47. Cambridge, Mass., U.S.A.: Harvard University Press.
 Smith, E. L. (1951*b*). *Advance. Enzymol.* **12**, 191.
 Suzuoki-Ziro & Suzuoki-Tuneko (1954). *Nature, Lond.*, **173**, 83.
 Whittaker, V. P. (1951). *Physiol. Rev.* **31**, 312.
 Wilson, I. B. (1952). *J. biol. Chem.* **197**, 215.
 Zeller, E. A. (1945). *Helv. physiol. acta*, **3**, C47.

Nucleotides of Human Blood Platelets

BY P. FANTL AND H. A. WARD

Baker Medical Research Institute and Alfred Hospital Clinical Research Unit, Melbourne, Australia

(Received 21 February 1956)

Blood platelets take part in several reactions connected with haemostasis, and haemorrhagic conditions have been recognized in which platelet abnormalities appear to be the cause of the disturbance. The diagnosis of the defect is supported by microscopic examination of platelets and by laboratory tests designed to determine platelet activity in haemostatic reactions. During a study of the latter,

it was found necessary to obtain information about purine and pyrimidine derivatives in platelets. There is considerable uncertainty in the literature concerning the presence of nucleic acids and their breakdown products in blood platelets. The earlier literature has been reviewed by Maupin (1954). Platelets are believed to contain ribonucleic acid (RNA) by Tinacci & Cessi (1952) on the basis of