REFERENCES

- Aberdalden, E. & Heyns, K. (1934). *Ber. dtsch. chem. Ges.* **67**, 530.
- Adams, P. T. & Tolbert, B. M. (1952). *J. Amer. chem. Soc.* **74**, 6272.
- Arnstein, H. R. V. & Clubb, M. E. (1955). *Biochem. J.* **60**, xxxiv.
- Arnstein, H. R. V., Clubb, M. & Grant, P. T. (1954). *Radio-isotope Conference*, vol. 1, p. 306. Ed. by Johnston, J. E. London: Butterworths Scientific Publications.
- Arnstein, H. R. V. & Grant, P. T. (1954*a*). *Biochem. J.* **57**, 353.
- Arnstein, H. R. V. & Grant, P. T. (1954*b*). *Biochem. J.* **57**, 360.
- Calvin, M., Heidelberger, C., Reid, J. C., Tolbert, B. M. & Yankwich, P. F. (1949). *Isotopic Carbon*, p. 177. New York: John Wiley and Sons Inc.; London: Chapman and Hall, Ltd.
- Cornforth, J. W. (1949). In *The Chemistry of Penicillin*, p. 783. Ed. by Clarke, H. T., Johnson, J. R. & Robinson, Sir Robert. Princeton: Princeton University Press.
- du Vigneaud, V. & Patterson, W. I. (1935). *J. biol. Chem.* **109**, 97.
- Emerson, E. L., Puziss, M. & Knight, S. G. (1950). *Arch. Biochem.* **25**, 299.
- Fischer, E. (1906). *Ber. dtsch. chem. Ges.* **39**, 2320.
- Greenstein, J. P. (1954). *Advanc. Protein Chem.* **9**, 121.
- Halliday, W. J. & Arnstein, H. R. V. (1956). *Biochem. J.* **64**, 380.
- Hockenhull, D. J. D., Ramachandran, K. & Walker, T. K. (1949). *Arch. Biochem.* **23**, 160.
- Horowitz, N. H. (1947). *J. biol. Chem.* **171**, 255.
- Humphrey, J. H. & Lightbown, J. W. (1952). *J. gen. Microbiol.* **7**, 129.
- Knight, S. G. (1948). *J. Bact.* **55**, 401.
- Mozingo, R., Wolf, D. E., Harris, S. A. & Folkers, K. (1943). *J. Amer. chem. Soc.* **65**, 1013.
- Neuberg, C. & Kerb, J. (1912). *Biochem. Z.* **40**, 498.
- Owen, L. N. (1949). *J. chem. Soc.* p. 236.
- Popják, G. (1950). *Biochem. J.* **46**, 560.
- Ramachandran, K. & Walker, T. K. (1951). *Arch. Biochem. Biophys.* **31**, 224.
- Rufenach, K. (1952). *Helv. chim. acta*, **35**, 762.
- Smith, E. L. & Hockenhull, D. J. D. (1952). *J. appl. Chem.* **2**, 287.
- Steiger, R. E. (1944). *J. org. Chem.* **9**, 396.
- Stein, W. H. & Moore, S. (1949). *Cold Spr. Harb. Symp. quant. Biol.* **14**, 179.
- Stevens, C. M., Inamine, E. & De Long, C. W. (1956). *J. biol. Chem.* **219**, 405.
- Stevens, C. M., Vohra, P. & De Long, C. W. (1954). *J. biol. Chem.* **211**, 297.
- Stevens, C. M., Vohra, P., Inamine, E. & Roholt, D. A. (1953). *J. biol. Chem.* **205**, 1001.
- Stevens, C. M., Vohra, P., Moore, J. E. & De Long, C. W. (1954). *J. biol. Chem.* **210**, 713.
- Vilenkina, G. Y. (1949). *C.R. Acad. Sci., U.R.S.S.*, **69**, 385.

Glycerylphosphorylcholine and Phosphorylcholine in Semen, and their Relation to Choline

By R. M. C. DAWSON

Biochemistry Department, A.R.C. Institute of Animal Physiology, Babraham, Cambridge

AND T. MANN AND I. G. WHITE

*A.R.C. Unit of Reproductive Physiology and Biochemistry, Molteno Institute
and Department of Veterinary Clinical Studies, University of Cambridge*

(Received 13 August 1956)

Mammalian seminal plasma represents one of the richest natural sources of free choline and water-soluble choline compounds (Mann, 1954). However, as shown by Kahane & Lévy (1936, 1937), free choline accumulates in human seminal plasma only after ejaculation, as the result of a hydrolytic process which takes place in voided semen, on standing. Investigations on the nature of the choline precursor, which were carried out independently in the laboratories of Kahane in France (Kahane & Lévy, 1938, 1945, 1949; Diamant, Kahane & Lévy, 1952, 1953; Diamant, 1954, 1955) and of Lundquist in Denmark (Lundquist, 1946, 1947*a, b*, 1953), led to somewhat conflicting results, particularly as regards (i) the ratio between free choline and

the simultaneously liberated inorganic phosphate, and (ii) the nature of the phosphorylated choline precursor. It is the purpose of this paper to report the results of an investigation of the nature and distribution of choline compounds in the semen and male accessory secretions of various species. As well as human semen, the study included material from monkey, ram, goat, bull, boar, stallion, rat, rabbit, hedgehog and cock.

EXPERIMENTAL

Collection of semen and of secretions from male accessory organs. Human semen from donors at the Fertility Clinic was kindly made available by Dr H. A. Davidson. The semen of ram, bull, boar, goat, rabbit and stallion was

collected by means of an artificial vagina (Walton, 1945); the two parts of semen, the sperm and seminal plasma, were separated by centrifuging. Cock semen was obtained by the method of abdominal massage (Burrows & Quinn, 1939). The secretory fluids from male accessory organs were obtained at autopsy.

Whenever possible semen was frozen in solid CO_2 (-78°) immediately after ejaculation and stored at this temperature until it was used for determinations of glycerylphosphorylcholine (GPC) or phosphorylcholine. Evidence was obtained that in human semen there was a breakdown of phosphorylcholine during storage for a few days in a deep-freeze (-15°).

Preparation of suspensions of washed spermatozoa. Ram semen (1.0–1.5 ml.) was diluted with 6–8 vol. of a diluent, in a conical centrifuge tube, the spermatozoa were separated by centrifuging at about 1000 *g*, and washed twice with 6–8 vol. of fresh diluent. The diluents were: (i) a calcium-free Ringer's solution, composed of 100 ml. of 0.9% NaCl, 4 ml. of 1.15% KCl, 1 ml. of 2.11% KH_2PO_4 , 1 ml. of 3.82% $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$ and 2 ml. of 1.3% NaHCO_3 ; (ii) the same solution but with phosphate omitted ('P-free Ringer solution'); (iii) a calcium-free Ringer's solution supplemented with 20 ml. of 0.25M phosphate buffer, pH 7.4 ('Ringer-phosphate', Mann, 1946); (iv) a 'phosphate-diluent' (White, Blackshaw & Emmens, 1954), containing 0.032M- $\text{NaH}_2\text{PO}_4 \cdot \text{H}_2\text{O}$, 0.048M- $\text{Na}_2\text{HPO}_4 \cdot 12\text{H}_2\text{O}$, 0.040M-NaCl and 0.004M-KCl. The final concentration of spermatozoa in the washed sperm suspensions was about 3×10^8 cells/ml.

Chromatographic detection of GPC and phosphorylcholine. The sample of semen or seminal plasma (0.1–0.75 ml. according to the acid-soluble P content) was introduced into 4 ml. of boiling water, and maintained at 100° for 3 min. Ethanol (7 ml.) was added and the mixture centrifuged. The slightly turbid supernatant was shaken with 30 ml. of CHCl_3 . The milky upper layer was centrifuged for 20 min. at 10 000 *g* and the clear supernatant was rapidly passed through a column of Amberlite IRC 50 (H) (8 cm. $\times$ 0.8 cm.²) to remove cations. The effluent (25 ml.) was shaken with 75 ml. of *isobutanol* and the clear aqueous layer which separated was adjusted to pH 7 with NH_3 soln., taken down to a small volume *in vacuo*, and then applied to the origin of the paper chromatogram. Sometimes, especially with bull, stallion and boar semen, it was difficult to apply the final solution to the chromatogram owing to the presence of a gummy material, but this had no noticeable effect on the quality of the chromatograms developed. The methods of chromatography used, and the detection of the P-containing spots, have been previously described (Dawson, 1954, 1955). The solvents used were phenol saturated with 0.1% NH_3 soln. in one dimension and ethanol- NH_3 soln. (sp.gr. 0.880)–water (6:3:1, by vol.) in the second. It was found that the best separations were obtained when the second solvent was run at a temperature below 20° . The GPC spot had approximate R_F values of 0.88 and 0.55 in the respective solvents, and those for phosphorylcholine were 0.89 and 0.27. The identity of the spots was confirmed by running the sample simultaneously with synthetic compounds as markers.

Determination of GPC and phosphorylcholine. The frozen sample was either rapidly thawed and then pipetted or it was weighed in the frozen state on a torsion balance. It was then introduced into 2 ml. of 0.5% serum albumin solution at 0° and immediately precipitated with 1 ml. of 10% (w/v)

perchloric acid solution. Serum albumin was added to facilitate the precipitation of lipid material. The suspension was centrifuged and the supernatant filtered and stored at 0° . Two samples (1 ml.) were each mixed with 0.25 ml. of 5N-HCl; one was then kept at 0° and the other hydrolysed at 100° for 20 min. to release the acid-labile choline from GPC. The free choline in the unhydrolysed and hydrolysed samples was estimated by the method of Appleton, La Du, Levy, Steel & Brodie (1953), the choline being directly precipitated with the tri-iodide reagent. The entire estimation, including colorimetry, was carried out at 0° . Recovery of choline and GPC, with standards of these two substances, was linear between 0 and 200 μg . and 0 and 450 μg . respectively. Recovery of GPC added to dialysed ram seminal plasma was linear between 0 and 360 μg .

Phosphorylcholine was estimated in human semen samples by estimating the choline present before and after incubation for 90 min. at 37° . Under these conditions phosphorylcholine is completely hydrolysed by the acid phosphatase in the semen and GPC is not affected.

Isolation of GPC from ram seminal plasma. Seminal plasma (40 ml.), obtained from ram semen by centrifuging, was precipitated with 6 vol. of ethanol, and the filtered water-clear ethanolic extract concentrated *in vacuo* to a volume of 6 ml. After removing the small water-insoluble residue by centrifuging, the concentrate was poured, under constant stirring, into 6 vol. of acetone, and 1 hr. later the oily, brownish coloured bottom layer which formed on standing was separated from the colourless supernatant solution by decantation. The oily fraction was extracted with three successive 5 ml. portions of boiling ethanol, the mixture being allowed to boil each time for a few minutes. The combined ethanolic extracts, representing a total of 52 mg. of total P, were filtered while still hot, and treated under constant stirring with 0.6 g. of cadmium chloride dissolved in 0.8 ml. of water and diluted with 20 ml. of ethanol; a precipitate began to form immediately. The mixture was left standing for a few hours first at room temp., and then at 0° , overnight. The slightly yellowish and mostly amorphous precipitate was separated by centrifuging, and washed with ethanol. After draining off all remaining ethanol, the precipitate was dissolved in 8 ml. of water, and to the aqueous solution 95% (v/v) ethanol was added gradually until crystallization just began. After 5 days (at 0°), the snow-white crystalline material was collected in a sintered-glass crucible, washed with cold 80% (v/v) ethanol and dried in air on filter paper to constant weight.

Glycerol determination. With protein-free filtrates from samples deproteinized with $\text{Ba}(\text{OH})_2$ and ZnSO_4 , glycerol was estimated by the colorimetric periodate method of Lambert & Neish (1950) as modified by Ryley (1955). In filtrates from semen, fructose, which interferes with determinations of glycerol, was removed by the copper-lime precipitation technique, but this procedure involved small losses of glycerol owing to adsorption upon the copper-lime precipitate. GPC, analysed directly by the periodate method, gives about one-half the colour of the equivalent amount of free glycerol. An additional means of distinguishing between free and bound glycerol was obtained through chromatographic analysis with the procedure of Trevelyan, Procter & Harrison (1950).

Other determinations. Phosphorus was estimated by the method of Berenblum & Chain (1938), after the organic P-compounds had been converted into orthophosphate by

oxidation with perchloric acid. Fatty acid ester bonds were determined by the procedure of Stern & Shapiro (1953) with triacetin as a standard. Water-soluble ninhydrin-reacting N was measured by the method of Moore & Stein (1948), with the precipitation procedure described by Dawson (1956*a*). The O_2 uptake was measured manometrically at 37°.

Chemicals. Ovolecithin was prepared on an alumina column by the method of Hanahan (1954). Any acid-soluble impurities which might have been present were removed by washing with 0.05M-MgCl₂ solution according to the procedure of Folch, Ascoli, Lees, Meath & LeBaron (1951). Lysolecithin was prepared enzymically with cobra venom (Hanahan, Rodbell & Turner, 1954). L- α -Glycerylphosphorylcholine was prepared by the method of Dawson (1956*b*), phosphorylcholine by the method of Plimmer & Burch (1937) and phosphorylethanolamine was synthesized by the procedure of Outhouse (1937).

Gifts of L- α -glycerylphosphorylethanolamine (Professor E. Baer), phosphatidylethanolamine (Drs Lea and Rhodes) and Indian cobra venom (Zoological Society of London) are gratefully acknowledged.

RESULTS

GPC and phosphorylcholine in semen and accessory gland secretions

All semen samples and accessory gland secretions obtained were examined for water-soluble phosphate esters by two-dimensional paper chromatography as described in the Experimental section. In every sample the major portion of the water-soluble P visible on the chromatogram could be accounted for as either GPC or phosphorylcholine, or both. Figs. 1*a* and *b* show typical examples of this, being chromatograms prepared from human and ram semen respectively. The amount of GPC and phosphorylcholine present in the semen varied considerably between the species, a roughly comparative assessment of the quantity as judged from the size and intensity of the chromatographic spots and the quantity of material used being given in Table 1. It is seen that the semen of the ram, goat, boar, stallion and rabbit contained exclusively GPC, whereas in man there was a small amount of GPC but a much larger quantity of phosphorylcholine. Our results concerning human semen are in agreement with those of Lundquist (1947*a, b*), who was the first to isolate phosphorylcholine from fresh human semen; they do not support the possibility of phosphorylcholine arising from GPC during the isolation procedure (Diamant, 1955). Phosphorylcholine was also observed by us in small amounts in bull and monkey vesicular secretions, hedgehog accessory secretions and cock semen. In the boar and in the bull there was a particularly large concentration of GPC in the epididymal secretion, much larger than in any other secretion of the same species. The high content of GPC in the epididymal secretion might be taken to indicate that either the epididymis or the testis of the boar

and bull is concerned with the production of seminal GPC. Because of this possibility, a few preliminary experiments were carried out with these two organs from a boar, and evidence was obtained that both contain phospholipase *B*, an enzyme capable of producing GPC from lysolecithin. Whether, however, this activity accounts for the formation of GPC under conditions *in vivo* remains to be ascertained.

The values for the GPC concentration, determined as acid-labile choline (Table 1), agree well with the comparative visual estimates obtained by chromatographic means. Although the iodide method for estimating choline is much more sensitive than the reineckate method it is rather unspecific, and consequently the values obtained with it have to be accepted with some reserve. Nevertheless, when GPC was determined in two samples of ram semen by the reineckate-precipitation method, the values were found to correspond closely with those obtained by iodide precipitation.

Fig. 1. Paper chromatograms prepared from the aqueous extracts of (a) human semen, (b) ram seminal plasma. I, Glycerylphosphorylcholine; II, phosphorylcholine; III, probably a nucleotide; IV, inorganic phosphate.

Moreover, in ram semen, which chromatographically shows very little acid-soluble P apart from GPC, the labile-choline values were in good agreement with the estimates of total acid-soluble P.

Glycerylphosphorylethanolamine, which accompanies GPC in nearly all mammalian tissues (Campbell & Work, 1952; Ansell & Norman, 1953; Dawson, 1955), was detected only occasionally on chromatograms, and then only in traces, amounting to less than 1% of the GPC content (Table 1). Phosphorylethanolamine was not detected in any of the specimens.

In view of the chromatographic and analytical results reported above, which showed that in several species, and particularly in the ram, GPC accounts for practically all the bound choline and acid-soluble P in seminal plasma, an attempt was made to isolate pure GPC from ram semen. The method used for this purpose was based on the procedure described by Baer & Kates (1948) for the preparation of the crystalline cadmium salt of

synthetic GPC; it is described in the Experimental section. The crystalline cadmium salt of glycerylphosphorylcholine isolated from 40 ml. of ram seminal plasma weighed 470 mg. It was free from free choline and inorganic phosphate, and contained 23.3% of acid-hydrolysable bound choline; calc. for $(C_8H_{22}O_7NP)(CdCl_2) \cdot 2H_2O$: 24.5% bound choline; the total phosphorus content was 6.00% P; calc.: 6.27%. Further proof concerning the purity of the substance isolated from ram seminal plasma was obtained by hydrolysing a sample for 20 min. at 100° in N-HCl. This resulted in complete hydrolysis to free choline and phosphoglycerol, which was shown chromatographically to be the sole P compound. No free fatty acids were liberated by hydrolysis. The small impurity present in the preparation of the cadmium salt of seminal GPC appeared to contain glutamic acid; no other amino acids were detected chromatographically, and the preparation was also free from carbohydrate, citric acid and phosphorylcholine.

Table 1. *Concentration of GPC, phosphorylcholine and glycerylphosphorylethanolamine in the semen and accessory-gland secretions of various species*

Species	Material	GPC		Phosphorylcholine		Glycerylphosphorylethanolamine. Chromatographic determination
		Chromatographic determination	Acid-labile choline (mg. GPC/100 ml.)	Chromatographic determination	Choline released on incubation at 37° (mg. phosphorylcholine/100 ml.)	
Ram	Semen	+++++	1185, 1362, 1942	0	—	Trace seen in one sample
	Seminal plasma	+++++	1601, 2040, 1770	0	—	—
Bull	Semen	+++	460, 237	Trace seen in one sample	—	Trace
	Seminal plasma	+++	496, 420, 375, 352, 110*	0	—	0
	Vesicular secretion	Trace	—	+ (2 samples)	—	0
	Epididymal secretion	+++++	1490*	Trace	—	0
	Ampullar fluid	++	94*	Trace	—	0
Goat	Semen	+++++	1382, 1550	0	—	Trace
Boar	Seminal plasma	++	108, 235	0	—	Trace
	Vesicular secretion	++	190	0	—	Trace
	Epididymal secretion	+++++	3060	0	—	Trace
Stallion	Semen	+	38, 113	0	—	Trace
	Ampullar secretion	+	120	0	—	0
Man	Semen	+	59, 59, 54, 90	+++	301, 286, 256, 380	0
Rat	Vesicular secretion	+++	654	Trace	—	0
Rabbit	Semen	++	215, 264, 370	0	—	0
Hedgehog	Secretion of 'prostate I and II'	++	—	++	—	0
	Secretion of 'prostate III'	0	—	+++	—	0
Monkey	Vesicular secretion	0	—	++	—	0
Cock	Semen	0	—	+	—	0

* Samples from the same bull.

Metabolic behaviour of GPC and phosphorylcholine

In view of the large amount of GPC detected in ram semen and seminal plasma it was of interest to investigate whether ram spermatozoa themselves could either produce or metabolize GPC. Washed suspensions of spermatozoa were therefore incubated for 3 hr. at 37°, aerobically or anaerobically, but no liberation of GPC, inorganic P or free choline was observed. Similarly, on incubating the washed sperm suspensions with added GPC there was no disappearance of the substrate or significant production of free choline or inorganic P. The same result was obtained when whole ram semen was incubated, either as such or after diluting it with P-free Ringer solution or with the phosphate diluent. The incubation of ram seminal plasma by itself had, again, no effect on the content of GPC, and it did not result in the appearance of free choline, glycerol or inorganic phosphate.

The semen and seminal plasma of the bull, like those of the ram, were inactive towards GPC. In three experiments in which bull semen or seminal plasma was incubated for periods of 3–24 hr., aerobically or anaerobically, with or without penicillin, there was no liberation of any appreciable quantities of free choline, glycerol or inorganic P, nor was there a disappearance of GPC (Table 2). This finding contrasts with the observation by Lundquist (1947*a, b*) that choline and inorganic P appear in bull seminal plasma on incubation.

Lundquist has demonstrated that, in human semen, phosphorylcholine breaks down rapidly to choline and inorganic phosphate within a short time after ejaculation, owing to the action of the powerful acid (prostatic) phosphatase which comes in contact with the substrate at the time of ejacula-

tion. This observation has been fully confirmed in the present study. When fresh human semen was incubated for 90 min. at 37°, phosphorylcholine disappeared completely, and at the same time, equivalent amounts of free choline and inorganic phosphate were formed. The ease with which phosphorylcholine breaks down was even more evident in an experiment where freshly collected human semen was stored for 7 days at –15° instead of –78°. This period of storage was sufficient to cause the complete breakdown of the phosphorylcholine in the semen. Phosphorylcholine added to human seminal plasma also undergoes a rapid dephosphorylation; its complete disappearance was observed, for instance, in an experiment in which 0.5 ml. of a solution containing 6 mg. of sodium phosphorylcholine was incubated for 2 hr. at 37°, with 0.1 ml. of human seminal plasma and 0.4 ml. of veronal buffer at pH 5.7, 7 and 8.5 respectively. The ability of human seminal plasma to dephosphorylate phosphorylcholine exceeds markedly that of ram seminal plasma and even more so that of the bull; this behaviour is in agreement with the known acid phosphatase activity levels in the semen of these three species.

Apart from the large quantity of phosphorylcholine, human semen contains some GPC, as was mentioned before (Table 1). Unlike phosphorylcholine, however, GPC does not undergo the same rapid breakdown on incubation, and it can be detected chromatographically even after prolonged incubation of human semen (6 hr. at 37°). The difference between phosphorylcholine and GPC becomes even more pronounced when one compares the behaviour of the two substrates after they have been added as pure substances to the seminal plasma; incubation for 2 hr. at 37°, which as stated

Table 2. *Effect of incubation at 37° on the content of free choline and inorganic phosphate in bull and ram semen*

Sample	Conditions of incubation	Free choline (mg./100 ml.)	Inorganic P (mg./100 ml.)
Bull semen	No incubation	24	—
	3 hr., aerobic	28	—
	3 hr., anaerobic	28	—
Bull semen	No incubation	24	—
	3 hr., aerobic	26	—
	3 hr., anaerobic	36	—
Bull seminal plasma	No incubation	21	3
	18 hr., aerobic with penicillin	19	4
	18 hr., aerobic	24	3
Bull semen	No incubation	47	—
	24 hr., aerobic with penicillin	47	—
	24 hr., aerobic	46	—
Ram semen	No incubation	65	17
	3 hr., anaerobic	68	19
Ram semen diluted with P-free Ringer	No incubation	47	20
	3 hr., aerobic	53	32

previously hydrolyses phosphorylcholine completely, caused no breakdown of an added equivalent quantity of GPC. Thus the human seminal plasma appears to be devoid of any GPC diesterase, an enzyme which is distributed widely in mammalian tissues (Dawson, 1956b). The fact that acid phosphatase by itself is unable to attack GPC has been stated also by Schmidt, Bessman & Thannhauser (1953).

Effect of lecithin and lysolecithin on spermatozoa

In view of the close relationship of GPC to lecithin and lysolecithin, and the recent work on the 'protective' action of lecithin and related lipids on the sperm cells (e.g. Mayer, 1955), a series of experiments was carried out in order to compare the metabolic behaviour of GPC with that of lecithin and lysolecithin.

When pure lecithin, emulsified in Ringer solution, was incubated aerobically with a suspension of washed ram spermatozoa, there was sometimes, but not always, a considerable increase in O_2 utilization, which exceeded the control values by 40–380%. Choline, GPC, phosphatidylethanolamine and triolein did not produce a similar effect on the respiration, and as the lecithin had been carefully purified from acid-soluble impurities such as amino acids, the effect must be considered as a specific stimulation by lecithin. However, the increased O_2 consumption brought about by lecithin was not associated with the liberation of free choline or GPC, or phosphorylcholine; and it did not result in any decrease in the number of fatty acid ester bonds in the lecithin itself. It would thus appear that the increased O_2 consumption was not due to the enzymic hydrolysis of ester bonds in the lecithin and the subsequent metabolic oxidation of the hydrolytic products. Similar stimulatory effects of lecithin, not involving any apparent oxidation of lecithin, have been observed in studies on the respiration of liver mitochondria (Rodbell & Hanahan, 1955; Marinetti & Stotz, 1956), but in these experiments similar effects were obtained with phosphatidylethanolamine and lysolecithin. In the present study with ram spermatozoa, phosphatidylethanolamine neither increased the respiration nor was it metabolized by the sperm cells to produce ethanolamine, phosphorylethanolamine or glycerylphosphorylethanolamine.

Lysolecithin, on the other hand, had a marked inhibitory action on the respiration of washed ram spermatozoa, even after the lysolecithin preparation had been autoclaved sufficiently to destroy the enzymic activity of any residual cobra venom which might have been left during its preparation. This decrease in respiration was associated with a loss of motility as well as with a definite morphological change in the sperm cells, namely, a detachment of

the sperm head from the midpiece–tail. Cobra venom itself, which, as is well known, contains an active phospholipase A, diminished the endogenous respiration of ram spermatozoa; although no acid-soluble P was liberated, there was a decrease in the fatty acid ester bonds of the sperm suspensions, indicating that the venom had probably attacked the lipid coating of the sperm cells. It is interesting to note in this connexion that the respiratory enzyme systems of isolated liver-mitochondria preparations are also known to be sensitive to a short incubation with phospholipase-containing venoms (Nygaard & Sumner, 1953; Nygaard, Diansani & Bahr, 1954).

When human semen was incubated with lecithin for 2–7 hr. at 37°, at three different pH values, namely, 5.7, 7 and 8.5, no liberation of any acid-soluble P compounds was observed, above that found in the controls. It was thus not possible to confirm the claim of Cuzzocrea (1955) that an active phospholipase C is present in human semen.

DISCUSSION

It has been recognized for some time that mammalian seminal plasma is characterized by a high content of a phosphorus compound soluble both in trichloroacetic acid and ethanol, which does not readily liberate orthophosphate on acid hydrolysis. The similarity of this compound to glycerylphosphorylcholine was first stressed by Kahane & Lévy (1938, 1945, 1949), but it was only in 1952 that the French investigators actually accomplished the isolation of glycerylphosphorylcholine, as a ferric salt, from the seminal vesicle secretion of rats (Diament *et al.* 1952). A similar result was obtained by Lundquist (1953), who studied the vesicular secretions of rat, guinea pig and rabbit. The Danish authors also succeeded, a few years earlier, in isolating from human seminal plasma another P compound that was difficult to hydrolyse and in identifying it as phosphorylcholine (Lundquist, 1947*a, b*). This finding, however, was criticized by the French investigators, who felt that the chemical method employed for the isolation of phosphorylcholine from human semen was so drastic that it could have caused a breakdown of glycerylphosphorylcholine to phosphorylcholine.

Our own results show that in certain species, notably man, phosphorylcholine is undoubtedly a natural, major constituent of the seminal plasma. Glycerylphosphorylcholine, however, also occurs in human semen but at a much lower concentration. In the bull, ram, goat, boar and stallion, on the other hand, the bulk of the 'bound choline' and 'difficultly-hydrolysable P' of seminal plasma was identified as glycerylphosphorylcholine, to the almost complete exclusion of phosphorylcholine.

The two phosphorus-containing compounds differ characteristically in their behaviour towards seminal phosphatase. Whereas phosphorylcholine is broken down readily immediately after ejaculation, giving rise to free choline and inorganic phosphate, glycerylphosphorylcholine is stable and does not break down appreciably. Although glycerylphosphorylcholine and phosphorylcholine accounted together for the major portion of the water-soluble organically bound phosphate in the seminal plasma of all species examined, chromatographic evidence was obtained for the existence in certain species of small amounts of some other P compounds. It is interesting to note, however, that glycerylphosphorylethanolamine, which accompanies glycerylphosphorylcholine in most mammalian tissues, is detectable in seminal plasma in trace amounts only.

The occurrence of phosphorylcholine and glycerylphosphorylcholine in the seminal plasma naturally raises the problem of the mechanism of their formation in the male accessory glands of reproduction. A possibility which merits particular attention is that these compounds may be derived from the metabolism of phospholipids in organs such as the epididymis and seminal vesicle.

SUMMARY

1. Glycerylphosphorylcholine and phosphorylcholine were found to account for the bulk of 'water-soluble bound choline' and 'acid-soluble, difficultly-hydrolysable phosphorus' in the seminal plasma of various mammals. In man, the content of phosphorylcholine exceeded considerably that of glycerylphosphorylcholine, whereas in ram, bull, boar, goat, stallion and rabbit practically the whole of bound choline was accounted for as glycerylphosphorylcholine. Bull and boar possess a particularly rich source of glycerylphosphorylcholine in the epididymis. Cock semen contained a small quantity of phosphorylcholine but no glycerylphosphorylcholine.

2. In addition to chromatographic means, glycerylphosphorylcholine was also identified by chemical isolation; 470 mg. of the crystalline cadmium salt was obtained from 40 ml. of ram seminal plasma. Glycerylphosphorylethanolamine was detected in trace amounts in some samples of seminal plasma, but phosphorylethanolamine was altogether absent.

3. Phosphorylcholine was readily dephosphorylated in semen after ejaculation. Glycerylphosphorylcholine was neither dephosphorylated nor changed in any other way by spermatozoa. Lecithin usually stimulated the oxygen uptake of spermatozoa, but this was not associated with an enzymic hydrolysis of ester bonds or the liberation of acid-

soluble metabolic products. Lysolecithin, on the other hand, had a marked inhibitory effect on sperm respiration and motility.

REFERENCES

- Ansell, G. B. & Norman, J. M. (1953). *Biochem. J.* **55**, 768.
 Appleton, H. D., La Du, B. N., Levy, B. B., Steel, J. M. & Brodie, B. D. (1953). *J. biol. Chem.* **205**, 803.
 Baer, E. & Kates, M. (1948). *J. Amer. chem. Soc.* **70**, 1394.
 Berenblum, I. & Chain, E. (1938). *Biochem. J.* **32**, 295.
 Burrows, W. H. & Quinn, J. P. (1939). *Artificial Insemination of Chickens and Turkeys. Dep. Circ. U.S. Dep. Agric.* no. 525.
 Campbell, P. N. & Work, T. S. (1952). *Biochem. J.* **50**, 449.
 Cuzzocrea, G. (1955). *G. Biochim.* **4**, 363.
 Dawson, R. M. C. (1954). *Biochim. biophys. Acta*, **14**, 374.
 Dawson, R. M. C. (1955). *Biochem. J.* **59**, 5.
 Dawson, R. M. C. (1956a). *Biochem. J.* **64**, 13.
 Dawson, R. M. C. (1956b). *Biochem. J.* **62**, 689.
 Diamant, M. (1954). Doctor Dissertation: University of Paris.
 Diamant, M. (1955). *Bull. Soc. Chim. biol., Paris*, **37**, 1195, 1205.
 Diamant, M., Kahane, E. & Lévy, J. (1952). *C.R. Acad. Sci., Paris*, **235**, 1058.
 Diamant, M., Kahane, E. & Lévy, J. (1953). *Arch. Sci. physiol.* **7**, 45.
 Folch, J., Ascoli, I., Lees, M., Meath, J. A. & LeBaron, F. N. (1951). *J. biol. Chem.* **191**, 833.
 Hanahan, D. J. (1954). *J. biol. Chem.* **211**, 321.
 Hanahan, D. J., Rodbell, M. & Turner, D. L. (1954). *J. biol. Chem.* **206**, 431.
 Kahane, E. & Lévy, J. (1936). *C.R. Acad. Sci., Paris*, **202**, 2186.
 Kahane, E. & Lévy, J. (1937). *Bull. Soc. Chim. biol., Paris*, **19**, 959.
 Kahane, E. & Lévy, J. (1938). *C.R. Acad. Sci., Paris*, **207**, 747.
 Kahane, E. & Lévy, J. (1945). *Bull. Soc. Chim. biol., Paris*, **27**, 544, 553.
 Kahane, E. & Lévy, J. (1949). *J. Physiol. Path. gén.* **41**, 183.
 Lambert, M. & Neish, A. C. (1950). *Canad. J. Res. B*, **28**, 83.
 Lundquist, F. (1946). *Nature, Lond.*, **158**, 710.
 Lundquist, F. (1947a). *Acta physiol. scand.* **13**, 322.
 Lundquist, F. (1947b). *Acta physiol. scand.* **14**, 263.
 Lundquist, F. (1953). *Nature, Lond.*, **172**, 587.
 Mann, T. (1946). *Biochem. J.* **40**, 481.
 Mann, T. (1954). *The Biochemistry of Semen*. London: Methuen.
 Marinetti, G. & Stotz, E. (1956). *J. biol. Chem.* **217**, 745.
 Mayer, D. T. (1955). *Reproduction and Infertility, Mich. St. Univ. Centennial Symp. Rep.*, p. 45.
 Moore, S. & Stein, W. H. (1948). *J. biol. Chem.* **176**, 367.
 Nygaard, A. P., Dianzani, M. U. & Bahr, G. F. (1954). *Exp. Cell Res.* **6**, 453.
 Nygaard, A. P. & Sumner, J. B. (1953). *J. biol. Chem.* **200**, 723.

- Outhouse, E. L. (1937). *Biochem. J.* **31**, 1459.
 Plimmer, R. H. A. & Burch, W. J. N. (1937). *Biochem. J.* **31**, 398.
 Rodbell, M. & Hanahan, D. J. (1955). *J. biol. Chem.* **214**, 595.
 Ryley, J. F. (1955). *Biochem. J.* **59**, 353.
 Schmidt, G., Bessman, M. J. & Thannhauser, S. J. (1953). *J. biol. Chem.* **203**, 849.
 Stern, I. & Shapiro, B. (1953). *Brit. J. clin. Path.* **6**, 158.
 Trevelyan, W. E., Procter, D. P. & Harrison, J. S. (1950). *Nature, Lond.*, **166**, 444.
 Walton, A. (1945). *Notes on Artificial Insemination of Sheep, Cattle and Horses*. London: Holborn Surgical Instrument Co.
 White, I. G., Blackshaw, A. W. & Emmens, C. W. (1954). *Aust. vet. J.* **30**, 85.

Glycerol Metabolism by Spermatozoa

BY T. MANN AND I. G. WHITE

*A.R.C. Unit of Reproductive Physiology and Biochemistry, Molteno Institute,
and Department of Veterinary Clinical Studies, University of Cambridge*

(Received 13 August 1956)

Although the inclusion of glycerol in media for freezing and long-term storage of bull semen is now a generally established practice, very little has been known hitherto about the mechanism underlying the uptake and metabolism of glycerol by spermatozoa. Recently, however, it has been shown by White, Blackshaw & Emmens (1954) that although glycerol does not affect the motility and respiration of spermatozoa in a fructose-containing medium, it increases markedly the oxygen uptake of washed ram- and bull-sperm suspensions, giving rise under certain conditions to an accumulation of lactic acid. Evidence was also brought forward recently by O'Dell, Flipse & Almquist (1956) that [^{14}C]glycerol is taken up readily and utilized by bovine spermatozoa, with the production of $^{14}\text{CO}_2$, and that the rate of glycerol uptake is higher in the absence than in the presence of seminal plasma. Further experiments on glycerol metabolism by ram spermatozoa are described in the present study, together with new observations on the metabolic behaviour of certain related substrates, and, particularly, phosphoglycerol, dihydroxyacetone and sorbitol (preliminary communication, Mann & White, 1956). These findings are discussed in the light of another recently completed study (Dawson, Mann & White, 1957), which describes the occurrence in ram seminal plasma of large quantities of bound glycerol in the form of glycerylphosphorylcholine.

MATERIAL AND METHODS

Semen. This was collected from ten rams by means of an artificial vagina and pooled for each experiment. Washed sperm suspensions were made by diluting 1.0–1.5 ml. of semen to 10 ml., in a tapered centrifuge tube, with one of the diluents listed below, centrifuging at about 1000 g, and washing the spermatozoa twice with the same amount of fresh diluent. The diluents were (i) the 'phosphate diluent' of White *et al.* (1954), containing 0.032 M- $\text{NaH}_2\text{PO}_4 \cdot \text{H}_2\text{O}$,

0.048 M- $\text{Na}_2\text{HPO}_4 \cdot 12\text{H}_2\text{O}$, 0.040 M- NaCl and 0.004 M- KCl , and (ii) the calcium-free 'Ringer solution for spermatozoa' (Mann, 1946); in some experiments phosphate was left out from the Ringer solution ('P-free Ringer solution'), and in some other experiments each 100 ml. of the Ringer solution was supplemented further with 20 ml. of 0.25 M phosphate buffer, pH 7.4, prepared by mixing 190 ml. of an aq. soln. of 7.1 g. of Na_2HPO_4 (or 17.9 g. of $\text{Na}_2\text{HPO}_4 \cdot 12\text{H}_2\text{O}$) with 10 ml. of N-HCl ('Ringer-phosphate solution').

Determinations. Glycerol was determined in protein-free filtrates from samples deproteinized with $\text{Ba}(\text{OH})_2$ and ZnSO_4 , by the method of Lambert & Neish (1950) as modified by Ryley (1955), and in some cases the identification of glycerol was carried out also by chromatography by the method of Trevelyan, Procter & Harrison (1950). The methods used for the determination of fructose, lactic acid, phosphorus, sperm respiration (in air, at 37°) and fructolysis (in N_2 , at 37°) were as described by Mann (1946). The chromatographic identification of fructose was carried out by the method of Jermyn & Isherwood (1949).

RESULTS

Aerobic formation of lactic acid from glycerol by spermatozoa

The rate at which glycerol (0.01 M) is utilized by washed ram spermatozoa at 37° was determined in nine experiments, a fresh sperm suspension in 'phosphate diluent' being used each time, with about 3×10^8 cells/ml. The disappearance of glycerol under these conditions was nearly linear over the 3 hr. experimental period. The rate of disappearance was 0.5–1 mg. of glycerol/ 10^9 sperm/hr., with an average of 0.8 mg. Throughout the period of 3 hr., the utilization of glycerol was accompanied by an increase in O_2 uptake of the order of 0.1–0.25 c.c. of O_2 / 10^9 sperm/hr. over that of the glycerol-free control sample. The disappearance of glycerol was accompanied by the accumulation of lactic acid, at a fairly linear rate of 0.2–0.6 mg./ 10^9 sperm/hr. Assuming that one atom of oxygen was required for