sulphone and the amino groups of α -amino acids proceeds exclusively *via* the formation of two vinyl groups in the sulphone, followed by the addition of the amine to the ethylenic bonds.

We are indebted to the British Empire Cancer Campaign for grants which have partly covered the cost of these investigations. We are also grateful to Dr T. E. Banks for advice and assistance in the use and maintenance of the mass spectrometer and to Dr V. C. E. Burnop for help with one stage of the synthesis of deuterio-mustard-gas sulphone and also for his assistance in some of the early investigations. The microanalyses were carried out by Miss J. Cuckney and the staff of the Microanalytical Laboratories of the Imperial College.

REFERENCES

- Banks, T. E., Bournsell, J. C., Francis, G. E., Hopwood, F. L. & Wormal, A. (1946). *Biochem. J.* **40**, 745.
Bournsell, J. C., Francis, G. E. & Wormal, A. (1946a). *Biochem. J.* **40**, 737.
Bournsell, J. C., Francis, G. E. & Wormal, A. (1946b). *Biochem. J.* **40**, 743.
Bournsell, J. C., Francis, G. E. & Wormal, A. (1946c). *Biochem. J.* **40**, 768.
Dubbs, C. A. (1953). *Analyt. Chem.* **25**, 828.
Francis, G. E., Mulligan, W. & Wormal, A. (1955). *Biochem. J.* **60**, 118.
Graff, J. & Rittenberg, D. (1952). *Analyt. Chem.* **24**, 878.

The Quantitative Estimation of Pyruvic and α -Oxoglutaric Acids by Paper Chromatography in Blood, Urine and Cerebrospinal Fluid

By B. McARDLE

Member of Scientific Staff, Medical Research Council, Guy's Hospital Medical School, London, S.E. 1

(Received 10 September 1956)

It is now recognized that the estimation of pyruvate by the Friedemann & Haugen method (1943) carries with it an element of uncertainty owing to the non-specific nature of this technique. This lack of specificity may well be unimportant for estimations in human blood and cerebrospinal fluid, in which, under normal conditions, other α -keto acids are present in small or negligible amounts, but if a clearer insight is to be gained into the disturbances of the Krebs cycle in disease a more specific determination of the individual α -keto acids is required. This is especially true if α -oxoglutaric and other α -keto acids are present in significant amounts as, for example, in the blood of certain animals such as the rat, in certain tissues and in urine. A specific procedure involving the formation of the 2:4-dinitrophenylhydrazones of the α -keto acids and their separation by paper chromatography was devised by Cavallini, Frontali & Toschi (1949a, b), and since then a number of modifications of their method have been published. These methods, however, with few exceptions have been tedious and time-consuming, and in few is it possible to assess their accuracy. The present method has been developed from that of El Hawary & Thompson (1953) and has given satisfactory results over three years. The number of extractions has been greatly reduced, and the need for evaporating the extracted hydrazones to dryness eliminated. The addition of ethanol to the ethyl acetate used for extracting the hydrazones has further shortened the procedure by greatly reducing interference due to emulsions, which are particularly apt to develop during blood determinations.

METHOD

Reagents. 2:4-Dinitrophenylhydrazine, 0.2% in 2N-HCl; ethanol, absolute; ethyl acetate (A.R.); ammonium hydroxide, 2.5N; metaphosphoric acid, 5 and 10% (w/v) freshly prepared from a stock 50% solution; solvent mixture: *n*-butanol-ethanol-N-NH₃ soln. (13:2.5, by vol.); sodium hydroxide-sodium carbonate solution, freshly prepared by mixing equal parts of 2.5N-NaOH (A.R.) and 10% (w/v) Na₂CO₃ (A.R.).

Blood. Immediately after withdrawal 5-7 ml. of blood is mixed with 20 ml. of 5% metaphosphoric acid in a weighed tube. After a few minutes the mixture is centrifuged.

Urine. If the specific gravity is 1.02 or over, 3 ml. of urine is added to 13 ml. of 5% metaphosphoric acid. Below this specific gravity, increasing quantities of urine are added to decreasing amounts of more concentrated metaphosphoric acid, the amount of urine depending on the specific gravity, and the final concentration of the acid being not less than 4%. If the urine contains protein this must first be precipitated and removed by centrifuging.

Cerebrospinal fluid. A volume (3 ml.) is mixed with 14 ml. of 5% metaphosphoric acid, and centrifuged.

Preparation and extraction of keto acid hydrazones. A volume (2 ml.) of the 2:4-dinitrophenylhydrazine in 2N-HCl is added to 16 ml. of the centrifugate or, with protein-free urine, of the clear solution, and the mixture is incubated for 20 min. at 38°. The mixture is then shaken in a 30 ml. separating funnel with 5 ml. of ethyl acetate and 5 ml. of ethanol for 0.75 min., and the layers are allowed to separate. The aqueous phase is re-extracted successively with 5 ml. and then with 1 ml. of ethyl acetate. With blood filtrates some degree of emulsification usually interferes with the complete separation of the two phases during the second and third extractions. If this occurs the emulsion is added to the first ethyl acetate extract and centrifuged at 2000 rev./min. for 1-2 min. The aqueous layer is returned to the separating funnel and extracted for the third time. The contents of the

separating funnel are then transferred to a centrifuge tube and spun for 1–2 min. The aqueous layer is sucked off and the ethyl acetate layer is added to the earlier extracts. The combined ethyl acetate phase is then washed by shaking with 20 ml. of water made faintly acid with 1–2 drops of N -HCl. If necessary, it is centrifuged for 1 min. and all the aqueous layer is carefully sucked off and discarded. Aqueous $2N$ - NH_3 soln. (1.5 ml.) is then added to the ethyl acetate and oxygen or compressed air is bubbled through for 2.5 min. When separation is complete, the aqueous layer is pipetted into a suitable tube.

Chromatographic procedure. From 0.1 to 0.6 ml. of the final extract, depending on the expected keto acid content, is then placed on a strip of Whatman no. 3 MM filter paper from a micrometer syringe fitted with a stainless-steel hypodermic needle having a plastic mount (Ballbrook Surgical Instrument Co. Ltd.). This type of needle is not attacked by ammonia. For urine 0.2 ml. is usually ample, but for blood and cerebrospinal fluid 0.4 and 0.6 ml. respectively are taken because of the low α -oxoglutarate levels. Much time and trouble are saved by placing two spots on the strip simultaneously from two Agla syringes, a single micrometer actuated by a suitably geared motor being used, which stops automatically when the required volume has been delivered. At the same time air from a hair dryer at a temperature not exceeding 40° is directed on to the spots. The paper is hung in a sealed-glass tank, and the solvent is allowed to ascend for about 16 hr. The solvent mixture used by El Hawary & Thompson (1953), n -butanol-ethanol-0.5 N - NH_3 soln. (7:1:2), gives a wider separation of the spots between α -oxoglutarate and pyruvate, but this is at the expense of any hydrazones with a lower R_F value than α -oxoglutarate. It is thought that the solvent with the higher ammonia content used in the present work is more generally useful.

Identification. The paper strips are dried and the areas occupied by the hydrazones outlined with pencil, preferably under ultraviolet light. Although the sequence of the hydrazones on the paper is the same whether run in solvent containing 0.5 or N - NH_3 soln., the R_F values differ; α -oxoglutarate hydrazone has an R_F value, when run in the N - NH_3 soln. solvent, of approx. 0.28, the main pyruvate spot about 0.5 and the other pyruvate spot about 0.67. The spot of the hydrazone of acetoacetate, R_F about 0.57, is not as compact as those of pyruvate and α -oxoglutarate, but, although the upper border is rather indefinite, the lower edge is usually well defined. Quantitative measurement of acetoacetate has not been attempted since the proportion of the acid finally appearing as a hydrazone spot is very low. Thus 10 mg. of acetoacetic acid/100 ml. gives the same final colour intensity as 0.22 mg. of pyruvic acid/100 ml. Moreover, since the acetoacetate hydrazone spot is separated from that of pyruvate, small amounts of acetoacetate do not interfere with the quantitative determination of pyruvate. If, however, large amounts of this acid might be expected to be present, it is probably wise to destroy the acetoacetate hydrazone either by conversion into the fast-running acetone hydrazone (R_F 0.95) by heating the dried spot on the paper before chromatography in an oven at 60° for 20 min., or by adding 1 ml. of a 2% (w/v) solution of p -dimethylaminobenzaldehyde in 1.3 N -HCl 3 min. before the first ethyl acetate extraction. Preliminary observations have shown that the first procedure destroys about 90% of the acetoacetic acid hydrazone but leaves the α -oxoglutarate

hydrazone unaffected and only causes a loss of about 5% in the pyruvate hydrazone; with the second procedure 3 mg. amounts of acetoacetate added to blood filtrates are completely removed, but there is a rather larger loss of pyruvate (17%) and α -oxoglutarate (9%) hydrazones. Introduced into the Friedemann & Haugen method when ketonaemic blood is analysed, the second procedure reduces the increment in pyruvate estimation resulting from an acetoacetate concentration of 100 mg./100 ml. to 0.02 mg./100 ml. However, in these circumstances determinations of standard pyruvate and α -oxoglutarate solutions show a constant loss of 13 and 8% respectively. Allowance for this loss of pyruvate must therefore be made in the calculation.

The chromatograms of the more concentrated urines may show small amounts of unidentified substances absorbing ultraviolet light; these do not interfere with the identification of the pyruvate or α -oxoglutarate hydrazones. One of these spots has an R_F value about 0.31 and another about 0.37.

Elution and quantitative estimation. The area of paper occupied by the given hydrazone is cut into small pieces and placed in 5 ml. of the Na_2CO_3 - $NaOH$ mixture and the paper is ground with a glass rod until it has disintegrated. The mixture is then poured into a sintered-glass micro funnel (porosity 3), into which a small disk of filter paper (Whatman no. 42) has previously been inserted, and filtered under pressure into a small test tube. The coloured solution is refiltered through the same funnel. A solution blank is similarly filtered. Fifteen minutes after the addition of the alkali to the paper the red colour is measured in a 2 cm. cuvette with an Ilford Spectrum Violet 601 filter, and compared with a standard curve derived by eluting known amounts of the corresponding keto acid hydrazones from paper without subjecting them to chromatography. The optical density of the solution blank is also determined and a small correction added for the paper blank. For pyruvate spots this is 0.003 and for the smaller α -oxoglutarate spots 0.002. These values are the means of many estimations obtained by eluting, in the absence of added hydrazones, paper that has been treated overnight in the solvent mixture.

Colorimetric estimation of keto acids. The method of Friedemann & Haugen (1943) was also used for estimating the keto acids in blood. Ethyl acetate was used as solvent.

RESULTS

The extraction and chromatographic procedures result in some loss of α -keto acid hydrazones. The loss is greater for pyruvate than for α -oxoglutarate. The extent and constancy of the loss has been determined by comparing the final colour intensities given by known amounts of pyruvate and α -oxoglutarate with that of comparable amounts of synthetic hydrazones spotted on paper and immediately eluted. In thirty-five determinations of solutions containing 24–60 μ g. of pyruvate/16 ml. the mean recovery was 64.3% (s.d. 2.81). The corresponding value for α -oxoglutarate in thirty-five determinations of solutions containing 20–60 μ g./16 ml. was 80.9% (s.d. 2.04). Part of this loss is due to the slight increase in the volume of the NH_3 soln. phase, caused by the solubility of ethyl

acetate in the aqueous phase. Thus the 1.5 ml. of NH_3 soln. originally added increased to 1.6 ml. or over after mixing of the phases. This increase in volume is not allowed for in the calculation and leads to an apparent loss of hydrazone. A further source of loss was the incomplete (88 %) extraction of the hydrazones into the relatively small quantity of NH_3 soln. used; larger volumes of NH_3 soln. have, however, not given sufficiently greater extraction to offset the resulting decrease in concentration.

The mean values for recovery given above have been used in calculating all pyruvate and α -oxoglutarate values. The accuracy of the method in this series of thirty-five estimations is given in Table 1. Results for the recovery of 30 μg . amounts of pyruvate and α -oxoglutarate added to the deproteinized filtrates of blood, or to diluted urine, are given in Table 2.

As a further check on the method a comparison has been made, in 100 determinations on metaphosphoric acid filtrates of human blood, between the α -keto acid content as estimated by the Friede-

mann & Haugen method and the combined pyruvate and α -oxoglutarate content (expressed as pyruvate), determined by the chromatographic technique. The results are set out in Table 3. Good agreement was found over a wide range of pyruvate values, in specimens taken after fasting, after strenuous exercise, or after the ingestion of glucose both in normal subjects and in diseased states. The mean level of α -oxoglutarate in this series was 0.17 mg./100 g. (S.D. 0.06). In general the chromatographic values tended to be slightly lower than the values obtained by the Friedemann & Haugen method. This difference is certainly well within the limits of error of the methods, but it may none the less represent a real difference, since traces of other α -keto acids are present in blood.

With the present method a faint spot (R_f about 0.75), which is not given by standard solutions, is constantly obtained from blood, cerebrospinal fluid and urine. In blood, the substance giving this spot seems to be present entirely or almost entirely in plasma. One other faint spot (R_f about 0.83) has been rarely observed. These two spots probably correspond with the similar faint spots described by Kulonen, Carpen & Ruokolainen (1952) and by Gey (1954).

Acetoacetate has been detected occasionally in blood taken from non-diabetic subjects in the post-absorptive state, but only in very small amounts. Its presence may have occasionally contributed to the slightly higher blood α -keto acid levels found by the Friedemann & Haugen method, but it is more probable that the unidentified hydrazone of R_f 0.75 is responsible, since it is present in amounts which, on elution, give a colour intensity comparable with that given by α -oxoglutarate.

Table 1. *Analysis of solutions containing pure pyruvic and α -oxoglutaric acids*

No. of determinations	Present ($\mu\text{g.}$)	Found ($\mu\text{g.}$)		
		Mean	Range	s.d.
Pyruvate				
15	24	23.9	22.2-26.5	1.22
15	40	39.6	37.1-40.7	1.33
5	60	62.5	60.7-63.5	—
α -Oxoglutarate				
15	20	19.9	18.7-22.0	0.58
15	40	40.1	38.2-42.1	1.07
5	60	60.1	59.1-61.2	—

Table 2. *Recoveries of 30 μg . amounts of pure pyruvic or α -oxoglutaric acids added to 16 ml. amounts of metaphosphoric acid filtrates of blood or of diluted urine*

	No. of determinations	Content before additions ($\mu\text{g.}$)	Recovery	
			Mean	S.D.
		Blood filtrates		
Pyruvate	12	14.9-20.8	93.8	5.7
α -Oxoglutarate	12	3.9-4.2	98.6	4.1
		Urines		
Pyruvate	9	16.8-21.5	93.2	4.2
α -Oxoglutarate	9	24.8-30.2	109.6	6.5

Table 3. *Comparison between the α -keto acid content of 100 consecutive samples of human blood as determined by the Friedemann & Haugen method and the combined pyruvate and α -oxoglutarate content estimated chromatographically*

Results are expressed as mg. of pyruvate/100 g.

	Mean	Range	S.D.
A. Friedemann & Haugen method	1.01	0.42-3.31	0.528
B. Chromatographic method	0.97	0.33-3.34	0.542
C. Differences B - A	-0.04	+0.27 to -0.35	0.102

Table 4. *Pyruvate and α -oxoglutarate contents of blood from resting healthy subjects in the post-absorptive state, and of cerebrospinal fluid from patients with neurological disease*

The number of subjects is given in parentheses

	Pyruvate (mg./100 g.)			α -Oxoglutarate (mg./100 g.)	
	Mean	Range	S.D.	Mean	Range
Blood (8)	0.56	0.36-0.81	—	0.12	0.08-0.18
Cerebrospinal fluid (71)	0.96	0.49-1.69	0.25	<0.02	tr.-0.06

Table 5. *Excretion rates of pyruvate and α -oxoglutarate in the urine of healthy subjects*

Collection period	Subjects		Pyruvate (mg./hr.)			α -Oxoglutarate (mg./hr.)		
	No.	Sex	Mean	Range	S.D.	Mean	Range	S.D.
Overnight	25	M.	0.37	0.14-0.64	0.09	0.92	0.27-1.70	0.38
Overnight	25	F.	0.33	0.23-0.50	0.07	1.60	0.68-3.18	0.55
24 hr.	6	M.	0.42	0.24-0.87	—	1.05	0.45-1.46	—

The values given in Table 3 are for blood obtained under diverse circumstances, and can only be compared therefore with other published values for human blood in a general way. To allow a more exact comparison, results are given in Table 4 for post-absorptive human blood taken from healthy subjects who had been lying on a couch for 30 min. or more. The sum of the mean levels of pyruvate and α -oxoglutarate (expressed as pyruvate) are lower than in the larger series. Values for these acids were also estimated in the cerebrospinal fluid of seventy-one patients with neurological disorders (Table 4). The striking feature is the low level of α -oxoglutarate in the cerebrospinal fluid as compared with that of the blood.

In contrast to blood and cerebrospinal fluid, the α -oxoglutarate content of urine is usually greater than that of pyruvate. In Table 5 the hourly nocturnal excretion rates for these acids have been given; the overnight urine (formed during 7-9 hr. of sleep) was chosen to exclude as far as possible the effects of varying amounts of exercise and of diet. It can be seen that a woman, despite her smaller size, excretes almost twice as much α -oxoglutarate in her urine as a man. To facilitate comparison with other published figures, the results are given of the mean hourly excretion rates of six healthy men over 24 hr. In these six men the mean excretion of pyruvate was 56%, and of α -oxoglutarate 29%, higher during the day than at night.

DISCUSSION

The results obtained show that it is possible with this simplified procedure to estimate the levels of pyruvate and α -oxoglutarate in biological fluids with reasonable accuracy. It is true that there is an appreciable loss, but the loss is constant and some of it due to the increase in volume of NH_3 soln. used

for the extraction of the hydrazones from the ethyl acetate. Incomplete extraction of the hydrazones by the small volume of NH_3 soln. is a further source of loss, but it is probable that, as with the sodium carbonate extraction in the Friedemann & Haugen method, the proportion extracted remains constant. It seems likely also that the greater loss of pyruvate as compared with α -oxoglutarate is due to the conversion of a proportion of the pyruvate hydrazone into the isomer with R_f 0.67, since the colour equivalent of the more rapidly moving isomer after the addition of the $\text{NaOH}-\text{Na}_2\text{CO}_3$ mixture is considerably less than that of the other isomer (Ishewood & Cruickshank, 1954; Markees, 1954, 1955). The relative proportion of the isomer of R_f 0.67 tends to be rather variable, but in the last fifty estimations on blood the proportion present has only been 12.5% (S.D. 5.15).

The values for pyruvate and α -oxoglutarate in human blood correspond fairly closely with the mean levels of pyruvate (0.35-0.92 mg./100 ml.) and of α -oxoglutarate (0.095-0.16 mg./100 ml.) found by other workers who have used chromatographic methods (Seligson, McCormick & Soborov, 1952; El Hawary & Thompson, 1953; Henneman, Altschule, Goncz & Alexander, 1954; Smith & Taylor, 1955). The α -keto acids do not appear to have been examined chromatographically in the cerebrospinal fluid before, but since pyruvate is the only one present in significant amounts, it is possible to make a reliable comparison with results obtained by colorimetric methods. The mean level of 0.96 mg./100 ml. found in the present series is in good agreement with comparable groups of patients reported by Bueding, Wortis & Stein (1942), and Amatuzio & Nesbitt (1950).

Chromatographic separation of the different α -keto acids is particularly necessary in urine, since in human urine α -oxoglutarate tends to be present

in greater amount than pyruvate, and other α -keto acids are also present in small amounts. Kulonen *et al.* (1952) reported a mean daily excretion in five subjects of 16.9 mg. of pyruvate and 14.0 mg. of α -oxoglutarate. The level of activity of the subjects is not mentioned, but it seems likely that the high pyruvate excretion may have been the result of exercise, since it has been shown by Johnson & Edwards (1937) and by McArdle (unpublished work) that the excretion of pyruvate is greatly increased after strenuous exertion. A point of some interest that is being further investigated is the higher excretion rate of α -oxoglutarate in women.

SUMMARY

1. A simplified procedure is described for the separation of the 2:4-dinitrophenylhydrazones of α -keto acids by paper chromatography, and for the quantitative estimation of the separated hydrazones of pyruvate and α -oxoglutarate.

2. Values are given for pyruvate and α -oxoglutarate in post-absorptive blood from healthy resting human subjects, in cerebrospinal fluid and in urine.

3. The rate of excretion of α -oxoglutarate in urine is greater in women than in men.

This work was carried out on behalf of the Medical Research Council. I wish to thank Professor R. H. S.

Thompson and Dr M. F. S. El Hawary for help and advice. I am greatly indebted to Miss H. Pels and Miss F. Crew for their skilled assistance.

REFERENCES

- Amatuzio, D. S. & Nesbitt, S. (1950). *J. clin. Invest.* **29**, 1486.
 Bueding, E., Wortis, H. & Stein, M. (1942). *J. clin. Invest.* **21**, 85.
 Cavallini, D., Frontali, N. & Toschi, G. (1949*a*). *Nature, Lond.*, **163**, 568.
 Cavallini, D., Frontali, N. & Toschi, G. (1949*b*). *Nature, Lond.*, **164**, 792.
 El Hawary, M. F. S. & Thompson, R. H. S. (1953). *Biochem. J.* **53**, 340.
 Friedemann, T. E. & Haugen, G. E. (1943). *J. biol. Chem.* **147**, 415.
 Gey, K. F. (1954). *Hoppe-Seyl. Z.* **294**, 128.
 Henneman, D. H., Altschule, M. D., Goncz, R. M. & Alexander, L. (1954). *Arch. Neurol. Psychiat., Chicago*, **72**, 688.
 Isherwood, F. A. & Cruickshank, D. H. (1954). *Nature, Lond.*, **173**, 121.
 Johnson, R. E. & Edwards, H. T. (1937). *J. biol. Chem.* **118**, 427.
 Kulonen, E., Carpen, E. & Ruokolainen, T. (1952). *Scand. J. clin. Lab. Invest.* **4**, 189.
 Markees, S. (1954). *Biochem. J.* **56**, 703.
 Markees, S. (1955). *Experientia*, **11**, 205.
 Seligson, D., McCormick, G. J. & Soborov, V. (1952). *J. clin. Invest.* **31**, 661.
 Smith, M. J. H. & Taylor, K. W. (1955). *Lancet*, **1**, 27.

Studies on the Particulate Components of Rat Mammary Gland

1. A METHOD FOR DETERMINING THE COMPOSITION OF THE RETAINED FLUID

BY A. L. GREENBAUM AND T. F. SLATER
Department of Biochemistry, University College, London

(Received 15 November 1956)

The mammary gland of the rat as dissected from the animal is a complex system containing not only mammary tissue itself but also the secreted fluid retained in the alveoli and duct system of the gland (colostrum in pregnancy, milk during lactation). Tissue suspensions of rat mammary gland prepared at different stages of the lactation cycle contain a variable quantity of this retained fluid which, in the later stages of lactation, may account for up to 60 % of the gross weight of the gland (Folley & Greenbaum, 1947). As the retained fluid has neither the same chemical constitution nor the same enzymic activity as the parent mammary tissue it is essential, in quantitative studies on mammary tissue, to correct analyses on the total gland for the amount of

retained fluid found in the wet tissue when it is excised from the body.

A method has been published for the estimation of retained milk in suspensions of rat mammary tissue (Folley & Greenbaum, 1947) in which the percentage of lactose in milk obtained from lactating rats was determined and then used to calculate the percentage milk retained in tissue suspensions at all stages of the lactation cycle. While this method gave reasonable values for milk retention during lactation, where the percentage of lactose in milk was shown to be approximately constant, it was almost certainly in error when applied to rats at other stages of the lactation cycle such as late pregnancy, shortly after parturition or early in