We thank Dr W. R. Rees and Mr T. R. Reynolds for information about the paper-chromatographic separation of glucose and sorbitol, and Dr B. J. Bines for assistance in part of the work. This work was made possible by grants from the Agricultural Research Council, to whom we express our thanks.

REFERENCES

- Bacon, J. S. D. & Edelman, J. (1951). *Biochem. J.* **48**, 114.
 Cifonelli, J. A. & Smith, F. (1954). *Analyt. Chem.* **26**, 1132.
 Hanes, C. S. & Cattle, M. (1938). *Proc. Roy. Soc. B*, **125**, 387.
 Pazur, J. H. (1955). *J. Amer. chem. Soc.* **77**, 1015.
 Pazur, J. H. & Budovich, T. (1956). *J. biol. Chem.* **220**, 25.
 Peat, S., Whelan, W. J. & Hinson, K. A. (1952). *Nature, Lond.*, **170**, 1056.
 Peat, S., Whelan, W. J. & Hinson, K. A. (1955). *Chem. & Ind.* p. 385.
 Peat, S., Whelan, W. J. & Hobson, P. N. (1951). *J. chem. Soc.* p. 1451.
 Peat, S., Whelan, W. J. & Jones, G. (1957). *J. chem. Soc.* p. 2490.
 Peat, S., Whelan, W. J. & Kroll, G. W. F. (1956). *J. chem. Soc.* p. 53.
 Peat, S., Whelan, W. J. & Rees, W. R. (1956). *J. chem. Soc.* p. 44.
 Pirt, S. J. & Whelan, W. J. (1951). *J. Sci. Fd Agric.* **2**, 224.
 Somogyi, M. (1945). *J. biol. Chem.* **160**, 61.
 Trevelyan, W. E., Procter, D. P. & Harrison, J. S. (1950). *Nature, Lond.*, **166**, 444.
 Whelan, W. J. & Morgan, K. (1955). *Chem. & Ind.* p. 1449.
 Woolf, L. I. (1953). *Nature, Lond.*, **171**, 841.
 Zill, L. P., Khym, J. X. & Cheniae, G. M. (1953). *J. Amer. chem. Soc.* **75**, 1339.

Toxicity of Autoxidized Squalene and Linoleic Acid, and of Simpler Peroxides, in Relation to Toxicity of Radiation

By V. J. HORGAN, J. ST L. PHILPOT, BARBARA W. PORTER AND D. B. ROODYN
Radiobiological Research Unit, A.E.R.E., Harwell, Didcot, Berks

(Received 8 January 1957)

Dam & Granados (1945) found that in rats deficient in vitamin E there was an increase in organic peroxide-like material (hereinafter abbreviated to 'organic peroxide'). They suggested that it resulted from the well-known chain reaction by which unsaturated fats, rubber, etc. 'autoxidize' in air to form peroxides. Though this interpretation is not the only possible one it was the first plausible indication that autoxidation could occur *in vivo* as well as *in vitro*, and was consistent with the known 'antioxidant' properties of vitamin E (for reviews see Mattill, 1947; Holman, 1954). According to the accepted mechanism chain autoxidation is initiated by free radicals having a high content of chemical free energy, and may be terminated by formation of low-energy free radicals such as semiquinones, as happens in the presence of vitamin E and other antioxidants. In absence of antioxidants one high-energy free radical can give rise to several hundred molecules of organic peroxide before the chain is terminated by mutual reaction of two free radicals. In many chemical systems the ions produced by 'ionizing radiations' (X-rays, etc.) give rise to high-energy free radicals, e.g. OH from water (Weiss, 1944). Thus X-rays would be expected to induce autoxidation, and Mead (1952) has found *in vitro* that they do, though the conditions seem to be critical because V. J. Horgan & J. St L. Philpot (unpublished work) failed to repeat the experiment. The

question whether X-rays induce autoxidation *in vivo* as well as *in vitro* was provisionally answered in the affirmative by Horgan & Philpot (1953, 1954*a*, *b*), who extracted mice that had received a lethal dose (950 r.) of X-rays. With various methods of peroxide estimation, they sometimes found a significant increase ($P < 0.01$; apparent mean chain length about 80) as compared with unirradiated mice. The technique was, however, unreliable and is still being developed. Dubouloz & Dumas (1954) found that in rat skin, 0.1 mm. thick, 5000 r. of X-rays produced 2×10^{-8} and 20 000 r. produced 1.3×10^{-8} mole/cm.² of peroxide (for references from 1950 onwards see Dubouloz & Dumas, 1954). These figures correspond to mean chain lengths of about 14 and 23 respectively; but it is not clear what fraction of this peroxide was in the living part of the skin. Thus there is some evidence that X-rays can initiate autoxidation *in vivo*, though the observed peroxide could be formed in some other way.

There are not many substances in living matter with a structure suitable for chain autoxidation. The most suitable are the 'methylene-interrupted' [m , ($m + 3n$) diene and polyene] fatty acids, linoleic, linolenic and arachidonic, i.e. the essential fatty acids. Other possibilities are the monoenes, notably oleic acid, the carotenoids [conjugated, i.e. m , ($m + 2n$) polyenes] and squalene [m , ($m + 4n$) polyene]. Though all these substances might be

invoked to explain the observations, Horgan & Philpot tried to narrow the field on the basis of an unexpected observation that roughly half the peroxide in their extracts seemed to be less reactive than typical hydroperoxides and behaved more like disubstituted peroxides. This suggested that the substance predominantly concerned might be squalene, whose primary autoxidation product was shown by Bolland & Hughes (1949) to contain one hydroperoxide and one disubstituted (cyclic) peroxide group, as opposed to the primary product from oleic or linoleic acids, which has only a hydroperoxide group (Farmer & Sutton, 1943; Swift, Dollear & O'Connor, 1946; Bolland & Koch, 1945). With linoleic acid, however, the double bonds are shifted into the conjugated position during autoxidation and can then apparently add a second molecule of oxygen to form a cyclic or polymeric peroxide group (cf. Bergström, Blomstrand & Laurell, 1950). The work of Bergström *et al.* was not known to Horgan & Philpot, and as they could find no data concerning a second molecule of oxygen they ignored the possibility. The high toxicity of radiation might be explained if squalene diperoxide, formed by autoxidation, could act as an antimetabolite in blocking cholesterol synthesis, for which squalene was shown to be an intermediate by Langdon & Bloch (1953). If a specific blocking mechanism like this existed squalene diperoxide might be far more toxic than ordinary peroxides. The testing of this possibility was the original purpose of the work described below.

At the same time, apart from the comparison with squalene, data were needed on the toxicity of injected linoleic acid hydroperoxide and peroxides in general. It is already known (Wyss, Clark, Haas & Stone, 1948; Dickey, Cleland & Lotz, 1949; Ross, 1950; Loveless, 1951) that peroxides are mutagenic; but little is known as to their toxicity on injection. Apart from toxicity the autoxidation of essential fatty acids, if extensive, might cause them to be depleted (cf. Sinclair, 1956). In support of this view Cheng, Kryder, Bergquist & Deuel (1952), following earlier work of Decker, Fillerup & Mead (1950), showed that the survival time of chronically X-irradiated rats was less on a fat-free diet than on the same plus methyl linoleate. If radiation toxicity is due to autoxidation of essential fatty acids, giving either toxic products or deficiency, it is surprising, though possibly explicable, that vitamin E fails to protect (Herve & Bacq, 1949; Horgan & Philpot, 1954*b*; Haley, McCulloch & McCormick, 1954).

Our toxicity tests on peroxides of squalene and linoleic acid required a knowledge of their concentration, since we did not separate them from the parent substances. The occasion was used to study the behaviour of autoxidized pure linoleic acid towards the latest form of our reagent, and to

compare it directly with autoxidized squalene. Previously we had available only some doubtfully pure ethyl oleate.

EXPERIMENTAL

Estimation of peroxides. The method of peroxide estimation used here has been developed primarily for butanol extracts of mice, where oxygen must be rigorously excluded to avoid possible interference by aerobic catalysts such as parahaematin, which might be present in the extracts. When strictly anaerobic, most reagents fail to react even with peroxide; but leuco-brilliant cresyl blue in a suitable buffer is an exception. Under our present conditions it can distinguish two types of reactivity according to whether a cuprous salt is needed ('type 2') or not ('type 1'). Further differentiation can be made according to the rate of reaction, and also according to the rate of destruction of peroxide by the buffer in absence of cuprous salt. This classification is purely empirical and has no simple relation to structure or to the division into 'reactive' and 'less reactive' made on a temperature basis by our earlier reagents; but it is nevertheless useful analytically.

The procedure is as follows: 1.4 ml. of about 5×10^{-5} M-peroxide in butanol is placed in a 2 ml. stoppered spectrophotometric cell and oxygen-free nitrogen is bubbled through the solution for 4 min. The cell is mounted beside a similar control cell, at the bottom of a small glass hood with an opening at the top allowing additions to be made against a flow of nitrogen. Through the opening 0.5 ml. of saturated brilliant cresyl blue (approx. 8×10^{-4} M) in butanol buffered with 0.5 M-lutidine plus 0.25 M-acetic acid is run into the cell from a burette fitted with a U-tube containing a reducing column of cadmium filings. Any failure of anaerobic conditions is shown by blue replacing the yellow of the leuco-dye in the control cell, but this is rare. The absorption at 620 m μ in this cell and in the test cell after various lengths of time at room temperature is measured in a spectrophotometer after inserting the stoppers through the hood and removing the cells. Under these conditions $\log I_0/I$ for a 5 mm. thickness is 1.0 for every 6.95×10^{-5} mole of reoxidized brilliant cresyl blue. The above procedure measures the extent of the type 1 reaction. For the total reaction the cuprous salt is added before the buffered leuco-dye, and the extent of the type 2 reaction is estimated from the total by difference. Destruction by buffer is tested by adding the cuprous salt at various times after the buffer. Tests with mixtures of known peroxides failed to reveal interaction, so that one type can apparently be estimated in the presence of another.

The cuprous salt (substantially lithium cuprobromide) is prepared by passing 0.02 ml. of 5×10^{-3} M-cupric chloride (hydrated) plus 0.1 M-lithium bromide plus 0.02 M-lithium acetate in butanol through a burette fitted with a reducing column of copper powder.

Materials

Brilliant cresyl blue

This was freed from the red impurity previously described (Horgan & Philpot, 1954*a*), by Soxhlet extraction with *n*-butyl acetate.

2:6-Lutidine

This was distilled before use and stored under nitrogen.

n-Butanol

This was refluxed for an hour over cadmium filings plus lithium bromide to remove peroxide, and distilled. It was used within 24 hr.

Disuccinoyl peroxide

This was kindly provided by Dr Latarjet.

Other peroxides

These were made by approximately standard methods (cf. Tobolsky & Mesrobian, 1954), except *tert*.-butyl hydroperoxide and dibenzoyl peroxide, which were bought. The solid ones were recrystallized and the liquid ones redistilled shortly before use.

Hydrogen peroxide

This was standardized by the use of an iodometric technique (cf. Vogel, 1951).

Squalene

This was a commercial product which we purified by passage through an alumina column. A highly purified specimen kindly provided by Dr Heller behaved similarly on autoxidation but the toxicity of its product has not yet been measured.

Linoleic acid and methyl linoleate

These were highly purified products kindly provided at cost price by the Hormel Foundation (Austin, Minn., U.S.A.).

Autoxidation. The squalene and linoleic acid were originally autoxidized by slowly bubbling air through them without a catalyst at room temperature (about 20°). In later experiments with linoleic acid and methyl linoleate a roller-tube was used for aeration. Some experiments with linoleic acid were done also with a thin film spread on glass, diffusion of air being relied upon. Most experiments were done in a slow stream of air (from the compressed-air line), dried with silica gel and magnesium perchlorate ('anhydron') and filtered through glass wool.

Animals. Three strains of mice were used: CBA, hairless albino and R [a cross between PCT female and (C3H × 101) male]. R mice were all 3–3½ months old and had a mean wt. of 22.7 ± s.e.m. 0.37 g. (37 mice). They were fed on a standard cube ration.

Toxicity determinations. Solutions of the peroxides were injected intraperitoneally into female mice in volumes of 0.1–0.4 ml. Disuccinoyl peroxide was dissolved in water. The others, being insoluble in water, were dissolved in ethyl palmitate or in squalene (for comparison with squalene diperoxide, which was prepared in squalene by autoxidation). The doses of *tert*.-butyl hydroperoxide, autoxidized linoleic acid, methyl linoleate and squalene were determined by peroxide estimation; those of benzoyl peroxide, hydroxycyclohexyl hydroperoxide, tetralin hydroperoxide and disuccinoyl peroxide were determined by weight and by peroxide estimation; and those of di-*tert*.-butyl peroxide by weight alone. The doses were chosen by trial so as to approach the LD₅₀ with as few mice as possible. A number of mice were then injected with doses near the LD₅₀ until it was determined with sufficient accuracy. This procedure sufficed for present purposes instead of a formal probit analysis, because of the sharpness of the dose-response curves. The mice were kept for 30 days after injection and inspected twice daily for the first two days, then daily.

RESULTS

Types of reactivity of peroxides

Table 1 gives the percentage of theoretical colour development with known peroxides under the conditions described above for distinguishing two types of reaction. The figure for hydrogen peroxide assumes that the iodimetric method gives 100% reaction. The other figures assume that the substances were pure. The present form of reagent, applied to extracts of mice, shows the presence of both types as expected from previous results, but we have not studied them properly.

The behaviour of di-*tert*.-butyl peroxide was puzzling and needs further study. The first specimen of this that we made gave a type 2 reaction to the extent of half an equivalent with no destruction by buffer. A mixture with *tert*.-butyl hydroperoxide (type 1) behaved additively, i.e. the total colour was equivalent to the hydroperoxide in the absence of cuprous salt and to this plus half the di-*tert*.-butyl peroxide in the presence of cuprous salt. After storage for a month at 3° the di-*tert*.-butyl peroxide failed to react under any conditions, although the substance is said to be exceptionally stable. A second specimen was prepared and behaved exactly like the first, reacting when fresh but not after storage. A third specimen failed to react even when fresh, although it and the other two all boiled at 111°, which is the accepted b.p. The third specimen had m.p. –37° (accepted –40°). Neither the second nor the third specimen reacted with iodide even when fresh, showing that hydroperoxide was absent. The first specimen was not tested with iodide. The material used for toxicity tests was specimen 2, freshly prepared. Diethyl peroxide has a slow type 2 reactivity. In the presence of cuprous salt it oxidizes the leuco dye with a half-reaction time of about 35 min. at 23°, and if the buffer is added 5 min. before the cuprous salt about 12% of

Table 1. *Leuco-brilliant cresyl blue reagent tested with various peroxides*

Type 1 reaction in the absence, type 2 in the presence, of cuprous salt.

Peroxide	Percentage of theoretical colour development	
	Type 1	Type 2
Disuccinoyl peroxide	100–101	0
1-Hydroxycyclohexyl-1-hydroperoxide	94	—
<i>tert</i> .-Butyl hydroperoxide	97	—
Hydrogen peroxide	91	—
Tetralin hydroperoxide	1–11	92–99
Diethyl peroxide	2	40 (99 in 90 min.)
Di- <i>tert</i> .-butyl peroxide	0	0 or 47 (see text)

the peroxide is destroyed. With tetralin hydroperoxide, destruction by buffer is complete in a few minutes, but nevertheless cuprous salt enables it to oxidize the leuco dye before it is destroyed.

Autoxidation of squalene and linoleic acid

Fig. 1 shows a log plot of the peroxide contents of squalene and linoleic acid during autoxidation at room temperature in a bubbling tube with untreated air from the compressed-air line. The linoleic acid had been imperfectly anaerobic at 0° for a week after opening, which may have shortened the induction period. The straight lines (except for the first squalene point which was too low for accuracy) show that the kinetics are ideally autocatalytic, in accordance with the slow branched-chain reaction mentioned in the introduction, with no complications such as change in length of the fast linear-reaction chains. Each substance has two straight lines, one for type 1 peroxide reaction only and the other for total reaction. The difference in ordinates (0.3, i.e. approx. log 2) shows that the total is apparently almost exactly twice the type 1 in both cases. Thus there seem to be two types of peroxide group present in both cases, in equivalent amounts if they both give equal yields of oxidized dye. Autoxidation was still continuing at the steadily increasing rate when the products were used for toxicity tests. The total molarity of peroxide groups was then 0.92 for linoleic acid and 0.24 for squalene.

Unfortunately for simplicity, we repeated the linoleic acid autoxidation primarily in order to measure the oxygen uptake and thereby to verify that a second molecule of oxygen was combining to form a cyclic or polymeric peroxide group, which would explain the existence of two types of peroxide reaction as with squalene. The oxygen uptake was measured by weighing on a microbalance, with a light shallow-glass dish supported by three thin legs on a second dish, to avoid loss by creeping, as reaction vessel. The result is shown as a linear plot in Fig. 2. Clearly only one molecule of oxygen is taken up, along a smooth S-shaped curve; but more surprising is the behaviour of the peroxide. Through the major part of the oxygen uptake only the type 1 peroxide reaction is perceptible; but near the end there is a sudden replacement of half of the type 1 reaction by its equivalent of type 2, without further uptake of oxygen. This change seemed, qualitatively, to be accompanied by a sudden increase in viscosity. The conjugated diene was estimated from the absorption at 231 m μ , and is plotted in the same figure. In later repetitions with the above reaction vessel (called a 'cake-stand' from its appearance), with a roller-tube and with the original bubbler-tube, the behaviour was roughly like that of Fig. 2, but the sudden loss of type 1

peroxide reaction occurred irregularly at one or more earlier stages, and was not necessarily accompanied by an equivalent increase in type 2 reaction. Duplicate assays never disagreed by more than 10 %, which is small compared with the irregularities in question. In one roller-tube experiment the type 2 reaction was in the opposite direction, i.e. less colour was developed in the presence of cuprous salt than in its absence. This was shown to be due to a substance that could reduce brilliant cresyl blue in the presence of the cuprous salt; thus the measured type 2 reaction is

Fig. 1. Autoxidation of linoleic acid and squalene with air in bubbler-tube (moles of peroxide/mole of starting material). $\times$, Peroxide determined in the absence of Cu^+ ion; $\circ$, peroxide determined in presence of Cu^+ ion.

Fig. 2. Autoxidation of linoleic acid in air in a thin layer, unstirred (moles of peroxide/mole of starting material). $\blacksquare$, Oxygen absorbed (gravimetric); $\times$, peroxide determined in the absence of Cu^+ ion; $\bullet$, peroxide determined in the presence of Cu^+ ion minus peroxide determined in absence of Cu^+ ion; $\blacktriangle$, conjugated diene estimated from absorption at 231 m μ . At 115 hr. the viscosity seemed to increase suddenly.

the net result of two opposing reactions, either of which can predominate. A typical roller-tube experiment where the net type 2 reaction was positive is illustrated in Fig. 3, where in addition to the measurements already described the absorption developed by thiobarbituric acid at $520\text{ m}\mu$ (cf. Bernheim, Bernheim & Wilbur, 1948) is shown. In the butanol solvent used by us thiobarbituric acid reacts with autoxidized linoleic acid but fails to react with the simpler peroxides. The colour is orange if developed in air, or red in nitrogen turning to orange on admitting air. The main absorption peak in nitrogen is at about $540\text{ m}\mu$ and in air at $460\text{ m}\mu$.

Toxicities

Fig. 4 shows dose-mortality curves for R mice and Table 2 shows the estimated LD_{50} for all the peroxides except diethyl peroxide. The latter was originally intended as a general model of disubstituted peroxides, which might help in estimating the importance of the cyclic or polymeric peroxide groups in the diperoxides of linoleic acid and squalene. It was abandoned for fear that its volatility might give misleading results, though two mice given $460\text{ }\mu\text{moles}$ both died. Later we hoped that di-*tert*.-butyl peroxide would be a more suitable model, but it is less toxic despite its lower volatility. The chief points to note in Table 2 are the normal toxicity of autoxidized squalene and methyl linoleate and the high toxicity of autoxidized linoleic acid as compared with the other peroxides. The three strains of mice are about equally sensitive, except that hairless-albino mice seem to be more sensitive than the others to disuccinoyl peroxide.

It was hoped that injected peroxides might give the delayed deaths characteristic of radiation toxicity, but the delay, though perceptible, is much less. Of 275 R mice that died after peroxide treatment 188 died on the first day, 50 on the second and 34 on the third to fifth day. Only three deaths occurred later than this (on the 11th–14th days). Two of these occurred with autoxidized linoleic acid and one with di-*tert*.-butyl peroxide. For comparison, of 177 R mice that died in various radiation experiments after 850–1000r. of whole-body X-irradiation, none died in the first 3 days, 38 died between 4 and 8 days, 121 died between 9 and 16 days and 15 died between 17 and 30 days after treatment.

Table 3 summarizes the results of control tests on solvents and substances related to the peroxides in various ways, and shows that the toxicities in Table 2 are mainly due to the peroxide groups themselves. It is unlikely that the toxicities of the autoxidized samples of squalene and methyl linoleate were due to unautoxidized material present, since injection of unautoxidized materials in doses several times higher than the LD_{50} doses did not cause any deaths. There were some deaths with

higher doses of unautoxidized linoleic acid, acting possibly as an anionic detergent, but Table 3 shows that the acid is far less toxic than its autoxidation product. It is also unlikely that material other than peroxide is responsible for the toxicity of autoxidized linoleic acid. Thus Fig. 4 (a) and Table 2 show that four samples of linoleic acid at different stages of autoxidation (and therefore differing widely in their content of peroxide), unchanged linoleic acid and decomposition products all have the same toxicity on a peroxide basis.

Fig. 3. Autoxidation of linoleic acid in roller-tube in air (moles of peroxide/mole of starting material). ■, Oxygen absorbed (gravimetric); x, peroxide determined in the absence of Cu^+ ion; ○, peroxide determined in the presence of Cu^+ ion; ▲, conjugated diene estimated from absorption at $231\text{ m}\mu$; △, extinctions/ 10^{-5} mole of starting material (1 cm. , $520\text{ m}\mu$) developed by thiobarbituric acid.

Fig. 4. Toxicities of various peroxides. Curves were obtained with R mice. For further details see Table 2. Curve (a) is from four samples of autoxidized linoleic acid; ●, sample I; ○, sample II; x, sample III; ■, sample IIIa (see footnote to Table 2). (i) Autoxidized linoleic acid; (ii) disuccinoyl peroxide; (iii) hydroxycyclohexyl hydroperoxide; (iv) benzoyl peroxide; (v) autoxidized squalene; (vi) autoxidized methyl linoleate; (vii) tetralin hydroperoxide; (viii) *tert*.-butyl hydroperoxide.

Table 2. *Toxicity of peroxides in three strains of mice*

For method see text.					
Compound	Strain of mouse	Solvent	LD ₅₀ Peroxide (μ moles/mouse)	Original autoxidizable substance (μ moles/mouse)	No. of mice used in test
Autoxidized linoleic acid*					
Sample I	R	Ethyl palmitate	<4	<13.0	28
Sample II	R	Ethyl palmitate	6	12.8	35
Sample III	R	Ethyl palmitate	7	10.8	47
Sample IIIa	R	Ethyl palmitate	6-8	17.6-23.5	80
Disuccinoyl peroxide	R	Water	12	—	16
	CBA	Water	10-12	—	20
	Hairless albino	Water	4-6	—	29
1-Hydroxycyclohexyl hydroperoxide	R	Ethyl palmitate	15	—	37
	CBA	Ethyl palmitate	18-20	—	16
	CBA	Squalene	15-16	—	20
Dibenzoyl peroxide	R	Ethyl palmitate	20	—	47
	CBA	Ethyl palmitate	20	—	16
Autoxidized squalene*	R	Squalene	22	162	29
	Hairless albino	Squalene	50	—	10
Autoxidized methyl linoleate*	R	Ethyl palmitate	45-60	69-92	70
Tetralin hydroperoxide	R	Ethyl palmitate	52	—	32
	CBA	Ethyl palmitate	36	—	16
<i>tert.</i> -Butyl hydroperoxide	R	Ethyl palmitate	58	—	24
	CBA	Ethyl palmitate	60-70	—	8
Di- <i>tert.</i> -butyl peroxide†	R	Ethyl palmitate	1080	—	39

* The three samples of linoleic acid used had autoxidized to the extent of 0.31, 0.47 and 0.65 mole of total peroxide/mole of original linoleic acid respectively. Sample IIIa was obtained by allowing sample III to stand in air for a week until its total peroxide had fallen to 0.34 mole of total peroxide/mole of original linoleic acid. The samples of squalene and methyl linoleate were autoxidized to the extent of 0.14 and 0.65 mole of total peroxide/mole of starting material respectively.

† Dose was calculated from the weight of injected material and not from peroxide estimations.

Table 3. *Control tests of toxicities of solvents and compounds related to peroxides*

All experiments were on female R mice. See Table 2 for toxicities of related peroxides.

Compound	Dose (μ moles/mouse)	No. of mice treated	No. of deaths
Linoleic acid	32.5	10	0
	65.0	5	0
	89.5	10	2*
	195	5	1
Methyl linoleate	78	5	0
	156	5	0
Squalene	635	5	0
<i>cyclo</i> Hexanol	20	5	0
<i>tert.</i> -Butanol	80	5	0
Ethyl palmitate (solvent)	0.6 ml.	5	0

* One of these deaths was probably not due to the treatment.

Table 4. *Toxicities of butyl halides on injection into CBA-strain mice*

Compound	LD ₅₀ (μ moles/mouse)	No. of mice used
<i>tert.</i> -Butyl chloride	600-700	15
<i>n</i> -Butyl chloride	100-150	16
<i>tert.</i> -Butyl bromide	250-300	11
<i>n</i> -Butyl bromide	100-200	19

Table 4 contains data on some butyl halides, which were tested for comparison because it was thought that the peroxides might be acting in a manner roughly analogous to that of alkylating agents, i.e. by chemical substitution. Thus *tert.*-butyl hydroperoxide might be comparable in its effects with *tert.*-butyl chloride or bromide. In fact the butyl halides have very low toxicity, including *tert.*-butyl chloride, which reacts like mustard gas by the '*S_N1*' mechanism (Hughes, 1935). This agrees with the usual experience that monofunctional alkylating agents have only slight biological activity (cf. Loveless & Revell, 1949; Loveless, 1951).

DISCUSSION

Four points have emerged during the present work which suggest that there is no need to invoke squalene rather than linoleic acid in attempting to explain radiation toxicity in terms of autoxidation. These are as follows.

(1) The best reagent that we know so far is unable to distinguish one of our samples of autoxidized linoleic acid from autoxidized squalene, at any stage in the autoxidation. Both products give the two types of peroxide reaction to approximately

equal extents, though in other samples of autoxidized linoleic acid the type 1 reaction predominates and the type 2 reaction may even be negative.

(2) Bergström *et al.* (1950) have shown clearly that under some conditions the addition of a second molecule of oxygen to the conjugated double bonds of linoleic acid hydroperoxide can be comparable in speed with the primary reaction. Their peroxide data suggest, however, that under their conditions (aqueous emulsion at pH 9) the diperoxide decomposed as fast as it was formed. Under our acidic anhydrous conditions of autoxidation (*in vitro*) the diperoxide might easily survive till tested, and it should then behave chemically just like squalene diperoxide, in agreement with our first results. Thus the chief difference in the autoxidation of linoleic acid and squalene may be not in the nature of the product but in the kinetics of its formation, since theoretically the two groups appear simultaneously with squalene and successively with linoleic acid. Our later experiments suggest, however, that the type 2 peroxide reaction can be given by linoleic acid which has been autoxidized under conditions where apparently only one molecule of oxygen is taken up. The abruptness and irreproducibility of the formation of type 2-reacting material, the fact that no extra oxygen is taken up when it is formed, and the fact that it seems to be formed from type 1-reacting material, suggest that it may be formed by some non-oxidative chain reaction such as polymerization of the conjugated double bonds to form a rubber-like polymer, still containing the original hydroperoxide groups but with altered reactivity. The origin of the negative type 2 reactivity observed in one experiment is completely obscure. Until the reactions *in vitro* are clearer and more controllable it is impossible to predict what could happen *in vivo*, but it seems likely that all our analyses of mouse extracts will be explicable in terms of autoxidation of essential fatty acids rather than of squalene.

(3) Squalene diperoxide is no more toxic than several simple peroxides, so that the idea that it might act as a specific antimetabolite receives no support.

(4) On a peroxide basis our specimen of autoxidized linoleic acid was more toxic than any of the other peroxides tried, and the average LD₅₀ for the four samples used (0.26 μ mole of peroxide groups/g.) was slightly less than the mean increased peroxide extracted by Horgan & Philpot from lethally irradiated, as compared with unirradiated, mice (0.22 μ mole/g.). Thus without the need to postulate incomplete extraction by butanol or inefficient distribution of injected peroxide from the peritoneum to sensitive sites there is evidence that radiation produces enough peroxide (if derived mainly from linoleic acid) to account for the death

of the animal. However, the evidence is tentative until a thorough study of mouse extracts has been made with the improved reagent. The chief discrepancy in comparing toxicities of peroxides and of radiation is in the time scale, since injected peroxides kill more quickly than radiation. This discrepancy, and the fact that vitamin E fails to protect appreciably against radiation *in vivo*, may perhaps be explicable in terms of a difference in distribution of peroxide. Hard X-rays produce chain-initiating free radicals fairly uniformly throughout the animal, and any unsaturated fats (e.g. in the phosphatides of cell membranes) will be liable to autoxidation except in places where vitamin E is stored (e.g. the fat depots). Injected peroxides may be carried by the blood more efficiently to some sites than to others. Thus equal total doses of injected and radiation-produced peroxide may kill different groups of cells, and the animal may survive for longer when deprived of one group than of the other. For the acutely lethal effects of radiation, the cells concerned are predominantly the haemopoietic and the dividing intestinal ones, which die within 24 hr. With injected peroxide it is not known which cells die first, or whether any die before the animal as a whole.

Thus all the facts known to us are consistent with the hypothesis that radiation toxicity in mammals is due to initiation of chain autoxidation of essential fatty acids producing lethal doses of peroxides in sites not reached by vitamin E.

The fact that autoxidized methyl linoleate is far less toxic than autoxidized linoleic acid was quite unexpected. Possibly the ionized carboxyl group anchors the linoleic acid to some sensitive site such as a cell membrane. We hope soon to measure the toxicity of autoxidized phosphatides containing linoleic acid.

SUMMARY

1. An anaerobic reagent is described, consisting of leuco-brilliant cresyl blue in butanol buffered with lutidine and lutidine acetate, with and without lithium cuprobromide. This shows two types of reaction with various organic peroxides, according to the effect of the cuprobromide, with further differentiation according to kinetics.

2. Both types of reaction were obtained equally with the products of uncatalysed autoxidation of squalene and linoleic acid at room temperature, at all stages observed, but the autoxidation of linoleic acid was highly erratic.

3. The previous finding of two types of peroxide in extracts of X-irradiated mice can now be explained in terms of linoleic acid as well as of squalene.

4. On a basis of total peroxide content, autoxidized linoleic acid, unlike autoxidized squalene, was

more toxic by intraperitoneal injection than any of the simple peroxides with which it was compared. Autoxidized methyl linoleate was less toxic than most of the simple peroxides.

5. The LD_{50} of autoxidized linoleic acid ($0.26 \mu\text{mole}$ of peroxide/g.) was only slightly higher than the mean increase of peroxide previously found in mice after 950r. of X-rays ($0.22 \mu\text{mole/g.}$).

6. Injected peroxides kill more quickly than radiation, but this may be a question of distribution.

7. The evidence is consistent with the view that radiation toxicity is due to initiation of chain autoxidation of essential fatty acids producing lethal doses of peroxides in sites not reached by vitamin E.

We wish to thank Dr John H. Heller for highly purified squalene, Dr R. Laterjet for disuccinoyl peroxide, Mr M. J. Corp for X-radiation, Mr R. T. Beadle and Mr J. W. P. Phelpsstead for efficient technical assistance, and the animal-house staff for care of the animals.

REFERENCES

- Bergström, S., Blomstrand, E. & Laurell, S. (1950). *Acta chem. scand.* **4**, 245.
- Bernheim, F., Bernheim, M. L. C. & Wilbur, K. M. (1948). *J. biol. Chem.* **174**, 257.
- Bolland, J. L. & Hughes, H. (1949). *J. chem. Soc.* p. 492.
- Bolland, J. L. & Koch, H. P. (1945). *J. chem. Soc.* p. 445.
- Cheng, A. L. S., Kryder, G. D., Bergquist, L. & Deuel, H. J. jun. (1952). *J. Nutr.* **48**, 161.
- Dam, H. & Granados, H. (1945). *Acta physiol. scand.* **10**, 162.
- Decker, A. B., Fillerup, D. L. & Mead, J. F. (1950). *J. Nutr.* **41**, 507.
- Dickey, F. H., Cleland, G. H. & Lotz, C. (1949). *Proc. nat. Acad. Sci., Wash.*, **35**, 581.
- Dubouloz, P. & Dumas, J. (1954). *Proc. 1st Int. Photo-biological Congr., Amsterdam.*
- Farmer, E. H. & Sutton, D. A. (1943). *J. chem. Soc.* p. 119.
- Haley, T. H., McCulloch, E. F. & McCormick, W. G. (1954). *Science*, **119**, 126.
- Herve, A. & Bacq, Z. M. (1949). *C.R. Soc. Biol., Paris*, **143**, 1158.
- Holman, R. T. (1954). *Progress in the Chemistry of Fats and other Lipids*, chap. 2. Ed. by Holman, R. T., Lundberg, W. O. & Malkin, T. London: Pergamon Press Ltd.
- Horgan, V. J. & Philpot, J. St L. (1953). *Trans. Faraday Soc.* **49**, 324.
- Horgan, V. J. & Philpot, J. St L. (1954a). *Brit. J. Radiol., N.S.*, **27**, 63.
- Horgan, V. J. & Philpot, J. St L. (1954b). *Radiobiology Symposium, Liège*. London: Butterworth.
- Hughes, E. D. (1935). *J. chem. Soc.* p. 255.
- Langdon, R. G. & Bloch, K. (1953). *J. biol. Chem.* **200**, 129, 135.
- Loveless, A. (1951). *Nature, Lond.*, **167**, 338.
- Loveless, A. & Revell, S. (1949). *Nature, Lond.*, **164**, 938.
- Mattill, H. A. (1947). *Annu. Rev. Biochem.* **16**, 177.
- Mead, J. F. (1952). *Science*, **115**, 470.
- Ross, W. C. J. (1950). *Nature, Lond.*, **165**, 808.
- Sinclair, H. M. (1956). *Lancet*, **1**, 381.
- Swift, C. E., Dollear, F. G. & O'Connor, R. T. (1946). *Oil & Soap*, **23**, 355.
- Tobolsky, A. V. & Mesrobian, R. B. (1954). *Organic Peroxides*, 2nd ed., pp. 2-55. New York: Interscience.
- Vogel, A. G. (1951). *A Text Book of Quantitative Analysis*, 2nd ed., p. 348. London: Longmans, Green and Co.
- Weiss, J. (1944). *Nature, Lond.*, **153**, 748.
- Wyss, O., Clark, J. B., Haas, F. & Stone, W. S. (1948). *J. Bact.* **56**, 51.

The Enzymic Hydrolysis of Adenosine Triphosphate by Liver Mitochondria

I. ACTIVITIES AT DIFFERENT pH VALUES

By D. K. MYERS* AND E. C. SLATER

Laboratory of Physiological Chemistry, University of Amsterdam, Netherlands†

(Received 4 March 1957)

Interest in the hydrolysis of adenosine triphosphate (ATP) by mitochondria was stimulated by the suggestion that it may be due to a reversal and diversion of the reactions which are responsible for the synthesis of ATP during the process of oxidative phosphorylation (Hunter, 1951; Lardy & Wellman, 1953; cf. Lardy & Elvehjem, 1945). When the mito-

chondria are carefully isolated from liver homogenates in isotonic sucrose solution at 0° , they exhibit little or no hydrolytic activity towards ATP (Kielley & Kielley, 1951). The oxidation of a number of substrates by these mitochondria is obligatorily coupled with the phosphorylation of two or three molecules of adenosine diphosphate (ADP) to ATP for each atom of oxygen consumed. The P:O ratio is decreased and the rate of hydrolysis of added ATP is increased when the structure of the mitochondria is damaged, e.g. by exposure to hypotonic solutions,

* Present address: Suffield Experimental Station, Ralston, Alberta, Canada.

† Postal address: Jonas Daniël Meyerplein 3, Amsterdam-C, Netherlands.