which represents 20% of the total myofibrillar protein, is extracted from the myofibril together with myosin by Guba–Straub solution containing ATP. In our hands this fraction represented 3–7% of the whole and it might be supposed that it consists in part of tropomyosin, which in all makes up at least 10–12% of the myofibril. de Villafranca (1956) has further studied this fraction, and although the effect of salt on its viscosity is remarkably like the behaviour of tropomyosin the protein is precipitated at lower ammonium sulphate concentrations; he concludes therefore that it is probably not tropomyosin. If that is so then the protein studied by Villafranca must constitute part of the residue, representing 60–65% of the myofibrillar protein left after extraction of the soluble fraction. This seems rather unlikely, as the results of the analysis of whole muscle (Hasselbach & Schneider, 1951) suggest that this residue must consist almost entirely of myosin.

SUMMARY

1. On exhaustive extraction of rabbit myofibrils with buffers of low ionic strength, pH 7.1–8.3, 25–36% of the total myofibrillar protein passed into solution. This extract contained tropomyosin, an inactive form of actin and an unidentified protein provisionally named component C.

2. Tropomyosin has been shown to account for at least 10–12% of the total myofibrillar protein.

3. The inactive form of actin extracted from myofibrils was electrophoretically identical with F actin inactivated by dialysis against 0.078M-borate buffer, pH 7.1, but not with F actin inactivated by treatment with potassium iodide.

4. When the extract obtained by treating myofibril with solutions which are considered to remove the A band selectively was dialysed against

0.04M-potassium chloride containing 6.7 mM-phosphate, pH 7.2, 3–7% of the total myofibrillar nitrogen remained in solution.

We wish to express our thanks to Mr T. C. Grey for skilled technical assistance and to the Medical Research Council for a research expenses grant (to S.V.P.). We are also indebted to Mr B. R. Slater for help in the determination of the diffusion constant of component B.

REFERENCES

- Bailey, K. (1948). *Biochem. J.* **43**, 271.
 Bailey, K. (1951). *Biochem. J.* **49**, 23.
 Boxler, E. (1954). *J. gen. Physiol.* **38**, 53.
 Corsi, A. & Perry, S. V. (1958). *Biochem. J.* **68**, 12.
 de Villafranca, G. W. (1956). *Arch. Biochem. Biophys.* **61**, 378.
 Finck, H., Holtzer, H. & Marshall, J. M. (1956). *J. biophys. biochem. Cytol.* **2**, Suppl. p. 175.
 Guba, F. & Straub, F. B. (1943). *Studies Inst. med. Chem. Univ. Szeged*, **3**, 46.
 Guba, F. & Szent-Györgyi, A. (1945). *Acta physiol. scand.* **9**, Suppl. 25.
 Hanson, J. & Huxley, H. E. (1953). *Nature, Lond.*, **172**, 530.
 Hasselbach, W. (1953). *Z. Naturf.* **8b**, 449.
 Hasselbach, W. & Schneider, G. (1951). *Biochem. Z.* **321**, 462.
 Hodge, A. J. (1955). *J. biophys. biochem. Cytol.* **1**, 361.
 Jacob, J. J. C. (1947). *Biochem. J.* **41**, 83.
 Krebs, H. A. & Hems, R. (1953). *Biochim. biophys. Acta*, **12**, 172.
 Morales, M. F., Botts, J., Blum, J. J. & Hill, T. L. (1955). *Physiol. Rev.* **35**, 475.
 Perry, S. V. (1953). *Biochem. J.* **55**, 114.
 Perry, S. V. (1955). In *Methods in Enzymology*, vol. 2, p. 582. Ed. by Colowick, S. P. & Kaplan, N. O. New York: Academic Press.
 Perry, S. V. & Grey, T. C. (1956a). *Biochem. J.* **64**, 184.
 Perry, S. V. & Grey, T. C. (1956b). *Biochem. J.* **64**, 5P.
 Szent-Györgyi, A. G., Mazia, D. & Szent-Györgyi, A. (1955). *Biochim. biophys. Acta*, **16**, 339.
 Watanabe, S. (1955). *Arch. Biochem. Biophys.* **54**, 559.

Some Observations on the Localization of Myosin, Actin and Tropomyosin in the Rabbit Myofibril

BY A. CORSI* AND S. V. PERRY

Department of Biochemistry, University of Cambridge

(Received 4 February 1957)

In the previous paper (Perry & Corsi, 1958) it has been shown that at pH 7–8, under conditions of low ionic strength, about 35% of the total protein can be extracted from the isolated myofibril. As myosin was absent from this extract, which consisted mainly of actin and tropomyosin, it seemed likely that a study of the morphological changes involved

* Present address: Istituto di Patologia Generale, Piazza S. Eufemia, 19, Modena, Italy.

during the extraction of myofibrils under these conditions would provide useful information on the localization of actin and tropomyosin on the one hand and myosin on the other. In the communication which follows, the results of such an investigation are described together with those of complementary morphological and chemical studies under conditions in which myosin was the main component to be extracted from the myofibrils.

METHODS

Preparation and extraction of myofibrils. Myofibrils were prepared and extracted by the methods described in the preceding paper (Perry & Corsi, 1958). The stock pyrophosphate solution contained 0.1 M- $\text{Na}_4\text{P}_2\text{O}_7$, 10 mM- KH_2PO_4 , and N-HCl (approx. 83 ml./l.) was added to bring the final pH to 7. Suspensions extracted with pyrophosphate were centrifuged for 15 min. at 14 000 g, and those extracted with 5 mM-acetate pH 4.3 for 60 min. at 105 000 g. The clear supernatants were used for adenosine triphosphatase (ATPase) and actomyosin assays.

Assay of myosin adenosine triphosphatase. The ATPase activity of extracts and suspensions was determined in 0.05 M-2-amino-2-hydroxymethylpropane-1:3-diol (tris)-HCl buffer, pH 7.4, 0.25 M-KCl, 5 mM- CaCl_2 and 5 mM-adenosine triphosphate (ATP). Incubations were carried out at 20° for 5 min. with three different concentrations of enzyme within the linear-response range. In the extraction with 5 mM-acetate buffer, pH 4.3, and with pyrophosphate solutions the whole suspensions and extracts were dialysed at 0° for 2 days against a solution containing 0.5 M-KCl and 0.039 M-borate, pH 7.1, which was changed several times. The percentage of myosin extracted from the myofibril was taken as 100 times the ATPase activity/ml. of the extract divided by the ATPase activity/ml. of the whole suspension.

Actomyosin test. The extract was dialysed for 2 days against a frequently changed solution containing 0.5 M-KCl and 0.04 M-phosphate buffer, pH 7.2. (Efficient dialysis after pyrophosphate extraction was essential to remove the last trace of pyrophosphate, which would dissociate any actomyosin present.) Viscosities were measured at 0° with an Ostwald viscometer (time of flow for water, 60 sec.) in the presence and absence of 0.5 mM-ATP (Bailey & Perry, 1947). A fall in viscosity on the addition of ATP indicated the presence of actomyosin.

Contraction test. Myofibrillar residues, washed free of the extraction medium, were treated with solutions containing 0.025–0.1 M-KCl, 5 mM-ATP and 5 mM- MgCl_2 , with the pH adjusted if necessary to 7 with dilute alkali. Intact myofibrils contracted violently under these conditions. The test was either carried out by irrigating with the appropriate solution, the myofibrils being kept under observation on the microscope slide, or by inducing contraction in a small test tube and then examining the suspension microscopically.

Microscopic technique. All observations were carried out with the Cooke, Troughton and Simms Ltd. phase-contrast microscope and an oil-immersion achromatic objective, N.A. 1.3, and a $\times 15$ eyepiece. A green filter was used for photography.

RESULTS

Characteristics of myofibril preparations

From fresh muscle in 0.025 M-KCl-borate. When prepared in this medium usually about 30 % of the myofibrils showed reversal of striation; on occasions the proportion of reversely striated myofibrils was even higher (Pl. 1: 1–4). In the majority of the myofibrils the I and A bands were readily recognized, the I band was usually relatively much shorter than the A band and in most cases the sarcomere length ranged from 1.9 to 2.3 μ . The H space, defined as an area of lower optical density in the centre of the

A band of myofibrils possessing discrete A and I bands, was usually present and corresponded to one-third or less of the A band. Very occasionally the H space was not visible and a dark line which probably corresponded to the M line was visible in the centre of the A band (Pl. 1: 2). This dark line was never observed when the H space was apparent.

From fresh muscle in ethylenediaminetetra-acetate-KCl. If the KCl concentration of the borate medium used for the preparation of myofibrils was increased to 0.1 M many more of the myofibrils were reversely striated. If 5 mM-ethylenediaminetetra-acetate (EDTA) was added to 0.1 M-KCl and the pH adjusted to 7 (see Perry & Corsi, 1958), with very few exceptions all myofibrils showed distinct I and A bands (Pl. 1: 5–8) and the sarcomere length ranged from 1.7 to 2.5 μ (average length 2.2 μ). Although reversal of striation was only rarely observed there was a tendency for myofibrils prepared in EDTA to aggregate in suspension. Usually the A band was longer than the I, but frequently myofibrils of sarcomere length 2 μ or less, with the I longer than the A band, could be seen (Pl. 1: 6). The H space was apparent in practically all myofibrils and generally occupied one-third or less of the A band. In some cases it represented one-half or more of the A band (Pl. 1: 7, 8) but in these cases there was no obvious increase in length of the I band. In our preparations there was no clear correlation between the width of the H space and the length of the I band on the one hand and the sarcomere length on the other as demanded by the model of Hanson & Huxley (1955).

From glycerol-treated muscle in 0.1 M-KCl. Excellent myofibrils with distinct I and A bands were readily obtained from glycerol-treated muscle which had been stored for 2–3 weeks at -10° . (Pl. 1: 9, 10). If the glycerol-treated muscle was stored for shorter periods some reversal of striation occurred during the preparation, due no doubt to the endogenous ATP which was still present. The A band was usually longer than the I, and the H space was present in practically all myofibrils. On some occasions it was very narrow and only very rarely could not be recognized at all. Usually when the I band was longer than average the H space was likewise wider, but in our preparations there appeared to be exceptions to this rule.

It seems worthy of note, particularly in view of some of the other findings presented here, that in myofibrils isolated from both fresh and glycerol-treated muscle the Z line varied considerably in density and thickness. Usually the appearance was characteristic for a given myofibril (Pl. 1: cf. 5 with 3 or 7). A number of other authors (e.g. Cajal, 1888; Bennett & Porter, 1953) have commented on the variation of appearance of the Z line in fixed and stained preparations.

From the study of a large number of preparations obtained from both fresh and glycerol-treated muscle the A band did not appear to be constant in length. Sarcomeres with discrete I and A bands ranged in length from 1.7 to 3.0 μ and in general the longer the A band the greater the sarcomere length.

Extraction of components other than myosin from the myofibril

Low ionic strength, pH 7.1–8.3. On prolonged extraction of fresh myofibrils with 0.078 M-borate, pH 7.1, a fraction representing about 25 % of the total protein and consisting mainly of tropomyosin and actin was removed (Perry & Corsi, 1958). In Pl. 2: 11 and 12, which illustrate borate-extracted myofibrils, it can be seen that the Z line has disappeared and the I band also appears empty, although sufficient material remains to hold the A bands together along the myofibril. The region which corresponded to the H space in extracted myofibrils appeared less dense and usually wider than the original H space. Borate-extracted myofibrils still contracted on addition of ATP, although slowly, and the contracted material appeared disorganized.

When glycerol-treated myofibrils were extracted with borate the Z line was much more resistant; it usually survived 20–50 days of treatment, although it often appeared somewhat granular in appearance after such periods. The I band of glycerol-treated myofibrils was likewise more resistant to borate extraction than that of fresh myofibrils. In the former, strands of material would be quite clearly distinguished in the I band after prolonged borate extraction (Pl. 2: 13).

More rapid and exhaustive extraction of tropomyosin and actin was achieved with 5 mM-tris buffer, pH 7.7, (Perry & Corsi, 1958). After extraction under these conditions the picture was similar to that obtained with borate, the edges of the A band being the only prominent features of the original myofibril which were apparent (Pl. 2: 14). These edges were thinner in most cases than the original area bounded by the junction of the I and A bands and the edge of the H space. The I band was completely empty and in many cases it appeared to have collapsed to give the appearance illustrated in Pl. 2, 14 and 15. On continued extraction at pH 7.7 and with shorter periods at pH 8.3 the myofibrillar structure disintegrated. Myosin did not pass into solution under these conditions although about 35 % of the total protein was extracted.

When 35 % of the total protein had been extracted at pH 7.7 application of ATP to the washed residue produced a rather slow but obvious contraction, which suggested that some actin was still present in the residue.

Low ionic strength, pH 6.45. When fresh myofibrils were treated with 5 mM-phosphate buffer, pH 6.45, actin, tropomyosin and component C passed into solution, although much less protein was extracted than at higher pH values. The I band appeared to have been selectively extracted, for the Z line disappeared and the general density of the I band was much reduced (Pl. 2: 16–18). On the other hand, little change was apparent in the A band. In some preparations the I bands were so weakened that the myofibril broke up into individual A bands (Pl. 2: 18).

Extraction by procedures removing mainly myosin

Acetate buffer, pH 4.3. Little morphological change could be detected in myofibrils stored for 10 days in 5 mM-acetate buffer at pH 5.2. At pH 4.3 rapid extraction and marked structural changes could be observed within a matter of minutes when extraction was carried out by irrigation on the

Fig. 1. Electrophoresis diagrams of myosin-containing extracts from myofibrils prepared from fresh muscle. Conditions: 0.25 M-KCl, 0.05 M-potassium phosphate buffer, pH 6.8. Preparation numbers refer to those listed in Table 1. (a) Myofibrils (174 FV, EDTA-prepared) dialysed against 5 mM-acetate buffer, pH 6.45, for 18 hr. Extract represented 63.9 % of total protein and contained tropomyosin equal to at least 2 % of total myofibrillar protein; actin present. Boundaries photographed after 720 min. (b) Myofibrils (177 FV, EDTA-prepared) extracted for 20 min. with 2 vol. of pyrophosphate solution. Extract represented 56.6 % of total proteins and contained tropomyosin equal to at least 1 % of total myofibrillar protein; actin present. Boundaries photographed after 900 min.

EXPLANATION OF PLATES 1-3

PLATE 1

1. Myofibrils 157F isolated from fresh muscle in 0.025M-KCl-borate and examined after 2 days. Normal myofibril with I, A, H and Z bands clearly visible and reversely striated myofibril.
2. Preparation as for no. 1. Myofibril with M line and a partly contracted myofibril (sarcomere = 1.5μ) with pronounced Z line and a double-line system (A band?) in the centre of sarcomere.
3. Preparation as for no. 1. Myofibril with thick Z line.
4. Myofibrils 158F isolated from fresh muscle in 0.025M-KCl-borate and examined immediately. Slightly shortened myofibril (sarcomere = 1.6μ) with prominent Z bands and double-band system (A band?) in the centre of sarcomere (cf. no. 2).
5. Myofibrils 174FV isolated in 0.1M-KCl-borate with 5 mM-EDTA and examined immediately. Typical myofibril.
6. Preparation as for no. 5. Myofibril of sarcomere length about 1.8μ with broad I and comparatively narrow A band with H space.
7. Preparation as for no. 5. Myofibril with broad H space, relatively narrow I and faint Z line. Sarcomere length = 2μ .
8. Myofibrils 162FV isolated in 0.1M-KCl-borate-EDTA. Examined immediately. Wide H space and faint Z line.
9. Myofibrils 159G isolated in 0.1M-KCl from glycerol-treated muscle. Dark line in the centre of A probably corresponding to the M line.
10. Myofibrils 154G isolated in 0.1M-KCl from glycerol-treated muscle. Typical example showing A, I, H and Z bands.

PLATE 2

11. Myofibrils 150F isolated in 0.025M-KCl-borate. Extracted for 28 days in 0.078M-borate, pH 7.1. 25% N extracted.
12. Myofibrils 129F isolated in 0.025M-KCl-borate. Extracted for 88 days with 0.078M-borate, pH 7.1. 24% N extracted.
13. Myofibrils 154G isolated in 0.1M-KCl from glycerol-treated muscle. Extracted for 31 days with 0.078M-borate, pH 7.1. Note remnants of I band and Z line.
14. Myofibrils 160F isolated in 0.025M-KCl-borate. Extracted for 123 days with 5 mM-tris buffer, pH 7.7.
15. Myofibrils 169FV isolated in 0.1M-KCl-borate-EDTA. Extracted for 7 days in 5 mM-tris buffer, pH 7.7. 29% N extracted.
16. Myofibrils 168FV isolated in 0.1M-KCl-borate-EDTA. Extracted for 17 days with 5 mM-phosphate buffer, pH 6.45. 11.6% total N extracted.
17. Preparation and extraction as for no. 16.
18. Preparation and extraction as for no. 16. Myofibril breaking up into separate A band fragments.

PLATE 3

- 19a-c. Myofibrils 174FV isolated in 0.1M-KCl-borate-EDTA. Progressive extraction of the same myofibril by irrigation with 5 mM-acetate buffer, pH 4.3. Final stage corresponded to extraction of 60-70% of total protein.
20. Myofibrils 168F isolated in 0.1M-KCl-borate-EDTA. Extracted for 4 days with 5 mM-acetate buffer, pH 4.3. Note granular Z band.
21. Myofibrils 174FV isolated in 0.1M-KCl-borate-EDTA. Extracted for 20 min. with 3 vol. of pyrophosphate, pH 7. This is the appearance on complete extraction with pyrophosphate. Note gaps in the centres of sarcomeres which are longer than the original H space.
22. Preparation and extraction as for no. 21.
23. Intermediate stage of extraction described in no. 21.
24. Myofibrils 178G isolated from glycerol-treated muscle in 0.1M-KCl-borate. Extracted for 2-3 hr. with 2 vol. of pyrophosphate. Note 'ballooning' effect in A band area, which was sometimes observed during pyrophosphate extraction.
25. Preparation and extraction as for no. 24. Final appearance of 20-30% of glycerol-treated myofibrils.
26. Preparation as for no. 24. Extracted with 2.4 vol. of pyrophosphate. Sarcomeres (length = 1.8μ) showing broad gaps.

A. CORSI AND S. V. PERRY—SOME OBSERVATIONS ON THE LOCALIZATION OF MYOSIN, ACTIN AND TROPOMYOSIN IN THE RABBIT MYOFIBRIL

A. CORSI AND S. V. PERRY—SOME OBSERVATIONS ON THE LOCALIZATION OF MYOSIN, ACTIN AND TROPOMYOSIN IN THE RABBIT MYOFIBRIL

A. CORSI AND S. V. PERRY—SOME OBSERVATIONS ON THE LOCALIZATION OF MYOSIN, ACTIN AND TROPOMYOSIN IN THE RABBIT MYOFIBRIL

microscope stage, but the process could be appreciably slowed down by dialysing myofibril suspensions against the buffer. After prolonged extraction 60–70 % of the total N passed into solution. Such extracts contained mainly myosin, together with some actin and tropomyosin. On electrophoresis in 0.25M-KCl and 0.05M-phosphate buffer, pH 6.8, two main peaks which probably corresponded to L-myosin and actomyosin were obtained (Fig. 1a). Under the conditions of the actomyosin test the relative viscosity of the extract studied in Fig. 1a was 3.36; in the presence of ATP it was 3.27. Although it can be concluded that actin is present, at first sight these figures suggest that the amount is small. This is not necessarily so as the ability of myosin to form actomyosin, which can be demonstrated viscometrically, is considerably reduced by exposure of myosin to pH values below 5 (Perry, unpublished work). Likewise, exposure to pH 4.3 considerably reduced the myosin ATPase activity, but preparations were sufficiently active after 2–3 days for satisfactory assays to be carried out. The results obtained are summarized in Table 1 and suggest that when 64–67 % of the protein is in solution all the myosin has been extracted.

When the extraction of individual myofibrils was observed on the microscope stage the first effect of 5 mM-acetate, pH 4.3, was to decrease the density of the A band. This band was eroded away from the centre, a process which was sometimes accompanied by swelling (Pl. 3: 19b, c). At about the same time the Z band appeared much thicker and more dense until it was the most striking feature of the myofibril. Later, in the middle of the A band a gap appeared which was broader than the original H space, with the result that there could be seen in the fragments which remained a rather thick and dense Z line and thin bands at the edges of the I band area.

These bands probably corresponded to the edges of the A band (Pl. 3: 19a–c). On prolonged extraction the area on each side of the Z line tended to become more even in density, although it was obvious that the density of this area appeared higher than that of the original I band (Pl. 3: 20). It may be suggested that the increase in relative darkness of the Z line and the I band area compared with the original unextracted myofibril is an optical effect due to the extraction of the strongly refracting material from the neighbouring A band. This explanation does not seem entirely satisfactory for the apparent increase in thickness of the Z line and it is possible that some translocation of protein may have occurred.

Pyrophosphate, pH 7. When fresh myofibrils were extracted with 2–4 vol. of $\text{Na}_4\text{P}_2\text{O}_7$ solution, pH 7 (see Methods), 56–60 % of the total N rapidly passed into solution and little further N was extracted by more prolonged treatment. On electrophoretic analysis two main components, which probably corresponded to L-myosin and actomyosin, were obtained (Fig. 1b). The presence of actomyosin was confirmed by the drop of viscosity obtained on the addition of ATP to the solution of the extract. After dialysis against the medium used for the actomyosin test the extract studied in Fig. 1b had a relative viscosity of 2.74, which was reduced to 2.23 on addition of ATP. ATPase assay of dialysed whole suspensions and extracts indicated that 95–100 % of the myosin was extracted.

Chemically the fraction extracted by the pyrophosphate solutions was similar with both fresh and glycerol-treated myofibrils, although in the latter there was some suggestion that less actin was extracted. After dialysis of pyrophosphate extracts of glycerol-treated myofibrils against 0.078M-borate

Table 1. Nitrogen and adenosine triphosphatase extracted from rabbit myofibrils by 5 mM-acetate buffer, pH 4.3, and pyrophosphate solutions

In the extraction with pyrophosphate the figures in the second column represent the volumes of solution added to 1 vol of myofibril suspension. Preparations 178G and 179G were made in 0.1M-KCl-borate from glycerol-treated muscle; the others were made from fresh muscle in the same medium, with EDTA present during initial homogenization.

Preparation no.	Extractant	Concn. (mg. N/ml.)	Extraction time	N extracted (%)	ATPase activity ($\mu\text{g. of P/min./ml.}$)	
					Whole	Extract
177FV	Pyrophosphate (2 vol.)	1.30	20 min.	57.8	56.6	60.0
177FV	Pyrophosphate (4 vol.)	0.771	20 min.	56.7	36.8	33.9
178G	Pyrophosphate (2.4 vol.)	1.027	4 hr.	59.4	36.5	38.7
179G	Pyrophosphate (2.4 vol.)	1.096	4 hr.	56.8	45.4	44.0
173FV	0.1M-KCl-borate	4.16	40 hr.	—	389	(Nil)
	5 mM-Acetate, pH 4.3*	—	40 hr.	38	159	100.4
174FV	5 mM-Acetate, pH 4.3	1.157	18 hr.	64	39.6	43.8
			43 hr.	67	24.4	23.4

* Slow extraction by dialysis.

buffer, pH 7.1, to precipitate myosin and actomyosin, tropomyosin could not be detected in the protein remaining in solution, whereas with fresh myofibrils an amount corresponding to at least 2% of the total N was present.

The appearance of myofibrils after complete extraction with pyrophosphate (Pl. 3: 21, 22) was very similar to that obtained after exhaustive extraction with acetate buffer, pH 4.3, as indeed the chemical evidence would indicate.

Nevertheless, important differences were apparent in the form the extraction took. Initially the H space disappeared and a dense line narrower than the original A appeared midway between the Z lines. Later this new band lost intensity so that the space between the Z lines acquired a more even density, apparently greater than that of the original I band. Finally the centre of the sarcomere, representing most of what was originally the A band, was extracted practically completely and the myofibrils appeared very similar to those shown in the photographs published by Hanson & Huxley (1955) after pyrophosphate extraction, and to those obtained on extraction with 5 mM-acetate, pH 4.3 (Pl. 3: 19, 20).

More prolonged extraction of glycerol-treated myofibrils with pyrophosphate solutions was required to obtain the same morphological picture as was obtained with fresh preparations (Pl. 3: 24-26). Even after 12 days some 20-30% of the myofibrils had a thin line midway between the Z lines (Pl. 3: 25). Some myofibrils had also a thin line on each side of the Z, so that, in addition to the latter, three lines could be seen in each sarcomere. A similar pattern was also sometimes observed during the extraction of fresh myofibrils (Pl. 3: 23).

After extraction of myofibrils and subsequent washing, first with pyrophosphate solution and then with 0.1 M-KCl-borate, in some cases it was possible to induce limited contraction by the addition of ATP. The effect was much less striking than that obtained with fresh myofibrils and indicated that myosin was still present in the residue. In view of the results of the ATPase experiments this must represent a small proportion of the whole of the myosin present in the intact myofibril.

DISCUSSION

Consideration of the results of the morphological studies in conjunction with the chemical investigations described in this and the preceding paper suggest that the following conclusions can be drawn about the localization of proteins within the myofibril.

Myosin

When the I band is extracted so that optically dense material cannot be detected by the phase-contrast microscope, conditions which lead to the

collapse of the I band or the breaking away of individual A bands, no myosin ATPase passes into solution. It can therefore be concluded that myosin is not present in the I band in any significant amount. Likewise it can also be concluded that this protein is absent from the Z band as it appears in uncontracted myofibrils, for when the Z band disappears myosin ATPase does not pass into solution. It follows that myosin is localized entirely in the A band and in this respect our results confirm and extend the findings of Hasselbach (1953), Hanson & Huxley (1953) and those more recently obtained by Finck, Holtzer & Marshall (1956), but do not support the views of Szent-Györgyi, Mazia & Szent-Györgyi (1955) who consider that there is a more even distribution of myosin along the myofibril.

It may be of interest to point out that the residue remaining after extraction of actin, tropomyosin and component C, and which represents about 65% of the total protein, probably contains at least 80% of myosin, but at the moment we feel that there is insufficient evidence to decide finally whether the dense edges of the A band, which are the main features of these extracted myofibrils, represent an accumulation of myosin or some other protein such as residual actin and tropomyosin.

Actin and tropomyosin

These proteins are taken together for we have no evidence to suggest that they are independently localized. On the contrary, the chemical evidence can be interpreted to indicate that they are associated *in situ*, for under a wide variety of conditions these proteins were extracted in fairly similar proportions. Both proteins were obtained when the I band was preferentially extracted at pH 6.45, and at higher pH values when they were also extracted from the A band. On exhaustive extraction at higher pH values the amount obtained was about thrice that extracted at pH 6.45. As normally the A band of myofibrils at equilibrium length is about twice as long as the I band these yields would fit in well with the view that actin and tropomyosin are evenly distributed along the sarcomere in the resting myofibril. Irrespective of the validity of this latter argument it can be concluded that actin and tropomyosin occur in both the I and A bands. Our investigations provide no evidence pointing to the absence of actin in the H space; the results suggest that extraction procedures which decrease the density of the H space extract actin. On treatment with pyrophosphate solutions fresh myofibrils were readily extracted to give the structures illustrated in Pl. 3: 20 and 21, which are similar to those described by Hanson & Huxley (1955). In our preparations the gaps produced in the centre of the sarcomere are wider than the H spaces and cannot be considered to represent the

original H spaces. Careful study of the figures published by Hanson & Huxley (1955) suggests that this fact is also true under their conditions of extraction. Furthermore, as analysis of the protein extracted by pyrophosphate solutions clearly indicates the presence of actin, it cannot be concluded from these experiments that the gap seen in pyrophosphate-extracted sarcomeres corresponds to a break in continuity of actin filaments in the intact myofibril. Continuity of the filaments remaining after extraction of myosin has been demonstrated by Hasselbach (1953).

Hasselbach (1953) has presented evidence that the Z band can be dissolved from the myofibril under conditions of higher ionic strength (e.g. at $I=0.6$). Under our conditions of extraction with low-ionic-strength buffers the Z band likewise disappears, and if it passes into solution then the protein arising from it should be found in the soluble fraction. In view of the fact that the amount of component C present can be much reduced by more extensive washing of the myofibril without the disappearance of the Z band, it seems unlikely that component C arises from this structure. Undoubtedly the main components of the soluble fraction are actin and tropomyosin but it is possible that this fraction contains small amounts of other proteins which are not readily detected by electrophoretic methods. At this stage the protein nature of the Z band must remain unresolved.

According to the model of the myofibril proposed by Huxley & Hanson (1954) and Huxley & Niedergeserke (1954) the H space disappears when the sarcomere length falls below 2μ . Such a model of myofibrillar structure is very difficult to reconcile with the appearance of many of the myofibrils isolated from fresh muscle. If the H space is defined as described above, it can readily be recognized in fresh myofibrils of sarcomere length less than 2μ (see Pl. 1: 6). Such myofibrils are not unusual, and the H space is frequently relatively wide (more than half of the A band). This model likewise demands that actin be absent from the H space, but the evidence presented here gives no indication of this; on the contrary it suggests that either actin or

tropomyosin or both are present in the H band region of the myofibril.

SUMMARY

1. Extraction of actin, tropomyosin and component C from isolated rabbit myofibrils was accompanied by a disappearance of the Z line, almost complete disappearance of material from the I band and a fall in density and widening of the H space.

2. Pyrophosphate solutions and 5 mM-acetate buffer, pH 4.3, extracted practically all the myosin together with some actin and tropomyosin. On exhaustive treatment with these solutions the A band was removed from the sarcomere almost completely.

3. In myofibrils isolated from fresh muscle the H space was apparent in sarcomeres of a wide range in lengths. No obvious correlation was apparent between the lengths of the I band and H space.

4. It is concluded that myosin is absent from the I band and Z line and that actin and tropomyosin are present in both I and A bands.

We wish to express our thanks to Mr T. C. Grey for skilled technical assistance and to the Medical Research Council for a research-expenses grant (to S. V. P.).

REFERENCES

- Bailey, K. & Perry, S. V. (1947). *Biochim. biophys. Acta*, **1**, 506.
- Bennett, H. S. & Porter, K. R. (1953). *Amer. J. Anat.* **93**, 61.
- Cajal, S. R. Y. (1888). *Int. Mschr. Anat. Physiol.* **5**, 205, 253.
- Finck, H., Holtzer, H. & Marshall, J. M. (1956). *J. biophys. biochem. Cytol.* **2**, Suppl. p. 175.
- Hanson, J. & Huxley, H. E. (1953). *Nature, Lond.*, **172**, 530.
- Hanson, J. & Huxley, H. E. (1955). *Symp. Soc. exp. Biol.* **9**, 228.
- Hasselbach, W. (1953). *Z. Naturf.* **8b**, 449.
- Huxley, A. F. & Niedergeserke, R. (1954). *Nature, Lond.*, **173**, 971.
- Huxley, H. E. & Hanson, J. (1954). *Nature, Lond.*, **173**, 973.
- Perry, S. V. & Corsi, A. (1958). *Biochem. J.* **68**, 5.
- Szent-Györgyi, A. G., Mazia, D. & Szent-Györgyi, A. (1955). *Biochim. biophys. Acta*, **16**, 339.