anhydrous hydrazine at 25°. This is probably due to reaction between the hydrazine and reactive peptide bonds in the protein as shown by the following calculation with glycylglycine. In this case $k_u = 12.4 \times 10^{-3} \text{ min.}^{-1}$ at 60° and, assuming $E = 17 \text{ kcal. mole}^{-1}$, it is found that, at 20°, 68 % of the glycylglycine would have reacted after 50 hr. whereas after the same time at 2° (the freezing point of hydrazine) only 15 % would have reacted. The reactivity towards hydrazine of this peptide bond in the protein is probably not very different from that in the dipeptide, hence it may be concluded that, in general, some degradation of proteins will occur even at 2° after prolonged contact with anhydrous hydrazine.

SUMMARY

1. The kinetics of the uncatalysed and acid-catalysed hydrazinolysis of five simple peptides was studied by an experimental technique involving hydrazinolysis, treatment with benzaldehyde, dinitrophenylation and paper chromatography.

2. First-order rate constants were obtained for the uncatalysed reaction and also for the catalysed hydrazinolysis in the presence of a large excess of catalyst. In *m*-hydrazine sulphate at 60° the reaction is about five times as fast as in anhydrous hydrazine.

3. On the basis of the rate constants it is possible to make recommendations about the use of the hydrazine method for the determination of the *C*-terminal amino acids of proteins.

4. The energies and entropies of activation calculated for two peptides show that the approximately 1000-fold range of rate constants encountered in the catalysed hydrazinolysis at 60° is due largely to steric effects. This may allow selective hydrazinolysis of peptide bonds in proteins.

I wish to thank Dr M. Lipson for continued interest in this work and Dr E. O. P. Thompson and Dr J. R. McPhee for helpful discussions. It is a pleasure to acknowledge the technical assistance of Mr H. Groll.

REFERENCES

- Akabori, S., Ohno, K., Ikenaka, T., Okada, Y., Hanafusa, H., Haruna, I., Tsugita, A., Sugae, K. & Matsushima, T. (1956). *Bull. chem. Soc. Japan*, **29**, 507.
- Akabori, S., Ohno, K. & Narita, K. (1952). *Bull. chem. Soc. Japan*, **25**, 214.
- Audrieth, L. F. & Ogg, B. K. (1951). *The Chemistry of Hydrazine*, p. 201. New York: John Wiley and Sons, Inc.
- Bradbury, J. H. (1956). *Nature, Lond.*, **178**, 912.
- Bradbury, J. H. (1958). *Biochem. J.* **68**, 482.
- Fraenkel-Conrat, H., Harris, J. I. & Levy, A. L. (1955). In *Methods of Biochemical Analysis*, vol. 2, p. 360. Ed. by Glick, D. New York: Interscience Publishers Inc.
- Harris, J. I., Cole, R. D. & Pon, N. G. (1956). *Biochem. J.* **62**, 154.
- Kay, L. M., Schroeder, W. A., Munger, N. & Burt, N. (1956). *J. Amer. chem. Soc.* **78**, 2430.
- Kroner, T. D., Tabroff, W. & McGarr, J. J. (1955). *J. Amer. chem. Soc.* **77**, 3356.
- Laidler, K. J. (1950). *Chemical Kinetics*, p. 76. New York: McGraw Hill.
- Lawrence, L. & Moore, W. J. (1951). *J. Amer. chem. Soc.* **73**, 3973.
- Levy, A. L. (1954). *Nature, Lond.*, **174**, 126.
- Locker, R. H. (1954). *Biochim. biophys. Acta*, **14**, 533.
- Niu, C. & Fraenkel-Conrat, H. (1955). *J. Amer. chem. Soc.* **77**, 5882.
- Ohno, K. (1953). *J. Biochem., Tokyo*, **40**, 621.
- Ohno, K. (1955). *J. Biochem., Tokyo*, **42**, 615.
- Penneman, R. A. & Audrieth, L. F. (1948). *Analyt. Chem.* **20**, 1058.
- Perrone, J. C. (1951). *Nature, Lond.*, **167**, 513.
- Rees, E. D. & Singer, S. J. (1956). *Arch. Biochem. Biophys.* **63**, 144.
- Sanger, F. (1945). *Biochem. J.* **39**, 507.
- Schlögl, K., Wessely, F. & Wawersich, E. (1954). *Mh. Chem.* **85**, 957.

The Hydrazinolysis of Insulin, Lysozyme, Wool Proteins and Wool

By J. H. BRADBURY

Wool Textile Research Laboratory, C.S.I.R.O., Geelong, Australia

(Received 29 April 1957)

Quantitative studies of the hydrazinolysis of proteins have been made by Ohno (1953, 1954, 1955), Locker (1954) and Niu & Fraenkel-Conrat (1955). Ohno (1953) reported yields of up to 68 % of leucine from lysozyme which corrected to about 100 % after allowance for decomposition of leucine during hydrazinolysis. Locker (1954) obtained only about 25 % yields of isoleucine and phenylalanine from tropomyosin and actin respectively,

and found that some amino acids were completely decomposed by the hydrazine treatment whereas others could be only partially recovered. Niu & Fraenkel-Conrat (1955) confirmed this observation and, in their work on a number of proteins, corrected for the amount of *C*-terminal amino acid decomposed by the hydrazine (usually more than half). They found that the recovery of amino acids from amino acid plus protein was not appreciably

different from that of the amino acid when treated alone. It is possible to minimize the decomposition of amino acids during hydrazinolysis by working under milder conditions in the presence of an acid catalyst (Bradbury, 1956). This should allow all the amino acids, with the exception of asparagine and glutamine, which are removed in the benzaldehyde treatment, to be determined quantitatively as *C*-terminal in proteins. It was to obtain information on this question, particularly for wool, that this investigation was made.

Preliminary investigations with insulin showed that small amounts of some non-*C*-terminal amino acids were observed in the final solution along with the *C*-terminal amino acid alanine. This probably accounts for the original workers in this field, Akabori, Ohno & Narita (1952), identifying glycine in the hydrazinolysate even although it is not a *C*-terminal residue in insulin. Locker (1954) always found small amounts of non-*C*-terminal amino acids, particularly glutamic acid, serine and alanine, which set a practical limit to the applicability of the method since with high-molecular-weight proteins they become comparable in amount with the quantity of *C*-terminal amino acid. On the other hand, Niu & Fraenkel-Conrat (1955) claim that for the 15 proteins which they investigated, small amounts (0.1 equiv./mole) of non-*C*-terminal amino acids were observed only in a few cases. In a recent paper, Akabori *et al.* (1956) consider the possibility of the formation of these non-*C*-terminal amino acids as a result of hydrolysis of hydrazides during and after hydrazinolysis.

In this paper the acid-catalysed hydrazinolysis method, which is useful for simple peptides (Bradbury, 1956, 1958), is applied to insulin, lysozyme, wool proteins and wool itself. A large number of experiments have been made with the first two in order to study the effect of variation of temperature, water content of the hydrazine and other factors on the yield of *C*-terminal and non-*C*-terminal amino acids. The results obtained with wool show the limitations of the method when it is applied to a protein having very few, if any, *C*-terminal groups.

EXPERIMENTAL

Materials

Samples of bovine insulin (Boots Pure Drug Co. Ltd., Nottingham) and lysozyme (Armour and Co., Chicago) were used without further purification. They were shown to be free of loosely bound amino acids or peptides since after dinitrophenylation no ethyl acetate-extractable dinitrophenyl-(DNP)-amino acids or DNP-peptides were detected on paper chromatograms.

A sample of Merino 64's greasy wool was separated into

base and tip and each section extracted in a Soxhlet extractor with light petroleum (Shell X 222) for 8 hr. After drying overnight the wool was immersed six times in water, each for 15 min. at 50°, and air-dried at room temperature.

The samples of wool proteins used were kindly supplied by Mr J. M. Gillespie of the Wool Textile Research Laboratory, Melbourne. The material Z29 was the insoluble residue left after successive extractions of wool with alkaline thioglycollate solution. The soluble material from one such extraction was treated with iodoacetic acid to produce *S*-carboxymethylkeratine 2 (SCMK 2) which was fractionated with $(\text{NH}_4)_2\text{SO}_4$ (Gillespie, 1956, 1957).

Procedure

About 1–2 μ moles (10–100 mg.) of material was weighed into a tube, followed by hydrazine sulphate (0–200 mg.). The material was dried for about 2 hr. at 80° at 10^{-4} mm. Hg pressure and an accurately known volume of hydrazine of known water content (see Bradbury, 1958) was distilled in. The tube was sealed under vacuum, heated and opened, the excess of hydrazine was removed and the residue shaken with benzaldehyde, to remove hydrazides, and dinitrophenylated and extracted as previously described (Bradbury, 1958). The ethyl acetate solution was evaporated to dryness and the bulk of the dinitrophenol removed by two treatments for 30 min. at 40–50° and 10^{-4} mm. Hg pressure, a cold-finger condenser packed with dry ice being used. Test experiments with various DNP-amino acids showed that no appreciable amount of sublimation of these materials occurred under these conditions. The residue was dissolved in acetone and applied as a spot to Whatman no. 3MM paper. The latter is preferred to no. 1 paper because it is more reproducible in behaviour and has a much higher capacity. The paper was run by the ascending method for 16 hr. in the first direction with the toluene solvent and then, after drying at 40° for about 3 hr., in the second direction with the phosphate solvent (Levy, 1954). The spots were marked, those which disappeared with HCl gas were disregarded (including dinitrophenol) and the others tentatively identified by their position with respect to dinitrophenol and a control which was run in the second direction. These spots were cut out, the colour was eluted with 2% NaHCO_3 soln., the optical density of the solution measured and the concentration of material in μ moles/g. of protein calculated from, in general, the extinction coefficients in Table 1 of the preceding paper. In most cases there was a brown-yellow streak due to DNP-hydrazides running almost the whole length of the paper in toluene but not moving at all in the second solvent. The aqueous layer from the extraction procedure was not examined, since only mono-DNP-histidine could be present and bis-DNP-histidine was never detected in the two-dimensional chromatogram.

The identity of the spots was checked by hydrolysis with twice-distilled 6*N*-HCl in a sealed tube at 110° for about 8 hr., and extraction with ethyl acetate. The aqueous layer was evaporated to dryness to remove HCl, and dinitrophenylated and extracted in the usual manner. The DNP-materials from both fractions were then identified by one-dimensional chromatography with the toluene, phosphate or 2-methylbutan-2-ol solvents (Fraenkel-Conrat, Harris & Levy, 1955).

RESULTS

The results are given in Tables 1-3.

DISCUSSION

From Table 1 it is clear that the yield of alanine from insulin is not affected by alteration of the conditions of either the benzaldehyde treatment or

the dinitrophenylation, within the limits recorded. The hydrazinolysis reaction is incomplete in the absence of acid catalyst at 60° but complete in *m*-hydrazine sulphate since no increase of yield occurs on raising the temperature to 80°. As the *C*-terminal sequence of insulin is lysylalanine this result would be predicted from the rate constants in the preceding paper for the uncatalysed and

Table 1. *Hydrazinolysis of insulin and lysozyme*

For definitions of the abbreviations for amino acids used in this table and in Table 3 see *Biochem. J.* (1953), 55, 5.

Material	Expt. no.	Conditions of hydrazinolysis				Yield of <i>C</i> -terminal amino acid* (%)	Non- <i>C</i> -terminal amino acids (μmoles/g. of protein)			
		Water in hydr-azine (% w/w)	<i>m</i> -Hydr-azine sulphate	Temp.	Time of heating (hr.)		Gly	Ser	Glu	Asp
Insulin	161 <i>b</i> , 169 <i>a</i> , 182 <i>a</i> †	0.0	1.0	60°	16	36±4	10±2	7±1	11±1	0
Insulin	244 <i>a</i> , 182 <i>b</i>	0.0	1.0	60	16	37	9	3	10	0
Insulin	161 <i>a</i> ††	0.0	1.0	60	16	31	8	13	5	0
Insulin	244 <i>b</i> §	0.0	1.0	60	16	35	5	4	8	0
Insulin	245	5.0	1.0	60	16	39	6	6	10	0
Insulin	260	35.0	1.0	60	16	21	14	9	15	0
Insulin	144†	0.0	0.0	60	16	16	Not measured			
Insulin	162 <i>b</i> †	0.0	1.0	80	16	39	11	19	13	0
Insulin	162 <i>a</i> ††	0.0	1.0	80	16	34	8	6	14	0
Insulin	145††	0.0	0.0	80	16	38	Not measured			
Insulin (1 μmole)	218	0.0	1.0	60	16	46	5	6	10	0
+ alanine (1 μmole)										
Alanine	201	0.0	1.0	60	16	93	0	0	0	0
Lysozyme	183 <i>a</i>	0.0	1.0	60	16	77	6	2	0	3
Lysozyme	183 <i>b</i> †	0.0	1.0	60	16	54	4	3	0	4
Lysozyme	210	0.0	1.0	60	2	15	1	2	0	1
Lysozyme	195†	0.0	0.0	60	16	64	0	0	0	v. small
Lysozyme	194 <i>a</i>	0.0	1.0	80	16	53	14	5	0	6
Lysozyme	233, 242	0.0	1.0	60	16	82	6	2	0	4
(1 μmole) + leucine (1 μmole)										
Leucine	243	0.0	1.0	60	16	95	v. small	v. small	0	0

* The *C*-terminal alanine is recovered from insulin (mol.wt. 6000) but not *C*-terminal asparagine, which is removed in the benzaldehyde treatment; the *C*-terminal leucine is obtained from lysozyme (mol.wt. 14 900).

† Dinitrophenylated by Levy's (1954) method; all others by Sanger's (1945) method.

‡ Benzaldehyde treatment with A.R. material in neutral solution. In all other cases treatment was in H₂SO₄ soln. (from N₂H₄·H₂SO₄ after reaction of the hydrazine with benzaldehyde at about pH 2).

§ Treatment with peroxide-free benzaldehyde prepared by double distillation of the A.R. material under reduced pressure of nitrogen (Mulcahy & Watt, 1953).

|| Insulin had not dissolved completely by the end of the treatment.

Table 2. *Yield of C-terminal amino acids from insulin and lysozyme*

	Insulin*	Lysozyme
Mean yield (% uncorr.)	36±1 (excluding Expts. 260, 144)	62±6 (excluding Expt. 210)
Overall yield (%) of <i>C</i> -terminal amino acid from protein plus amino acid added in equimolar amount	46 (Expt. 218)	82 (Expts. 233, 242)
Recovery (%) of the amino acid added to the protein†	56	102
Yield (% corr.)	36 × 100/56 = 64	62

* The other *C*-terminal amino acid, asparagine, is converted into the hydrazide and consequently removed by the benzaldehyde treatment.

† This quantity is calculated on the assumption that the yield of, for example, alanine from the insulin in Expt. 218 is 36%, hence the recovery of the alanine added is 2 × 46 - 36 = 56%.

Table 3. *Hydrazinolysis of wool and wool proteins in hydrazine containing m-hydrazine sulphate at 60° for 16 hr.*

Protein	No. of run	Yield of amino acids (μ moles/g. of protein)						
		Gly	Ser	Thr	Glu	Asp	Ala	Total
Wool (base)	216, 211	4 $\pm$ 2	6 $\pm$ 3	3 $\pm$ 1	5 $\pm$ 1	2 $\pm$ 1	4 $\pm$ 2	24 $\pm$ 10
Wool (base)*	217	3	3	3	2	2	0	13
Wool (tips)	262	3	5	4	2	2	0	16
SCMK 2 (Z34F)	281, 287	0	3 $\pm$ 1	0	4 $\pm$ 3	3 $\pm$ 2	0	10 $\pm$ 6
SCMK 2 (Z34F)†	282, 288	7 $\pm$ 3	5 $\pm$ 2	0	2 $\pm$ 2	10 $\pm$ 3	0	24 $\pm$ 10
Z 29	261, 273	0	8	0	0	6	1	15

* No hydrazine sulphate used.

† 5% of water in hydrazine.

catalysed hydrazinolysis at 60° of the similar peptide lysylglycine. However, for lysozyme the C-terminal leucine is liberated completely at 60° in the absence of catalyst (Expt. 195), hence the C-terminal peptide bond must be considerably more labile than that in lysylglycine or leucylglycine (see Fig. 1 of the preceding paper).

Ohno (1955) has proposed the C-terminal sequence Asp.Gly.Ala.Asp(NH₂).Leu [for definitions of these abbreviations see *Biochem. J.* (1953), 55, 5] for lysozyme on the basis of peptides separated after hydrazinolysis, but this result is at variance with a considerable amount of circumstantial evidence which points to Lys.Leu or Arg.Leu (Penasse, Jutisz & Fromageot, 1953; Thureauux & Acher, 1956; E. O. P. Thompson, private communication). Because of the observed lability of the C-terminal peptide bond of lysozyme towards hydrazine a Lys.Leu sequence does not seem likely.

Expts. 201 and 218 show that although alanine itself is not decomposed to any great extent by reaction with hydrazine, about 44% is decomposed in the presence of insulin (Table 2). On the other hand, leucine is not decomposed appreciably by reaction with hydrazine even in the presence of lysozyme. Niu & Fraenkel-Conrat (1955) found no appreciable difference in the recovery of amino acids after uncatalysed hydrazinolysis for 10 hr. at 100° in the presence or absence of proteins, but it must be stressed that their recoveries of alanine and leucine when hydrazinolysed alone were only 45 and 54% respectively. It is not clear why the corrected yield of C-terminal amino acid from insulin and lysozyme is only about 60%.

The formation of small amounts of non-C-terminal amino acids from insulin and lysozyme must be due to side reactions, the most likely being hydrolysis of hydrazides occurring during hydrazinolysis or in the subsequent treatments or in both (Locker, 1954; Akabori *et al.* 1956). The last-named authors measured the amount of hydrolysis of serine, glycine, alanine and leucine hydrazides after various treatments. They found that serine and glycine hydrazides were hydrolysed, to the

extent of about 0.6%, by heating in anhydrous hydrazine (purity > 99%) at 100° for 5 hr., but alanine and leucine hydrazides only to about 0.2%. Dinitrophenylation by Sanger's (1945) method for 2 hr. hydrolysed not more than 0.5% of these hydrazides.

Only a small fraction of the non-C-terminal amino acids can be produced by hydrolysis during dinitrophenylation because of the prior removal of the major part of the hydrazides by the benzaldehyde treatment. Akabori *et al.* (1956) suggest that oxidation of hydrazides to acids may be brought about by peroxides present in the benzaldehyde and used isovaleraldehyde instead. On comparing Expts. 182b and 244a with Expt. 244b in Table 1, it is clear that the use of twice-redistilled benzaldehyde may cause a slight but not significant lowering of the glycine content over that obtained with A.R. benzaldehyde. The only possible explanation of the formation of most of the non-C-terminal amino acids would seem to be that they are produced during hydrazinolysis in anhydrous hydrazine. If this were so one would expect that the amino acids observed should be those which are most readily produced from the hydrazide and, to a lesser extent, those which are most abundant in the protein. Thus the relative abundance of amino acids in insulin is leucine + isoleucine > glutamic acid > valine > glycine, tyrosine > serine, but because of the greater reactivity of the hydrazides of serine and glycine as compared with leucine and alanine (and, by analogy, valine too) the actual order observed is glutamic acid, glycine > serine. With lysozyme the relative abundance of amino acids is aspartic acid > leucine + isoleucine > glycine, arginine, alanine > serine and the non-C-terminal amino acids are glycine > aspartic acid, serine.

The obvious way in which the non-C-terminal amino acids could be produced from the hydrazides is by hydrolysis, but if so one would expect an increase as the water content of the hydrazine is increased. However, as shown in Table 1 for insulin, there is no change with hydrazine containing 5% of water and only a slight increase even with 35% of water present. Perhaps they are

produced by some reaction other than hydrolysis, but in either event it is seen that it has not been possible to eliminate these non-*C*-terminal amino acids even by the careful exclusion of water from the system by the use of high-vacuum techniques. In the measurement of the *C*-terminal amino acids of other proteins it is therefore necessary to make a correction for the formation of these non-*C*-terminal amino acids.

From the results in Table 3 it is clear that there are no more amino acids produced by hydrazinolysis of the wool proteins SCMK 2 and Z29 than from wool itself. The alkaline thioglycollate extraction, followed by treatment with iodoacetic acid (for SCMK 2), has therefore produced no new *C*-terminal amino acids, or new *N*-terminal amino acids (Thompson, 1957). Thus the alkaline thioglycollate treatment solubilizes wool by specific rupture of cystine bonds, there being no attack of peptide bonds.

By the use of the hydrazine method on wool, glycine, alanine, serine and threonine were detected qualitatively by Blackburn & Lee (1954). To these must be added glutamic acid and aspartic acid, as shown in Table 3. However, it is very difficult to make a quantitative estimate of the *C*-terminal amino acids because of the occurrence of non-*C*-terminal amino acids and also the losses of *C*-terminal amino acids on hydrazinolysis. The only way to correct for these two factors is to use the data obtained from insulin and lysozyme, although it is realized that this course is open to some objection. It is seen from Tables 1 and 3 that glycine, serine, glutamic acid and aspartic acid from insulin and lysozyme are obtained in slightly greater amount than from wool. Assuming that these are almost entirely non-*C*-terminal in wool leaves only 5 μ moles of *C*-terminal amino acid/g. of protein; about 10 μ moles/g. of protein after allowance is made for the non-quantitative recovery of *C*-terminal amino acids on hydrazinolysis. The *N*-terminal amino acids of wool are the same as those given in Table 3, with valine additional. Their amount has been estimated by various workers with the method of dinitrophenylation and is summarized by Thompson (1957); a total of 16–20 μ moles/g. seems most likely.

SUMMARY

1. The hydrazine method has been used to determine the *C*-terminal amino acids of insulin, lysozyme, wool proteins and wool.

2. The *C*-terminal amino acids, alanine from insulin and leucine from lysozyme, are obtained in 60% yield, together with small amounts of the non-*C*-terminal amino acids, glycine, serine, glutamic acid and aspartic acid.

3. The non-*C*-terminal amino acids appear to be produced during hydrazinolysis by a process which is independent of the amount of water in the hydrazine, at least up to 5%.

4. The total amount of the *C*-terminal amino acids, glycine, serine, threonine, alanine, glutamic acid and aspartic acid, in wool and the wool proteins studied is about 10 μ moles/g.

The author wishes to thank Dr M. Lipson for continued interest in this work, Dr E. O. P. Thompson for helpful discussions and Mr H. Groll for assistance with the experimental work.

REFERENCES

- Akabori, S., Ohno, K., Ikenaka, T., Okada, Y., Hanafusa, H., Haruna, I., Tsugita, A., Sugae, K. & Matsushima, T. (1956). *Bull. chem. Soc. Japan*, **29**, 507.
- Akabori, S., Ohno, K. & Narita, K. (1952). *Bull. chem. Soc. Japan*, **25**, 214.
- Blackburn, S. & Lee, G. R. (1954). *J. Text. Inst.* **45**, T487.
- Bradbury, J. H. (1956). *Nature, Lond.*, **178**, 912.
- Bradbury, J. H. (1958). *Biochem. J.* **68**, 475.
- Fraenkel-Conrat, H., Harris, J. I. & Levy, A. L. (1955). In *Methods of Biochemical Analysis*, vol. 2, p. 360. Ed. by Glick, D. New York: Interscience Publishers Inc.
- Gillespie, J. M. (1956). *Proceedings of the International Wool Textile Research Conference*, vol. B, p. 34. Melbourne: Commonwealth Scientific and Industrial Research Organization.
- Gillespie, J. M. (1957). *Aust. J. biol. Sci.* **10**, 105.
- Levy, A. L. (1954). *Nature, Lond.*, **174**, 126.
- Locker, R. H. (1954). *Biochim. biophys. Acta*, **14**, 533.
- Mulcahy, M. F. R. & Watt, I. C. (1953). *Proc. Roy. Soc. A*, **216**, 10.
- Niu, C. & Fraenkel-Conrat, H. (1955). *J. Amer. chem. Soc.* **77**, 5882.
- Ohno, K. (1953). *J. Biochem., Tokyo*, **40**, 621.
- Ohno, K. (1954). *J. Biochem., Tokyo*, **41**, 345.
- Ohno, K. (1955). *J. Biochem., Tokyo*, **42**, 615.
- Penasse, L., Jutisz, M. & Fromageot, C. (1953). *Bull. Soc. Chim. biol., Paris*, **35**, 376.
- Sanger, F. (1945). *Biochem. J.* **39**, 507.
- Thaureaux, J. & Acher, R. (1956). *Biochim. biophys. Acta*, **20**, 559.
- Thompson, E. O. P. (1957). *Aust. J. biol. Sci.* **10**, 225.