INHERITANCE IN THE ASEXUAL REPRODUCTION OF *CENTROPYXIS ACULEATA*

F. M. ROOT

Zoölogical Laboratory, Johns Hopkins University, Baltimore, Maryland

[Received June 9, 1917]

TABLE OF CONTENTS

	PAGE
INTRODUCTION	173
The "pure line" concept.....	173
Problems	175
Plan of present investigation	175
Material— <i>Centropyxis aculeata</i>	176
Descriptive	176
Previous breeding work	179
Methods	180
Cultural Technique	180
Statistical methods	180
Inheritance within a population	181
Existence of diverse races	181
Ancestral correlations within a population	183
Inheritance within single clones	185
Measured by the coefficient of correlation	185
Ancestral correlations in line 30.....	185
Ancestral correlations in the other clones	186
Inheritance proved to occur by selection experiments.....	188
Selection experiments 1 and 2; contradictory results of two brief experiments in mass selection	188
Selection experiment 3; quick effect of individual selection based on progeny	191
Selection experiment 4; decided effect of long-continued mass selection in a favorable race	192
CONCLUSIONS	195
Results of the present investigation	195
The present status of the pure line hypothesis.....	196
LITERATURE CITED	198
APPENDIX—Some important correlation tables	200

INTRODUCTION

The "pure line" concept

Since JOHANNSEN (1903), as a result of his studies on self-fertilized plants, first arrived at the conclusion that selection is ineffective within

"pure lines," the phenomenon of complete regression in such material has been shown, by a multitude of other investigators, to be of very general occurrence. It has been proved that the same general state of affairs exists, not only in self-fertilized plants, but also in plants reproducing vegetatively, by tubers or cuttings, and in animals reproducing by parthenogenesis, budding and fission. It is unnecessary to enter into the details of these investigations. I will mention only the papers of EAST (1910) on the potato, of AGAR (1914) and EWING (1914 a, 1914 b, 1916) on daphnids and aphids reproducing parthenogenetically, of HANEL (1908) and LASHLEY (1915, 1916) on Hydra, and of BARBER (1907), JENNINGS (1908) and WINSLOW and WALKER (1909) on protozoans and bacteria. All these investigators entirely corroborated the results and conclusions of JOHANNSEN.

And what are these conclusions? Simply that in organisms reproducing asexually and in homozygous organisms reproducing by self-fertilization, each species is not an ultimate unit, but is itself made up of a large number of diverse strains, called "biotypes," or "clones," differing from each other, but each one constant and invariable in its own hereditary constitution. This last idea, that every individual of any given pure line or clone is precisely the same in its genetic make-up as any other individual of the same pure line or clone, is the basic idea of the "pure line" concept. As a corollary, it follows that all the variations found within a single clone are purely somatic and not heritable. In this case, it is evident, selection working within a clone could have no effect. The testing of this hypothesis, then, is of the greatest importance for any general consideration of the mechanism of evolution.

This problem has already attracted the attention of numerous investigators. Some have completely confirmed the conclusions of JOHANNSEN. The more important of these papers have just been cited. But there are others who have maintained that hereditary variations do occur within clones, and that selection within a clone may have an effect.

PEARSON (1910), subjecting the data of the earlier pure line workers to a more accurate statistical analysis, found evidence indicating that such inheritance might exist. JOHANNSEN himself (1913) showed that sudden heritable variations (mutations) had appeared in several of his pure lines, and that these variations might be either minute or large. STOUT (1915), STOCKING (1915), MIDDLETON (1915), and JENNINGS (1916) have all succeeded in isolating, by selection, hereditarily diverse strains within single clones. The present paper adds another similar case to this list.

Problems

Accepting, as we must, the fact that there do exist, within a single species, diverse strains or races, there are two main problems which claim our attention. The main problem, undoubtedly, is the one which has already been discussed. Are "pure lines" really pure? Do heritable variations occur within the clone? It is this problem which is attacked in the present paper.

A second problem, intimately connected with the first, is presented by the very existence of diverse strains. How did these diverse strains originate? Or, to bring the problem within the horizon of the experimentalist, can new strains arise from old ones, and if so, how? The number of diverse strains which exist in nature is almost beyond belief. In my own work with *Centropyxis*, out of over thirty individuals, isolated from a single pipetteful of debris, taken from a single dead lotus stem, there were no two which, after a month's multiplication and careful study, could be surely assigned to the same clone. Where did these diverse races come from? How long have they been in existence? How do new races arise from existing races? To these questions we have, as yet, no answer.

Plan of the present investigation

The object of my work with *Centropyxis* was to find out whether heritable variations do or do not occur within a single clone. Since many investigators have reported that such variations are not to be found, it may be taken for granted that it will be no easy task to find them, even if they do occur. It is not sufficient simply to pick a character at random, and determine whether its variations are or are not hereditary. One must first show that diverse clones exist in the given species, and that these clones differ from each other in certain inherited characters. Then these characters may be studied, and those which are most favorable for showing the inheritance of variations within the clone, if it exists, may be determined. When this has been done, and not until then, one is in a position to put the question to a fair test.

There are two ways in which the inheritance of variations within the clone may be determined. The first is a statistical method, making use of the coefficient of correlation. The inheritance of diversities within the clone is always accompanied, unless some other factor intervenes, by a significant positive correlation between parents and their offspring. Also, as PEARSON (1910) has pointed out, if the size of the coefficient

of correlation diminishes as we go back to more and more remote ancestral generations, it indicates that inheritance of variations occurs within single clones. This criterion of a "diminishing ancestral correlation" applies to coefficients of correlation computed either from a single clone or from a population containing many clones.

All the more recent investigators have realized, however, that results expressed solely as coefficients of correlation are often inconclusive and hard to interpret. In particular, if an experiment extends over a considerable space of time, a gradual change in environmental conditions which affect the character studied will simulate a true inheritance of variations in the most perfect manner, so far as the resulting coefficients of correlation are concerned.

The real test for the occurrence of heritable variations within a clone must be the success or failure of an attempt to isolate, by selection, hereditarily diverse groups within a single clone.

In my own work on *Centropyxis*, both of these methods have been employed, and the evidence drawn from coefficients of correlation was used as a guide in the subsequent selection experiments.

In closing this brief introduction to the problem, I desire to express my deep indebtedness to Dr. H. S. JENNINGS, who suggested this problem to me, and who has greatly assisted both the experimental work and the preparation of this paper, by his kindly advice and criticism.

Material;—Centropyxis aculeata

Descriptive

The lobose rhizopod, *Centropyxis aculeata* Stein, is an excellent subject for genetic investigation. It is closely related to *Diffugia*, the subject of JENNINGS's recent (1916) paper, and, like *Diffugia*, its protoplasm is surrounded by a spine-bearing shell which fulfills all the requirements for genetic study. That is, it presents definitely measurable structural characters which are heritable, yet very variable, and which are entirely unaffected by growth or environmental influences during the life of the individual. The shape of the shell is even more favorable for study than that of *Diffugia*. The individuals used in this study correspond closely to the description of variety *discoides* PENARD (1902). In this form the chitinous shell, studded with sand grains and diatom tests, is decidedly flattened (see figure 2). The upper surface is convex and the lower slightly concave. Seen from above, as in figure 1, the outline of the shell is nearly circular, and the thin "anterior" edge, with its

smooth outline, is clearly distinguishable from the thick, rough "posterior" edge, which may bear from one to eleven spines, each tipped by a grain of quartz which looks like the cork in a bottle. In all but the oldest shells, the outline of the mouth, which is situated at the top of

FIGURE 1.—Dorsal view of living individual of *Centropyxis aculeata*.

a funnel-like infolding of the lower surface, is plainly visible through the shell, even when the organism is viewed from the upper side.

In order to permanently record the characters of an individual, then, it was necessary only to transfer the living organism to a minute drop of water on a slide and to trace the outline of shell and spines with a camera lucida, under the high power of the compound microscope. By focussing down a little the outline of the mouth was then added. Such camera outlines were made of every individual used in the statistical part of the work, and in one of the selection experiments. Some of them are reproduced, reduced in size, as illustrations for the present paper.

The chitinous shell surrounds an ameboid protoplast. At the time of cell division the protoplast swells, probably by the absorption of water, and projects out of the mouth of the old shell. This projection assumes the size and shape of a new shell, but one with its "antero-posterior" axis in just the opposite direction to that of the old shell. To the surface of this mold flow various sand-grains and diatom tests which have been picked up by the parent organism and stored within its body. Now,

over the entire surface of the exposed protoplast, a chitinous substance is secreted, in which, as it hardens, the foreign particles become firmly embedded. After the new shell is formed, the parent protoplast is par-

FIGURE 2.—Diagrammatic sagittal section of *Centropyxis aculeata*.

tially retracted into the old shell, the nucleus divides, and ultimately the protoplast pulls apart into two organisms, and the parent and its offspring separate.

The regular mitotic division of the nucleus during this process has been described and figured by SCHAUDINN (1903, 1911). Unfortunately, although a chromidial body is present (see figure 2), no account of its behavior during cell division has been published.

The new shell, when first formed, is almost white, in striking contrast to the dark brown color of old shells. The darker coloration is taken on gradually, first appearing as a faint yellowish tinge, which slowly deepens to brown. Even when four or five individuals have come from one in the course of a week or so, their relative ages, and therefore their parentage, is unmistakably indicated by their color. In practice, however, it was only very rarely that either of the two products of a division had time to divide again before they were separated. At ordinary laboratory temperatures, during the cooler months, *Centropyxis* divides about twice a week, at most. Inspection of the cultures at intervals of two or even three days was ordinarily sufficient to insure the isolation of parent and progeny before either had divided a second time.

In closing this brief description of the organism used in this experiment, I wish again to emphasize the fact that the new shell is completely produced by the parent organism before nuclear division begins. It is simply handed on to one of the products of the ensuing fission. Further, the new shell is in no sense molded on the old one. This is particularly

evident in *Centropyxis*, since, as figure 3 shows, the axes of old and new shells lie in exactly opposite directions. The number of spines and the shape of the shell, then, must be determined by precisely the same kind of determiners that underlie all the other hereditary characters of the species.

FIGURE 3.—Lateral view showing the position of old and new shells in the division of *Centropyxis aculeata*.

Previous breeding work on *Centropyxis*

The only other investigator who has studied the reproduction of *Centropyxis aculeata* is SCHAUDINN (1903, 1911). In his account of the life history of this species, he includes a few notes on the vegetative reproduction. Most of these statements I have been unable to confirm. He states, for instance, that the new shell is always larger than the old one. This I find not to be true. The new shell may be either slightly larger or slightly smaller than the old one. In line 30 (to be described later), with over 750 recorded fissions, the average size index of the parents was 21,969 sq. microns, that of the offspring 21,624 sq. microns. SCHAUDINN also says that he was never able to get an individual to divide more than once, and concludes that after this single fission the old individual lives on for some time and finally dies. In my cultures, individuals have repeatedly divided as many as ten times, some as many as fifteen times.

I have made no experiments on the sexual reproduction of *Centropyxis*, and will not discuss the life cycle. It is briefly described in SCHAUDINN's paper (1903). I can only state that my cultures have been repeatedly interrupted, usually in January or February, by what I take to be epidemics of sexual reproduction. At the beginning of such epidemics, the protoplasts of nearly every individual in the cultures

divide into a number of small amebae, which show no trace of a formed nucleus, but contain scattered chromatinic granules. These amebae escape from the shell, leaving it empty. I was unable to follow their further history.

Methods

Cultural technique

In keeping *Centropyxis* in cultures in the laboratory, my aim was to keep the environmental conditions as nearly like the natural conditions as possible. Each individual was isolated in the depression of a hollow-ground slide, in about five drops of pond water. To this water was added, at the time of isolation, a few strands of the *Oedogonium*, in the mats of which this rhizopod was living, as well as some debris taken from among the matted filaments of the alga. This debris furnished diatoms, the principal constituent of the food of the rhizopod, as well as sand grains to be used in the construction of new shells.

In these miniature aquaria the water was changed every two or three days, and once a week all the individuals were transferred to freshly prepared slides, to avoid any possibility of an injurious accumulation of either excretion products or bacteria. Whenever a fission occurred, the new individual was removed with a capillary pipette, outlined with the camera lucida, and transferred to a new culture slide.

Records were kept on small sheets of paper, each of which contained the number of a given individual, the date upon which it was isolated, its camera outline, and the numbers and dates of isolation of all of its progeny. Later, during the process of working up the data, the spine number and certain measurements of the outline drawing were added. Since these drawings, made from the living organism while the new shell was still clean and translucent, gave all the data required, no attempt was made to preserve the shells of individuals when they died or were discarded.

In some of the later work, when attention was centred wholly on the inheritance of spine number, the records were much less elaborate, consisting merely of the number of the specimen, the date on which it was isolated, its spine number, and the corresponding data for all of its offspring.

Statistical methods and use of terms

The following measurements were taken on each of the camera outlines: Diameter of shell and mouth in the "antero-posterior" axis.

Diameter of shell and mouth in an axis at right angles to the "antero-posterior" axis. To secure a number proportional to the area of the shell, the two diameters were multiplied together. This was called the "shell size index." In order to obtain a number related in some way to the form of the shell, the "antero-posterior" diameter was divided by the other diameter. This was called the "shell form ratio." The "mouth size index" and "mouth form ratio" were computed in the same way. All measurements were made in microns, giving the actual dimensions of the shell.

The coefficients of correlation given in this paper were computed by the improved methods devised by Dr. JENNINGS (see especially JENNINGS 1916, p. 416). So many of these constants are given that it seems inadvisable to attempt to publish even all of the correlation tables, much less the extensive pedigrees on which they are based. A few of the more important tables are included in the appendix.

A word as to terminology: In this paper, following the usage of SHULL (1912) and JOHANNSEN (1913), the term "pure line" will be considered as restricted to homozygous organisms reproducing by self-fertilization. The various strains of *Centropyxis* considered in this paper are properly to be spoken of as "clones," that is, families arising from a single ancestor by asexual reproduction. The more general terms "strains," "lines" and "races" are considered as synonymous and as having no specific meaning.

INHERITANCE WITHIN A POPULATION

Existence of diverse clones

The first requisite, in any study of inheritance within the asexual progeny of a single individual, is to be sure that the diverse clones, whose existence we assume in our population, really exist there. Next one must demonstrate that there exist between different clones hereditary diversities in the characters which are to be studied. Only too often (in the recent papers of EWING (1914 a, 1914 b, 1916), for example) this precaution has been neglected.

My first cultures of *Centropyxis aculeata* began in October, 1914, with the isolation of fifty individuals from a pipetteful of debris taken from a mat of *Oedogonium* growing on a lotus stem in the Homewood pond, Baltimore. About twenty of these lines died out during the first month, while I was learning the best culture methods. At the end of this time the thirty remaining lines all seemed to be distinct from each other. To keep the cultures within manageable bounds, only the four

largest clones were continued for two months more. In considering inheritance within a population, then, the population consists only of these four clones, numbers 9, 30, 41 and 43, containing in all 1049 individuals.

It was evident, from the very first, that at least two distinct strains existed in my material. A glance at table 1 or figure 4 shows clearly that line 30 differs most decidedly from the other three lines. Their small size, rapid fission rate and high spine number differentiates individuals of this line from the others at first sight.

The other three lines resemble each other more closely. The differences between them are more minute. Line 43 is the most clearly distinct of the three. Its higher fission rate alone would suffice to separate it from the others, and its distinctness is further shown by the slightly smaller size and lower spine number of its members.

TABLE I
Characters of the diverse lines.

Character	Line 9	Line 30	Line 41	Line 43
No. of individuals	93	749	126	81
Av. length	197.17	149.80	196.32	194.06
Av. breadth	188.25	144.50	188.33	184.02
Av. size index	371.25	216.48	370.19	357.23
Av. form ratio	1.048	1.038	1.044	1.056
Av. mouth size index	45.76	25.75	46.04	40.06
Av. mouth form ratio	.926	.920	.879	.923
Av. spine number	1.161	6.271	1.40	1.062
Av. number of days between two fissions	11.53	6.12	11.31	9.40

The other two lines, 9 and 41, are closely allied. The most striking difference between them lies in the higher average spine number of line 41, shown in table 1 and figure 5. Shell size and form are almost identical, as is also fission rate. The size and form of the mouth show slight differences. To anticipate the conclusions of this paper, it seems not improbable that lines 9 and 41 have been derived ultimately from a common ancestor.

It is to be noted, in connection with table 1, that the characters of size, spine number and fission rate show large differences between diverse clones, while the differences in form of shell and mouth are small.

When one recalls that the original ancestors of these lines were iso-

FIGURE 4.—Sample individuals (after camera outlines) of lines 9, 30, 41 and 43.

lated at the same time and from the same place, and that the individuals of the different clones were cultivated at the same time, under the same conditions, the slides containing them intermingled in the same moist chambers, it seems unnecessary to adduce any further evidence that diverse strains, differing in all the characters studied, do exist in *Centropyxis*.

Ancestral correlations within a population

In order to have a basis for comparison for the coefficients of correlation showing the inheritance of diversities within the single clones, coefficients of correlation were computed to show the degree of such

FIGURE 5.—Graphs of the percentage distribution of spine number in lines 9, 30, 41 and 43.

inheritance within the population. In table 2 are given the parental and grandparental coefficients of correlation for all the characters studied. In this table two points stand out clearly:

TABLE 2
Coefficients of correlation for the population.

Character	Parental correlation (1040 individuals)	Grandparental correlation (997 individuals)
Spine number..	.806 ± .007	.785 ± .008
Shell size	.903 ± .004	.773 ± .008
Shell form	.107 ± .022	.019 ± .021
Mouth size	.732 ± .010	.677 ± .011
Mouth form ..	.142 ± .021	.098 ± .021

First, the correlation is very high for shell size and high for spine number and mouth size, but low for the two form characters. This agrees with the conditions found in table 1. There the distinctions between the clones were definite in size and spine characters, but indefinite in form characters. So already two of the characters chosen for study have been practically eliminated for future work. It is perhaps signifi-

cant that these characters are ratios, a kind of character with which much biometrical work has been concerned.

In the second place, it is noteworthy that, in every case, the grandparental coefficient is decidedly smaller than the parental. As PEARSON (1910) has pointed out, according to the pure line hypothesis the offspring should resemble any other individual of the same clone, for example, their grandparent, just as much as they do their parent. PEARSON'S main argument against the evidence of the earlier pure line workers is this phenomenon of the diminishing ancestral correlation, which he considers satisfactory evidence that inheritance of variations within the pure line is occurring. It must be noted, however, that the objection to relying solely on the coefficient of correlation for proof of inheritance, because of the effect of a steadily changing environment (see page 177), applies with equal force to this phenomenon.

INHERITANCE WITHIN SINGLE CLONES

Measured by the coefficient of correlation

Ancestral correlations in line 30

Because of its rapid rate of division, many more individuals of line 30 were obtained than of any of the other races. The fact that this single clone contains 749 individuals makes it much the most suitable for statistical study. The other clones, lines 9, 41 and 43, with 93, 126 and 81 individuals, respectively, will be useful only for comparison. The numbers concerned are too small to admit of definite conclusions from these three lines.

In table 3 are given the coefficients of correlation for the first (parental), second (grandparental), third and fourth ancestral generations of line 30. It is at once evident that the five characters considered fall into two groups.

TABLE 3
Coefficients of correlation for line 30.

Character	Ancestral generations			
	1 (755 individuals)	2 (754 individuals)	3 (744 individuals)	4 (693 individuals)
Spine number	.082 ± .024	.034 ± .024	.009 ± .025	.007 ± .026
Shell size	.253 ± .023	.026 ± .025	.012 ± .025	.019 ± .026
Shell form	.062 ± .025	-.045 ± .025	-.056 ± .025	-.011 ± .026
Mouth size	.024 ± .024	.013 ± .025	-.033 ± .026	-.056 ± .026
Mouth form	.066 ± .024	-.043 ± .025	.066 ± .026	-.008 ± .026

The parental coefficients for spine number and shell size are significant, in comparison with their probable errors. All the coefficients for these two characters are positive, and there is a regular decrease in their size as one passes from more recent to more remote ancestral generations. This evidence points toward the inheritance of variations in size and spine number, even within the single clone.

The other three characters, shell form, mouth size and mouth form, show no such evidence of inheritance. The parental coefficients are less than three times their probable errors, many of the other coefficients are negative, and no diminishing ancestral correlation is to be found. It can be said only that while variations in these characters may possibly be inherited, this table gives no evidence leading to such a conclusion.

Ancestral correlations in the other clones

In table 4 are given, for comparison, the coefficients of correlation for the first and second ancestral generations in lines 9, 41 and 43.

In all of these lines, the numbers available were so small that no definite conclusions one way or the other can be drawn from them. As a glance at the table will show, they give no evidence for the inheritance of variations within the clone. Many of the coefficients, in both generations, are negative, hardly any are large enough to be significant, and in the few cases where there is a diminishing ancestral correlation, it is probably due to chance alone. One thing, only, is worth noting. The coefficients for spine number, in both generations, are all positive. This fact has no significance by itself, but, in comparison with the conditions shown by all the other characters, it may be considered to support the conclusion that variations in spine number may be inherited within the clone.

To sum up the purely biometrical part of the work, we may say:

1. Diverse clones exist in *Centropyxis*.
2. Within a population, diversities in size of shell and of mouth, and in spine number, are decidedly inherited, while diversities in form of shell and of mouth are only slightly inherited.
3. Within single clones, the evidence from coefficients of correlation indicates that variations in shell size and in spine number are inherited. The evidence that variations in the other characters are inherited is weak or lacking.

It was now decided to choose a single character for intensive study of the effect of selection. The statistical work has eliminated the two form characters, as showing only slight correlation even in a population.

There remain only spine number and the two size characters, which must be considered together, since they are closely correlated.

TABLE 4
Coefficients of correlation for lines 9, 41 and 43.
Line 9

Character	Parental correlation (93 individuals)	Grandparental correlation (81 individuals)
Spine number .	.148 $\pm$.069	.041 $\pm$.075
Shell size	-.121 $\pm$.069	-.060 $\pm$.075
Shell form	-.135 $\pm$.069	.120 $\pm$.074
Mouth size ...	-.071 $\pm$.069	-.155 $\pm$.073
Mouth form ..	-.053 $\pm$.070	-.252 $\pm$.070

Line 41

Character	Parental correlation (130 individuals)	Grandparental correlation (116 individuals)
Spine number .	.006 $\pm$.060	.125 $\pm$.060
Shell size	-.062 $\pm$.060	-.074 $\pm$.063
Shell form	-.057 $\pm$.060	.152 $\pm$.062
Mouth size ...	-.017 $\pm$.060	.117 $\pm$.059
Mouth form ..	.185 $\pm$.058	-.061 $\pm$.060

Line 43

Character	Parental correlation (81 individuals)	Grandparental correlation (69 individuals)
Spine number .	.144 $\pm$.073	.215 $\pm$.076
Shell size	.088 $\pm$.075	-.019 $\pm$.080
Shell form	-.082 $\pm$.075	-.053 $\pm$.080
Mouth size ...	-.148 $\pm$.074	.117 $\pm$.079
Mouth form ..	-.080 $\pm$.075	.240 $\pm$.075

Shell size, as has been shown, gives the highest parental coefficient of correlation, both in the population and in line 30. In the other lines

the coefficient was often negative. Spine number shows a decided correlation in the population and a significant one in line 30, and in all the other lines the coefficient was always positive. It is easy to imagine a way in which shell size could be "inherited" in a purely mechanical sense, since the old shell acts as a limit to the possible growth of the organism within it, and this organism then serves as a mold over the surface of which the new shell is secreted. Finally, spine number is the character which appears to have the least relation to the physiological activity of the organism, and is the one which can be most accurately and easily determined. For these reasons, spine number was chosen as the subject for selection experiments.

After the objections which have just been raised to the use of shell size as a subject for selection experiments, it is advisable to determine whether variations in shell size are correlated with variations in spine number. This is especially necessary, since in our population we find that the large lines have few spines and the small lines many spines. Within the clone, however, this does not hold. The coefficient of correlation between shell size and spine number for the 745 individuals of line 30 is $.019 \pm .025$, that is, for all practical purposes, zero.

Inheritance proved to occur by selection experiments

Selection experiments 1 and 2; contradictory results of two brief experiments in mass selection

In order to bring out clearly the reasons which led me to abandon mass selection for individual selection based on a progeny test, it seems worth while to give a brief account of two experiments in mass selection which came to an abrupt and unexpected end. The first of these experiments was ended by an epidemic of sexual reproduction, the second by a period of extremely hot weather which probably acted indirectly by accelerating the multiplication of undesirable bacteria.

Neither of these experiments lasted long enough for the application of the real test of their effectiveness, the free multiplication of two altered strains in comparison with each other. But since selection consists only in allowing certain individuals to breed and denying that privilege to others, a good idea of whether selection is being effective can be gained by comparing the average spine number of the progeny produced by high- and low-selected groups during the same period. Such comparisons for selection experiments 1 and 2 are given in tables 5 and 6.

Selection experiment 1. Table 5 shows the progressive effect of selec-

tion in a small many-spined clone closely resembling line 30. The progenitor of this race was isolated from algae taken from the Homewood pond on October 8, 1914. On November 9, 52 individuals had been secured by the multiplication of the original one, and selection began. On December 14 an epidemic of sexual reproduction put an end to the experiment.

TABLE 5

Selection experiment 1, mass selection.

High series: Parents with 6 or more spines.

Low series: Parents with 5 or less spines.

Mean of clone (52 individuals) when selection began: 6.56 spines.

Between selections	Progeny produced in				Difference between high and low means
	High series		Low series		
	Individuals	Mean spine number	Individuals	Mean spine number	
1 and 2.....	49	6.57 $\pm$.10	47	6.35 $\pm$.10	.22 $\pm$.14
2 and 3.....	30	7.36 $\pm$.12	23	5.75 $\pm$.14	1.61 $\pm$.18
3 and 4.....	9	7.76 $\pm$.22	7	5.50 $\pm$.25	2.26 $\pm$.32

In this experiment simple mass selection was used. The clone was divided into two parts, the individuals with six or more spines being classed as high, those with five or less as low. In each group only those progeny whose spine numbers lay on the same side of the mean (6.56 spines) as their parent's were allowed to reproduce.

As table 5 shows, the difference in mean spine number between the progeny produced by the high and low groups was only .22 of a spine after the first selection, but rose to 1.61 spines after the second, and to 2.26 spines after the third. Apparently, then, in this clone, three selections sufficed to isolate within the clone high and low strains with a mean spine number of about 7.75 and 5.5 spines, respectively. This quick and decided effect of selection was most unexpected, and it is most unfortunate that the onset of sexual reproduction ended this experiment so prematurely. The numbers involved are too small to allow of any definite conclusion from an experiment of such short duration.

Selection experiment 2. In table 6 is shown the course of a similar experiment in a clone with fewer spines and of larger size. The progenitor of this clone was isolated on July 3, 1915, from among algae at-

tached to a dead stick in an inlet of Nobska pond near Woods Hole, Mass. Selection was started on July 28, when the clone contained 55 individuals. The last individual of this race died on August 16, after a period of very hot weather.

TABLE 6

Selection experiment 2; mass selection.

High series: Parents with 2 or more spines.

Low series: Parents with 1 or 0 spines.

Mean of clone (55 individuals) when selection began: 1.51 spines.

Between selections	Progeny produced in				Difference between high and low means
	High series		Low series		
	Individuals	Mean spine number	Individuals	Mean spine number	
1 and 2.....	25	1.96 ±.14	29	0.31 ±.13	1.65 ±.19
2 and 3.....	5	1.40 ±.30	9	0.00	1.40 ±.30
3 and 4.....			5	0.40 ±.30	

When selection began, the mean spine number of the clone was 1.51 spines. The high and low groups therefore consisted of individuals with two or more and one or no spines, respectively. The method of selection was the same as in experiment 1. Although the clone died out before more than one or two selections could be made in most of the lines, table 6 shows that this clone behaved very differently from the one used in experiment 1. The first selection gave a decided difference (1.65 spines) between high and low groups, as against a very slight one in the first experiment (.22 spines). And after the first selection no further result was noticeable. This differs decidedly from the steady progressive effect which was found in experiment 1.

In both these experiments mass selection was used. In fact, in almost all the experiments on the effect of selection within pure lines or clones, mass selection has been used exclusively. Mass selection is based entirely on the external appearance of the individual, and it is evident that the degree of effectiveness of the selection depends very largely on the extent to which the genetic constitution of the individual resembles its external aspect. Formerly mass selection within populations was extensively practiced, and one of the greatest achievements of the pure line workers has been the pointing out that by selecting on a basis of the character of the progeny produced by an individual, the same results

can be obtained with much fewer selections. This is true, of course, because the average character of the progeny of an individual gives a much more accurate test of its hereditary constitution than does its external appearance.

If this is true in a population, may it not also be true within a clone? And if one seeks for an effect of selection within a clone, why not use the type of selection which has been found to be most effective in populations?

Selection experiment 3; quick effect of individual selection based on progeny

In accordance with the train of reasoning set forth above, individual selection based on the average spine number of the progeny was used in my next experiment. On October 10, 1915, an individual was isolated from the Homewood pond. After it had produced five progeny it was discarded. Each of these progeny, in turn, produced five offspring and was then discarded. Of these five sets of offspring, the set with the highest average spine number and the set with the lowest were allowed to produce five progeny each. The other three sets were discarded. Then another selection was made, the set of progeny with the highest average spine number being retained on the high side, and *vice versa* on the low side. This method of selection was continued for another generation.

TABLE 7

Selection experiment 3: Individual selection, based on the character of the progeny.

Generation 1:	5 (1.0)				
Generation 2:	5a (0.8)	5b (0.3)	5c (0.2)		
Generation 3:					
High series	5a1 (1.5)	5a3 (0.0)	5a4 (1.0)	5a5 (0.0)	
Low series	5c1 (0.3)	5c2 (0.2)	5c5 (1.0)		
Generation 4:					
High series	5a1a (1.2)	5a1c (0.0)	5a1d (1.0)		
Low series	5c2a (0.2)	5c2b (1.0)	5c2c (0.6)	5c2d (0.3)	
Generation 5:					
High series	5a1a1 (1.8)	5a1a2 (1.7)	5a1a4 (1.0)	5a1a5 (1.2)	
Low series	5c2a2 (0.2)	5s2a3 (0.4)	5s2a5 (0.6)		
Generation 6 and following generations:					
High series—56 individuals derived from 5a1a1—average spine number..	1.68±.09				
Low series—51 individuals derived from 5c2a2—average spine number..	0.22±.002				
Difference between high and low means.....	1.46±.085				

Table 7 shows the pedigree of this experiment. In this table, 5a, 5b and 5c are the progeny of individual 5; 5a1, 5a3, 5a4 and 5a5 are the

progeny of 5a; and so on. The numbers in parentheses, following the numbers which represent the individuals, are the average spine numbers of their five (in a few cases, six) progeny. All individuals which did not produce at least five progeny have been omitted from the table.

On December 14, after four selections had been made in both high and low groups, selection was discontinued, and the final individuals 5a1a1 and 5c2a2 and their progeny were allowed to multiply freely. Because of cold weather, the fission rate was very low, and when about fifty individuals had been produced in each group the experiment was ended on February 3, 1916.

The numbers concerned in this experiment are not large, but the difference between the means of the two final groups (1.46 spines) is so great in proportion to its probable error (.085) that there can be no doubt that in this case selection has isolated, within a single clone, two strains with decidedly different average spine numbers. In figure 6 are shown a few characteristic members of the two groups, from camera outlines.

In this experiment there is one possibility of error. When one is selecting individuals instead of masses of individuals,—a single line instead of a number of lines,—an excellent opportunity is offered for the multiplication of any mutation that may occur. Recognizing this fact, and being now quite convinced in my own mind that selection within the clone was having an effect, I turned to mass selection once more. In mass selection, with many parallel lines, there is no chance for the extreme multiplication of a mutation, and if one does occur it changes the final result only to a very slight degree. And if selection within a clone is really effective, mass selection on a large scale, in a favorable race, should show such an effect. (By a favorable race, I mean, of course, one in which the external appearance of an individual is a fairly accurate index of its genetic constitution.)

*Selection experiment 4; decided effect of mass selection
in a favorable race*

On June 29, 1916, an individual of *Centropyxis* was isolated from material collected from a freshwater pond on Cuttyhunk Island near Woods Hole, Mass., and was allowed to multiply. The course of this experiment is summarized in table 8. On August 1, when the line comprised 133 individuals, it was divided into two groups. This race was of large size, with a medium number of spines. On August 1 the average spine number was almost exactly 2.5 spines. All individuals with 0, 1

FIGURE 6.—Sample individuals of high and low groups at the end of selection experiment 3.

or 2 spines, 63 in number, were set aside as progenitors of low lines, and all individuals with 3 or 4 spines (there were 70 of these) became progenitors of high lines. As soon as any of these individuals divided, the new one was examined. In the low lines, if the new individual had 3 or 4 spines, it was destroyed; but if it had 0, 1 or 2 spines, it was retained and its parent destroyed. In the high lines, of course, the reverse procedure was followed. Such selection was continued until August 26, only a few days before I had to leave Woods Hole. From August 26 to September 3, inclusive, no selection was practiced, and all the remaining individuals of both groups were allowed to multiply freely.

TABLE 8
Selection experiment 4; mass selection.
High series: Parents with 3 or 4 spines.
Low series: Parents with 0, 1 or 2 spines.

Period of	Time	Number of individuals	Average spine number	Difference between high and low lines
Multiplication	June 29-July 8	9	3.11 $\pm$.22	
	July 9-18	28	2.39 $\pm$.14	
	July 19-30	96	2.44 $\pm$.08	
Selection	Aug. 1-10			
	High lines	146	2.71 $\pm$.05	.55 $\pm$.09
	Low lines	133	2.16 $\pm$.07	
	Aug. 11-20			
	High lines	167	2.55 $\pm$.05	.48 $\pm$.09
	Low lines	146	2.07 $\pm$.07	
Multiplication after selection	Aug. 21-25			
	High lines	89	2.77 $\pm$.06	1.00 $\pm$.09
	Low lines	79	1.77 $\pm$.07	
	Aug. 26-Sept. 3			
	High lines	192	2.75 $\pm$.05	.99 $\pm$.069
	Low lines	180	1.76 $\pm$.05	

During this time more than 175 offspring were produced by each group. The average spine numbers of these two sets of progeny differ by nearly a whole spine, with a probable error so small as to be negligible. This experiment, I think, proves conclusively that the effect of selection which has been demonstrated in experiment 3 is due to the inheritance of minute variations in spine number within the clone.

In this mass selection experiment, the appearance of a mutation has little effect on the ultimate result. Something comparable to a mutation did, in fact, appear in the low series of experiment 4. During the first period of multiplication, individual 1d1, with one spine, began producing progeny all of which had no spines. Progeny with no spines were comparatively rare among the other low lines. All these progeny of 1d1 also produced progeny with no spines, predominantly, but with a few one-spined progeny. The early appearance of this mutation gave it greater prominence than it would have enjoyed later on, since 1d1 and three of its offspring were included among the progenitors of low lines. In all, 39 individuals can be traced back to 1d1 as their original ancestor. Of these, 33 had no spines and 6 had one spine. But the effect on the final result of the experiment is inconsiderable. If all of the offspring of the mutation are left out of the table, the number of individuals produced by the low lines between August 26 and September 3 is reduced to 168, with an average spine number of 1.867. This still gives a difference between the average spine numbers of the high and low groups of $.88 \pm .07$.

The very favorable character of the race employed in experiment 4, that is, the close correspondence between its external appearance and genetic constitution, is shown by the high correlation between parent and progeny. The coefficient of correlation between parents and progeny, for spine number, is $.376 \pm .017$ for the 1277 individuals of this race. This is the highest coefficient of correlation within a clone that has been obtained in the course of my experiments. If the mutation 1d1 and its offspring are omitted, the coefficient becomes $.282 \pm .018$.

CONCLUSIONS

Results of the present investigation

The present study of inheritance in *Centropyxis* may be divided into two parts, the first concerned with a statistical analysis of inheritance within the population and within single clones, the second with an attempt to test the inheritance of variations in a single definite character within the clone, by the method of selection.

The results of the biometrical study of inheritance in *Centropyxis* may be summarized as follows:

1. There exist in the species, *Centropyxis aculeata*, many diverse strains or clones which, under identical environmental conditions, differ from each other decidedly in spine number, shell size, mouth size and fission rate, but which differ only slightly in shell form and mouth form.

2. In a population, the coefficients of correlation show that variations in spine number, shell size and mouth size are strongly inherited, while variations in shell form and mouth form are weakly inherited.

3. In a single clone, the coefficients of correlation show that variations in spine number and shell size are probably inherited.

4. If a diminishing ancestral correlation is considered a criterion of inheritance of variations within a clone, the coefficients of correlation of the population give such evidence of inheritance for all the characters studied, and the coefficients of correlation within a clone give such evidence for spine number and shell size.

5. From the evidence adduced under 2, 3 and 4 above, it was concluded that variations in spine number and shell size are inherited within the clone. Since shell size might be "inherited" without the mediation of the nucleus, spine number was considered the most favorable character for use in selection experiments.

The results of the experiments in the selection of variations in spine number may be summarized as follows.

6. In the brief experiments 1 and 2, mass selection based on external appearance gave contradictory results in two different races. In explanation of this, it is assumed that in different races the degree of correlation between the external appearance of the individual and its genetic constitution is diverse; in some the two are highly correlated, in others but little.

7. In experiment 3, individual selection based on the average spine number of the progeny of the individual under consideration resulted in a rapid isolation of two decidedly different strains within a single clone.

8. Since the result in experiment 3 might have been due to the appearance and preservation of a mutation, mass selection was again tried. In experiment 4, on a very favorable race (as is shown by its high parental correlation), mass selection based on the external appearance of the individual was sufficiently effective to isolate two distinct strains from within a single clone. In this experiment 1277 individuals were obtained, all coming originally from a single specimen.

9. In this experiment 4, something resembling a mutation did appear. Its 39 progeny were traced out, and it was shown that their elimination from the record makes no essential change in the final result.

The present status of the "pure line" hypothesis

In considering the bearing of this work and other similar investigations on the concept of the "pure line," one must carefully distinguish

between the two parts of which this hypothesis is composed. The term "pure line" was originally applied to groups of individuals derived by self-fertilization from a common homozygous ancestor. Such lines within a single species prove to be relatively constant diverse strains. That such relatively constant strains do occur in a great variety of plant and animal species, is an observed fact, and has been confirmed by every investigator who has entered this field. The further idea, that within one of these "pure lines" no variation in genetic constitution is possible, except by a sudden mutation, large or small, is not direct observation but hypothesis. It is an hypothesis, to be sure, which was based on all the experimental evidence available at the time it was formulated, but it is still an hypothesis, subject to revision at any time when new experimental results demand it.

For some years, the great majority of the experimental results tended to confirm this hypothesis, and it gained such a firm footing that all the contradictory cases were dismissed as aberrant types, results of a failure to properly control the environment, observational errors, etc. But of late the tide seems to be turning somewhat. STOUT (1915) in *Coleus*, STOCKING (1915) in *Paramecium*, MIDDLETON (1915) in *Stylonychia*, and JENNINGS (1916) in *Diffugia*, have all shown that minute heritable variations do occur within clones derived from a single ancestor by vegetative multiplication. In the present work on *Centropyxis*, the same conclusion is reached.

We must admit, it seems to me, that in these plants and protozoans reproducing asexually, small variations in genetic constitution do continually occur. How does this affect the "pure line" hypothesis? The issue can be met in two ways.

In the first place, it is still possible to reaffirm the entire validity of the original pure line hypothesis. The new conflicting results, like previous ones, can still be explained away as due to complicating factors. For example, the inheritance of abnormalities in *Paramecium* (STOCKING 1915) can be considered to follow from an abnormal condition of the macro-nucleus; the inheritance of variations in fission rate in *Stylonychia* (MIDDLETON 1915) might be due to the accumulation of waste products in the cytoplasm of the slowly-dividing, large-sized group. And in *Diffugia* (JENNINGS 1916) and *Centropyxis* (the present paper), until we know more about the process of cell division in these organisms, the observed inheritance of variations can be explained by the assumption that a process of somatic segregation or fractionation of the chromidial body takes place at each cell division, as has been suggested by MORGAN

(1916, p. 185). Other similar cases could doubtless be supplied with other explanations along the same lines.

But there is another viewpoint, alternative to the first one, which seems to me to be preferable. According to this idea, one may assert that the pure line concept is correct on the average, only; that it is a mathematically correct expression of the mean result of inheritance under natural conditions. But, just as the Galton-Pearson law of ancestral inheritance, though accurate mathematically, does not fit the details of Mendelian heredity, on the average result of which it is presumably based, so the pure line hypothesis, though true for average results, may be thought of as not holding in individual cases. Or, to look at the question from another side, just as species were found to consist of a number of pure lines, diverse from each other in minor details of genetic constitution, so we may think of pure lines as made up of a number of individuals, differing from each other in even smaller details which are, nevertheless, hereditary.

According to this idea, we can assume that each individual inherits, in general outlines, the genetic constitution of its species; in smaller details, that of its pure line; in still more minute details, that of its individual parent; and yet may differ from this parent in some hereditary points. There is a growing inclination among geneticists to designate all such hereditary differences, mutations, regardless of their magnitude, but I am opposed to this usage and would limit the word mutation to hereditary differences which are fairly large. If such differences are very minute, they can be termed genetic fluctuations, micro-mutations, or any other term which is found to meet the needs of those who work in this field.

So far as our present knowledge goes, either of these two views may be championed with equal success. The final decision between them must be left to the experimental results of the future.

LITERATURE CITED

- AGAR, W. E., 1914 Experiments on inheritance in parthenogenesis. Phil. Trans. Roy. Soc. London B **205**: 421-489.
- BARBER, M. A., 1907 On heredity in certain micro-organisms. Kansas Univ. Science Bull. **4**: 1-48.
- EAST, E. M., 1910 The transmission of variations in the potato in asexual reproduction. Conn. Exp. Sta. Rep. pp. 119-160.
- EWING, H. E., 1914 a Pure line inheritance and parthenogenesis. Biol. Bull. **26**: 25-35.
- 1914 b Notes on regression in a pure line of plant lice. Biol. Bull. **27**: 164-168.
- 1916 Eighty-seven generations in a parthenogenetic pure line of *Aphis avenae* Fab. Biol. Bull. **31**: 53-112.

- HANEL, ELISE, 1908 Vererbung bei ungeschlechtlicher Fortpflanzung von *Hydra grisea*. Jenaische Zeitschr. **43**: 321-372.
- JENNINGS, H. S., 1908 Heredity, variation and evolution in Protozoa. II. Heredity and variation in size and form in *Paramecium*, with studies of growth, environmental action and selection. Proc. Am. Phil. Soc. **47**: 393-546.
- 1916 Heredity, variation and the results of selection in the uniparental reproduction of *Diffugia corona*. Genetics **1**: 407-534.
- JOHANNSEN, W., 1903 Ueber Erblichkeit in Populationen und in reinen Linien. v + 68 pp. Jena: Gustav Fischer.
- 1913 Elemente der exakten Erblichkeitslehre. Zweite Ausgabe. xi + 723 pp. Jena: Gustav Fischer.
- LASHLEY, K. S., 1915 Inheritance in the asexual reproduction of *Hydra*. Jour. Exp. Zool. **19**: 157-210.
- 1916 Results of continued selection in *Hydra*. Jour. Exp. Zool. **20**: 19-26.
- MIDDLETON, A. R., 1915 Heritable variations and the results of selection in the fission rate of *Stylonychia pustulata*. Jour. Exp. Zool. **19**: 451-503.
- MORGAN, T. H., 1916 A critique of the theory of evolution. x + 197 pp. Princeton: Princeton University Press.
- PEARSON, KARL, 1910 Darwinism, biometry and some recent biology. I. Biometrika **7**: 368-385.
- PENARD, E., 1902 Faune rhizopodique du bassin du Léman. 714 pp. Geneva: W. Kündig et fils.
- SCHAUDINN, F., 1903 Untersuchungen über die Fortpflanzung einiger Rhizopoden. (Vorl. Mitt.) Arb. a. d. Kais. Gesundheits-Amptes. **19**: 547-576.
- 1911 FRITZ SCHAUDINN's Arbeiten. xv + 612 pp. Hamburg und Leipzig: Leopold Voss. (Contains paper of 1903 reprinted with illustrations added.)
- SHULL, G. H., 1912 Genotypes, biotypes, pure lines and clones. Science N. S. **35**: 27-29.
- STOCKING, R. J., 1915 Variation and inheritance of abnormalities occurring after conjugation in *Paramecium caudatum*. Jour. Exp. Zool. **19**: 387-449.
- STOUT, A. B., 1915 The establishment of varieties in *Coleus* by the selection of somatic variations. Carnegie Institution of Washington Publ. No. 218, 80 pp.
- WINSLOW, C. E. A., and WALKER, L. T., 1909 A case of non-inheritance of fluctuating variations in bacteria. Jour. Infect. Diseases. **6**: 90-97.

APPENDIX

Some important correlation tables

TABLE 9

Correlation between spine numbers of parents and their progeny in the population.

	Progeny											Totals
	0	1	2	3	4	5	6	7	8	9	10	
0	25	33	18	2	2							80
1	26	37	30	6	3							102
2	19	41	19	4	4	1						88
3	6	12	6	3	1	2	2	1				33
4		3		2		6	15	7	1			34
5		1		5	10	33	43	33	17	3	3	148
6				8	8	51	109	49	31	13	2	271
7				8	2	28	85	41	23	6	2	195
8				3	1	18	32	15	9	2	1	81
9					1		8	8	2	1		20
10							1	1	2			4
Totals	76	127	73	41	32	139	295	155	85	25	8	1056

TABLE 10

Correlation between shell size indices of parents and their progeny in the population.

	Progeny															Totals
	17	19	21	23	25	27	29	31	33	35	37	39	41	43	45	
17	2	5	16	5												28
19	8	33	71	16	4	1	4									137
21	6	69	192	71	26	1	7									372
23		20	85	48	15	6	3									177
25			15	6	3	2	1		1							28
27						1	1		1							3
29							1				1	1	2			5
31									7	5	5	8	1		1	27
33							3	2	8	14	7	13	2		4	53
35								3	12	10	12	21	2	1	3	64
37							6	2	12	15	7	23	3		2	70
39							1		6	8	7	20	3		3	48
41							1		1	3	6	5		1		17
43										1	2	4				7
45										1		2				3
Totals	16	127	379	146	48	12	27	7	48	57	47	97	13	2	13	1039

TABLE II

Correlation between shell size indices of grandparents and progeny in the population.

		Progeny															Totals
		17	19	21	23	25	27	29	31	33	35	37	39	41	43	45	
Grandparents	17		4	13	6	1	1	2									27
	19	1	22	71	15	4	14	10									137
	21	1	49	201	59	28	23	14									375
	23		21	96	30	11	12	9									179
	25		1	17	1	1	3	3									26
	27							1									1
	29							2				1			1		4
	31							1	1	5		4	7			4	22
	33							2		6	7	9	19			6	49
	35							3		5	9	11	21		1	7	57
	37							3	5	8	7	8	24			6	61
	39							1		2	6	5	19		1	5	39
	41							2		1	3	4	5			2	17
43							1	1	1				2		1	6	
45									1							1	
Totals		2	97	398	111	45	53	54	8	28	32	42	97	0	3	31	1001

TABLE 12

Correlation between spine numbers of parents and progeny in line 30.

		Progeny								
		3	4	5	6	7	8	9	10	Totals
Parents	3	1		2	2	1				6
	4	2		6	15	7	1			31
	5	5	10	33	43	33	17	3	3	147
	6	8	8	51	109	49	31	13	2	271
	7	8	2	28	85	41	23	6	2	195
	8	3	1	18	32	15	9	2	1	81
	9		1		8	8	2	1		20
	10				1	1	2			4
Totals		27	22	138	295	155	85	25	8	755

TABLE 13

Correlation between spine numbers of grandparents and progeny in line 30.

		Progeny							Totals	
		3	4	5	6	7	8	9		
Grandparents	3	1	1	1	2				5	
	4	6	3	6	6	6	4		31	
	5	7	5	21	70	33	7	3	146	
	6	22	11	45	119	55	10	9	271	
	7	16	9	39	81	37	10	4	196	
	8	6	3	14	36	14	6	2	81	
	9		1	3	10	4	1	1	20	
	10				3	1			4	
	Totals		58	33	129	327	150	38	19	754

TABLE 14
Correlation between the spine numbers of great-grandparents and progeny in line 30.

		Progeny							Totals
		3	4	5	6	7	8	9	
Great-grandparents	3	3	1	1					5
	4	5	1	5	14	3	1	1	30
	5	26	10	27	60	20	2		145
	6	39	16	52	128	26	5	2	268
	7	37	11	33	85	21	3	2	192
	8	14	2	17	37	6	2	2	80
	9	7		2	9	2			20
	10				4				4
Totals		131	41	137	337	78	13	7	744

TABLE 15
Correlation between shell-size indices of parents and progeny in line 30.

	Progeny														Totals
	16	17	18	19	20	21	22	23	24	25	26	27	28	29	
Parents	16							1							1
	17			1		3	1	1							6
	18	2		2	2	3	9	1	2						21
	19	2	1	6	9	11	8	2		2					41
	20	3	2	8	10	22	30	10	4	2		1		4	96
	21	2	1	11	25	44	53	21	9	12				3	181
	22	2	1	9	24	41	53	31	10	14		1		4	190
	23			2	11	21	38	23	10	7	3	4	1	3	123
	24			1	6	10	16	13	2	5		1			54
	25					4	8	3	1	2		2		1	21
	26					1	2	1	1	1					6
	27													1	1
	28											1			1
	29											1			1
Totals															
	11	5	40	87	160	218	107	39	45	2	11	1	16		743

TABLE 16

Correlation between spine numbers of parents and progeny in line 9.

	Progeny					Totals
	0	1	2	3	4	
0	10	11	6		1	28
1	7	8	12	2	1	30
2	5	11	8	1	1	26
3	1	2	2		1	6
4		2				2
Totals	23	34	28	3	4	92

TABLE 17

Correlation between spine numbers of parents and progeny in line 41.

	Progeny						Totals
	0	1	2	3	4	5	
0	9	8	8	2	1		28
1	16	14	8	2	2		42
2	12	15	7	2	3	1	40
3	5	7	3	2			17
4		1					1
5		1					1
Totals	42	46	26	8	6	1	129

TABLE 18

Correlation between spine numbers of parents and progeny in line 43.

	Progeny				Totals
	0	1	2	3	
Parents	0	6	14	4	24
	1	3	15	10	30
	2	2	15	4	22
	3		3	1	4
Totals	11	47	19	3	80

TABLE 19

Correlation between spine numbers of parents and progeny during selection experiment 4.

	Progeny					Totals
	0	1	2	3	4	
Parents	0	26	14			40
	1	11	116	82	78	311
	2		70	70	106	279
	3	1	56	82	204	441
	4		41	34	84	205
Totals	38	297	268	472	201	1276