bmj.com news roundup

Full versions of these stories are available at: bmj.com/content/vol331/issue7516/#NEWS_ROUNDUP

UK government fails to tackle weaknesses in drug industry

The UK government has failed to tackle the weaknesses in the pharmaceutical industry in its response to a report published earlier this year on the industry's role and influence, says a leading critic.

Ike Iheanacho, editor of the *Drug and Therapeutics Bulletin*, a publication from the consumer body Which?, told the *BMJ* that healthcare staff needed to know if they had been misled by drug advertisements. The government has not acted on many of the recommendations made in a highly critical report on the influence of the pharmaceutical industry on drug regulation, healthcare professionals, and the NHS, he said.

The report, published by the House of Commons' health select committee on 5 April, dealt with the growing influence of the industry on prescribing and the lack of transparency in the drug regulatory system (*BMJ* 2005;330:805).

"The pharmaceutical industry is entering a new era of openness. It is important that all parties work together so that the concerns about the industry expressed by the select committee are dispelled," said health minister Jane Kennedy.

Liza Gibson *London*

The Government Response to the Health Committee's Report on the Influence of the Pharmaceutical Industry is available at www.dh.gov.uk

Study highlights extent of medication errors in hospitals

A systematic review by Canadian researchers of 22 studies including 3755 patients showed that errors in prescription medication histories at hospital admission were "disturbingly common and potentially harmful to patients." Such errors occurred

Health consequences after Chernobyl disaster are less than feared

More than 4000 people could eventually die of radiation exposure from the disaster at the Chernobyl nuclear power station in 1986, but the long term health effects have been less than originally feared.

A report published this week by an international team of scientists, *Chernobyl's Legacy: Health, Environmental, and Socioeconomic Impacts*, found that fewer than 50 deaths so far could be directly attributed to radiation from the disaster. Nearly all of these were in rescue workers, who were exposed to high levels of radiation. The report predicts a further 3940 deaths from radiation induced cancer and leukaemia in the 200 000 emergency workers who were in the area in 1986-7, 116 000 evacuees, and 270 000 residents of the most contaminated areas, based on estimates of the radiation doses that they received. This is far lower than earlier speculations that radiation exposure might result in tens of thousands of deaths.

Figures analysed in the report show that about 4000 cases of thyroid cancer, mainly in people who were children and adolescents at the time of the disaster, have resulted from contamination.

Susan Mayor *London*

The report is available at www.iaea.org/NewsCenter/Focus/Chernobyl/pdfs/05-28601_Chernobyl.pdf

in up to 67% of cases in the studies, which were published between 1966 and April 2005 (*CMAJ* 2005;173:510-5).

Between 27% and 54% of patients had at least one medication history error and 19-75% of the discrepancies were unintentional. In six of the studies (n=588 patients) the investigators estimated that 11-59% of the medication history errors were clinically important.

The investigators searched Medline, Embase, and CINAHL for the articles and bibliographies of papers subsequently retrieved from the search. They reviewed all published studies with quantitative results that compared prescription medica-

tion histories obtained by physicians at the time of hospital admission with comprehensive medication histories. Three reviewers independently abstracted data on methodological features and results.

Prescription drugs most often involved in errors were cardiovascular agents (nitrates, digoxin, or β blockers), sedatives (benzodiazepines), and analgesics (non-steroidal anti-inflammatory drugs and opioids).

The study indicates a need for a systematic approach to ensuring that doctors take accurate drug histories at hospital admission and says that doctors may benefit from additional training.

David Spurgeon *Quebec*

Dutch government rejects tough measures on genital mutilation

Tough proposals to combat female genital mutilation—including examining the sexual organs of all girls up to the age of 14 and doctors being legally obliged to report all cases—have been rejected by the Dutch government. Several medical bodies have agreed with the government that such measures could deter parents and children, damaging school health services that currently reach 95% of the population.

The government has instead announced a £2m (\$3.7m; €2.9m) three year campaign geared towards prevention and information among at risk, largely immigrant, populations in the main cities. Pharos, a national organisation for refugees and health, will draw up the plans together with local and youth health services.

Medical professionals will have to follow a new national code when dealing with suspected cases, however. Prosecutions will also be made easier by delaying the start of the time limit on bringing a prosecution until the victim reaches 18.

Tony Sheldon *Utrecht*

Passive smoking increases children's risk of nasal cancer

Passive smoking in childhood may increase the risk of nasal cancer, says a study of children whose parents had lung cancer. The incidence of bladder and kidney cancer was also greater in the offspring, possibly as a result of exposure to tobacco carcinogens either in the womb or through breast feeding, say researchers writing online in the *International Journal of Cancer* on 10 August (<http://www3.interscience.wiley.com>, doi: 10.1002/ijc.21387).

Although the carcinogenic

effects of active smoking have been seen at many sites, the effects of passive smoking and exposure in pregnancy and breastfeeding are less well documented, say the researchers.

The authors used the nationwide Swedish family cancer database from 1958 to 2002 to see whether the children of parents with lung cancer are at risk of cancer that could not be explained by their smoking or familial risk. Most patients with lung cancer are smokers, and in populations with long term cigarette use, the population attributable risk of smoking is thought to be up to 90%.

The database used included 17 693 mothers and 41 838 fathers with lung cancer, and a total of 173 715 cancers were recorded in their offspring aged up to 70.

Roger Dobson *Abergavenny*

European public are unaware of heart failure symptoms

Only 3% of nearly 8000 randomly selected people from nine European countries could identify heart failure from a description of typical symptoms. This is despite 6% of people claiming that someone in their family had had the condition, an international survey of public awareness published in the *European Heart Journal* online on 31 August has found (<http://eurheartj.oxfordjournals.org>, doi:10.1093/eurheartj/ehi447).

The lead author, Willem Remme, professor of medicine and director of the Sticres Cardiovascular Research Institute, Rhooen, in the Netherlands, described the low awareness as "shocking" and "putting lives at risk."

"It has serious implications for individuals and for public health throughout Europe," he said. "If the public don't understand how common and how life threatening this condition is then they are not likely to seek medical help early, and they are also unlikely to demand appropriate measures from health-

care providers."

The level of awareness was described as dismal compared with that for transient ischaemic attack, which half of those surveyed identified from a list of symptoms, or angina, which almost a third correctly identified.

David Spurgeon *Quebec*

Imported syringes triggered meningitis in Sri Lanka

Health authorities in Sri Lanka have said that imported injection syringes might have triggered an outbreak of meningitis in patients who had received spinal anaesthesia in four hospitals.

Two maternity and two teaching hospitals reported 14 cases of meningitis associated with spinal anaesthesia or other procedures, such as lumbar puncture, during a three week period last month, government epidemiologists said. Four of the patients have died.

Investigations indicate that contaminated syringes and needles might have caused the outbreak, said a report released last week by the epidemiological unit of the Sri Lankan Ministry of Health. Among the 14 cases, eight patients had a recent history of spinal anaesthesia—five mothers who had undergone caesarean sections, a young boy who had undergone lumbar puncture for diagnosis of a fever of unknown origin, and two adults who had received intrathecal chemotherapy.

The report said that a possible source of contamination could have been unsatisfactory storage conditions. "Some imported syringes have been found [to be] contaminated," Athula Kahandalinayage, director general of health services told the *BMJ*. However, the source of contamination is still unknown and detailed investigations are now under way to determine whether this occurred during manufacture, transit, or storage, he said.

Ganapati Mudur *New Delhi*

UK research funding bodies unite against bioterrorism threat

Caroline White *London*

Three of the United Kingdom's largest research funding bodies have announced joint plans to curb the risk of research being misused for bioterrorism.

The Medical Research Council, the Biotechnology and Biological Sciences Research Council, and the Wellcome Trust have issued a joint policy statement announcing changes to their respective policies, guidance, and procedures for awarding grants to researchers.

The move, which is designed to identify at the grant application stage any research that could "conceivably result in harm," and to heighten general awareness of the issues, was prompted by growing public concern about the threat of potential abuse arising from legitimate studies of dangerous pathogens and toxins.

Each agency has already separately issued similar position statements on biomedical research and bioterrorism, covering areas such as funding decisions, international collaboration and training, and public trust.

But the joint statement signals the intention to intensify existing measures, and is the result of wide consultation among key stakeholders across the three agencies. It comes in the wake of increasing scrutiny from bodies, such as the US National Academies of Sciences and the UK Parliamentary Science and Technology Committee.

From January next year, applicants seeking funding from any of these bodies will be asked on their

grant application forms whether they and their institutions have fully assessed the risks of their research being misused in any way, including bioterrorism.

Guidance for funding committees on the risks of research misuse is also being developed, along with protocols on how to deal with cases which spark concerns. Good practice guidelines will include specific reference to the risks of research misuse.

The statement acknowledges that these measures alone will not be able to root out intentional abuse but that strong self governance "will ultimately provide the most effective means of managing the risks of misuse" while also ensuring that "responsibly conducted research is not unnecessarily obstructed."

It urges the scientific community to be proactive, calling for effective local (institutional) research governance, a cautionary approach to making appointments, and compliance with existing health and safety regulations.

Mark Walport, director of the Wellcome Trust, said, "We understand that many people are alarmed by the threat of bioterrorism. But we need to balance these fears with sensible and practical measures that will help us manage such risks." It is vital that scientists pursue legitimate research to understand and treat the many infectious diseases that pose a threat to public health. □

The statement will be on www.mrc.ac.uk, www.wellcome.ac.uk, and www.bbsrc.ac.uk.

Incidents like the anthrax attack in the US in 2001, in which US marines show clean-up techniques, illustrate the threat from terrorists