the compassionate use of an as yet unlicensed product, drug manufacturers should be prepared to supply the drug to the doctors of patients who have been shown to benefit. The costs would, arguably, be no greater than those for an open label study. Safety data would still be available from these patients through the usual monitoring systems.

Future action

Some of our concerns may also apply to other trial designs, but our focus here is on the particular problems of open label extension studies. Several of the problems could be resolved by unblinding the allocation of participants as they complete the phase III study. Investigators object to this on the grounds that it may introduce bias, but participants could be informed of their status in the double blind trial by someone not involved in the analysis of the trial. It has been argued that unblinding should occur even if no open label study follows because the subsequent treatment of participants may be harmed if decisions are made about further management in ignorance of their response to the new drug.⁴

Open label extension studies are currently being misused for marketing purposes or to enable compassionate use of new drug. We recommend that recruitment and consent are dealt with more openly if the study follows a double blind trial and that potential participants are told which arm of the trial they were in

before deciding whether to enrol. Research ethics committees should approve studies only if they address a genuine research question and have a clear rationale, end points, justification of sample size, time scale, and so on. If there is an argument for compassionate use, committees should advise use of mechanisms such as named patient prescriptions rather than open label extension studies.

We thank Susan Pope for providing information from the records of the multicentre ethics committee for Wales and the members of the committee for helping to form and challenge our opinions.

Contributors and sources: PJW was a member of the multicentre research ethics committee for Wales, and GJT is currently a member. GJT is also a member of the Bath local research ethics committee and a council member of the Association of Research Ethics Committees. The idea for the paper came from a discussion between the authors. GJT wrote the initial draft and both authors contributed to the final version. Both authors are guarantors.

Competing interests: None declared.

- 1 Micetich KC. The ethical problems of the open label extension study. *Camb Q Health Ethics* 1996;5:410-4.
- 2 Wainwright P. Consent to open label extension studies: some ethical issues. *J Med Ethics* 2002;28:373-6.
- 3 Council for International Organizations of Medical Sciences. *International ethical guidelines for biomedical research involving human subjects*. Geneva: World Health Organization, 2002.
- 4 Blader JC. Can keeping clinical trial participants blind to their study treatment adversely affect subsequent care? *Contemp Clin Trials* 2005;26: 290-9.

(Accepted 19 June 2005)

bmjlearning.com

The rules

"Dear Editor, I read with interest your reworking of my manuscript. Firstly here are a few things that you did well: you checked the spelling thoroughly and you did improve the grammar. Here's what you could have done better: it would have been better if you hadn't completely changed the tone and style and meaning of my sentences. Yours etc."

Pendleton's rules are rules that help trainers give balanced feedback to trainees.¹ The idea is that, when giving feedback, learners and teachers should concentrate on the positive first and then say what they thought could have been done better. If you've done an advanced cardiac life support course in Britain then you'll be familiar with the rules. In the past many trainers were thought to be too destructive in their criticisms, and Pendleton's rules are an attempt to correct this.

But the rules do have their critics. Many say that they add an *Alice in Wonderland* air to training. I've had a few complaints as an editor and as a clinician, but none of them ever read like the above letter. So why do we encourage doctors to give and receive feedback in a way that they will never experience in real life? It can also be difficult to think of positive things to say to trainees who are still in the low foothills of the learning curve or who, worse still, have turned up to courses without having done any preparatory work. How do you then "accentuate the positive"? Do you say, "Well, at least you turned up"? Perhaps the most annoying thing about the rules is the insistence of some training providers that you must follow them—regardless of the

learning style of the learners or the teaching style of the teacher.

Is there an alternative? Silverman et al have described a new way of giving feedback—called agenda-led, outcomes based analysis.² In this method you start with the learners' agenda and ask them what problems they experienced and what help they would like. Then you look at the outcomes that they are trying to achieve. Next you encourage them to solve the problems and then get the trainer and eventually the whole group involved. Feedback should be descriptive rather than judgmental and should also be balanced and objective.

If you are involved in teaching and learning then you may be interested in finding out more about these concepts. At BMJ Learning we are building up a resource of modules on learning and teaching. You can find out how to run a course, give a lecture, or even give feedback. You can give your views also. If you disagree with Pendleton's rules or with Silverman's ones or would even prefer a return to a 19th century "gloves off" approach to feedback then we are delighted to hear your views and to encourage debate.

Kieran Walsh *clinical editor, BMJ Learning*
(bmjlearning@bmjgroup.com)

- 1 Pendleton D, Scofield T, Tate P, Havelock P. *The consultation: an approach to learning and teaching*. Oxford: Oxford University Press, 1984.
- 2 Silverman JD, Kurtz SM, Draper J. The Calgary-Cambridge approach to communication skills teaching. Agenda-led, outcome-based analysis of the consultation. *Educ Gen Pract* 1996;7:288-99.