- Humphries, E. C. (1956). In *Modern Methods of Plant Analysis*, vol. 1, p. 481. Ed. by Paech, K. & Tracey, M. V. Berlin: Springer-Verlag.
- Iljin, W. S. (1951). *Plant & Soil*, **3**, 339.
- Jacobson, L. (1955). *Plant Physiol.* **30**, 264.
- Jacobson, L. & Ordin, L. (1954). *Plant Physiol.* **29**, 70.
- Krebs, H. A. & Eggleston, L. V. (1943). *Biochem. J.* **37**, 334.
- Laties, G. G. (1961). *Abstr. Commun. 5th int. Congr. Biochem., Moscow*, p. 324.
- Luggs, J. W. H. & Overell, B. T. (1948). *Aust. J. sci. Res. A*, **1**, 98.
- MacRobbie, E. A. C. & Dainty, J. (1958). *J. gen. Physiol.* **42**, 335.
- Pucher, G. W. & Vickery, H. B. (1949). *J. biol. Chem.* **178**, 557.
- Rasmussen, G. K. & Smith, P. F. (1961). *Plant Physiol.* **36**, 99.
- Roe, J. H. (1954). *Meth. biochem. Anal.* **7**, 118.
- Scott, R. O. & Ure, A. M. (1958). *Analyst*, **83**, 561.
- Sinclair, W. B. & Eny, D. M. (1947). *Plant Physiol.* **22**, 257.
- Taylor, T. G. (1953). *Biochem. J.* **54**, 48.
- Ure, A. M. (1958). *Brit. Commun. & Electronics*, **5**, 846.
- Vickery, H. B. & Wilson, D. G. (1958). *J. biol. Chem.* **233**, 14.

Biochem. J. (1963) **86**, 494

The Aerobic Inhibition of Glycolysis Associated with Brain Mitochondria

By R. BALÁZS

Neuropsychiatric Research Unit, Medical Research Council, Woodmansterne Road, Carshalton, Surrey

(Received 9 August 1962)

In recent years several attempts have been made to clarify the mechanism of the aerobic control of glycolysis by using cell-free preparations (Terner, 1954; Graffi & Schneider, 1956; Aisenberg, Reinafarje & Potter, 1957; Aisenberg & Potter, 1957; Von Korff & Twedt, 1957; Tiedemann & Born, 1958; Sisakian & Pinus, 1958; Von Korff, 1959; Gatt & Racker, 1959; Cremer, 1960; Schwartz & Lee, 1960; Tiedemann & Born, 1960). Only in some of these studies was evidence obtained of an interaction between respiration and glycolysis, as implied by the so-called 'Pasteur effect'.

In previous work from this Laboratory it was confirmed that about 10% of the glycolytic activity of whole-brain homogenates prepared with a blender is associated with the mitochondrial fraction obtained by the usual differential-centrifuging procedures. The aerobic inhibition and the site of inhibition of lactate formation in the glycolytic chain were also studied with this preparation (Balázs & Lagnado, 1959; Balázs, 1959).

In this paper results are given of an investigation of the interaction of glycolysis with respiration; experiments on the effects of respiratory inhibitors and uncouplers of oxidative phosphorylation are also described. It is concluded that more than one mechanism is operative in the aerobic control of glycolysis associated with brain-mitochondrial preparations. One mechanism involves a specific aerobic inhibition of glyceraldehyde phosphate dehydrogenase; another mechanism appears to be closely connected with the processes of oxidative phosphorylation. Part of this investigation has been published (Balázs & Richter, 1960a).

EXPERIMENTAL

Abbreviation. BAL, 2,3-Dimercaptopropanol.

Preparation of brain-mitochondrial fraction of rat brain. The method previously described (Balázs, 1959; Balázs & Lagnado, 1959) was followed. The two hemispheres were homogenized in a NIRD blender (Nelco Ltd., Shalford, Surrey), in 2 vol. of 0.25 M-sucrose containing 0.025 M-nicotinamide, for 45 sec. at 10 000 rev./min. The sucrose solution was then added to make a 10% homogenate and the suspension was homogenized for a further 30 sec. at the same speed. The nuclear fraction was sedimented at 1500g for 10 min. in a MSE 20 refrigerated centrifuge. The supernatant was centrifuged again at the same speed and time. The mitochondrial fraction was collected and washed once at 18 000g for 10 min. The final pellet was suspended in the nicotinamide-containing sucrose solution.

Media. The basic medium was prepared freshly from stock solutions and contained 6 mM-MgCl₂, 2 mM-EDTA (Na salt, pH 7.2), 25 mM-nicotinamide, 10 mM-KH₂PO₄, 20 mM-tris (adjusted to pH 7.2 with HCl) and sufficient KCl to bring the osmolarity to 0.3 after the addition of all the constituents to the final incubation medium. The pH of the basic medium was adjusted to 7.2 after the addition of the substrates. The pH of the medium was maintained generally by prior adjustment of the pH of the additives when necessary. A few compounds were used in alkaline solutions: sodium 5-ethylamylbarbiturate (Amytal), 2-n-heptyl-4-hydroxyquinoline N-oxide and dicoumarol. In these cases the pH of the basic medium was adjusted to 7.2 after the addition of these substances. The final volume in the main compartment of the Warburg flask in the absence and presence of the cofactors (2 mM-ADP and 0.33 mM-NAD⁺) was 1.5 and 2.0 ml. respectively.

Special chemicals and reagents. Sodium Amytal was a gift from Eli Lilly and Co. Ltd. Dicoumarol was a gift from Organon Laboratories Ltd. 2-n-Heptyl-4-hydroxyquin-

oline *N*-oxide was a gift from Dr J. W. Lightbown (National Institute for Medical Research, Mill Hill). This compound was dissolved in 1 mM-KOH and the concentration was determined with a Hilger Uvispek spectrophotometer by using the molar extinction coefficient at 346 m μ reported by Cornforth & James (1956). Antimycin A (Wisconsin Aluminium Research Foundation) was dissolved either in ethanol or dimethylformamide. Some experiments with this substance were carried out with a sample kindly supplied by Dr F. A. Holton (Medical Research Council, Hammersmith Hospital).

Crystalline glyceraldehyde phosphate dehydrogenase was prepared by the method of Cori, Slein & Cori (1948). Tris (0.03M) was used instead of pyrophosphate in the assay of the enzyme and the reaction was carried out at pH 7.2. The enzyme was six times recrystallized. Before use the enzyme suspension was centrifuged, the crystals were dissolved in 1 mM-EDTA, pH 8, and the solution was dialysed for 5 hr. at pH 8 against five changes of 1 mM-EDTA. The preparation had an activity of 22 μ moles of NADH formed/min./mg. of protein at 18°. The protein was determined by the method of Cori *et al.* (1948).

Pyruvic acid (AnalaR, British Drug Houses Ltd.) was redistilled and the sodium salt prepared by the method of Price & Levintow (1952):

The following chemicals were obtained from sources indicated: cytochrome *c*, α -oxoglutaric acid and tris (L. Light and Co. Ltd.); sodium salt of ADP, NAD⁺ and DL-glycerol 1-phosphate (sodium salt) (Sigma Chemical Co.); nicotinamide, ascorbic acid, L-cysteine hydrochloride, glutathione, BAL, fructose 1,6-diphosphate (tetrasodium salt), KOH, KCl, MgCl₂, KH₂PO₄, Na₂HAsO₄, NaF and sucrose (British Drug Houses Ltd.); EDTA, succinic acid, KCN and NaCN₃ (Hopkin and Williams Ltd.); solution of sodium salts of triose phosphoric acids, phosphoglycerate kinase, glycerol phosphate dehydrogenase and lactate dehydrogenase (C. F. Boehringer und Soehne, Mannheim, Germany).

Dimethylformamide was used for dissolving BAL and antimycin A. The solvent did not affect the metabolic processes under investigation, when used in 0.4% (v/v) final concentration.

Incubation and manometric technique. Incubation was carried out in conventional Warburg vessels at 37.5°. In the anaerobic experiments the vessels were gassed with N₂ (freed from traces of O₂ by passage through a heated copper tube) and the aerobic experiments were carried out in air. When consumption of O₂ was measured (by the Warburg technique; Umbreit, Burris & Stauffer, 1945) the centre well contained 0.1 ml. of 20% (w/v) NaOH. Temperature equilibration was carried out with open stopcocks for 6–10 min. and results were extrapolated for this preincubation period. Unless otherwise indicated the reaction was stopped after incubation for 40–60 min. by trichloroacetic acid, HClO₄ or Ba(OH)₂ according to the substances to be measured. The content of metabolites was measured usually in a zero-time control and the change in concentration was calculated by subtracting these values from the values obtained after incubation.

In the experiments with cyanide and azide the alkali was omitted from the centre well. Since cyanide and hydrazoic acid are volatile (Umbreit *et al.* 1945), the anaerobic vessels were gassed for 6.5 min. and the cyanide or azide was injected with an Agla microsyringe into the

vessels through a rubber tube that was fixed to the side arm. Then incubation was continued for the required length of time. Controls were incubated in air and N₂ for 7 min. and the value of the content of measured metabolites was subtracted from the experimental value.

In the experiments with glutathione and cysteine, these compounds were dissolved and the pH was adjusted to 7.2 just before incubation was started. They were added to the medium immediately before the addition of the mitochondrial suspension.

Analytical methods. Lactate was determined from perchloric acid- or trichloroacetic acid-treated filtrates (final concn. 3 and 5%, w/v, respectively) by the method of Barker & Summerson (1941) as described by Balázs (1959). Glucose was measured after Zn–Ba precipitation (Somogyi, 1945) with the reducing reagent of Somogyi (1952) by the method of Nelson (1944). Inorganic phosphate was measured by the method of Lohmann & Jendrassik (1926). Alkali-labile phosphate (in this system the sum of fructose diphosphate and triose phosphates) was determined by the method of Kiessling (1938). Protein was determined by the method of Lowry, Rosebrough, Farr & Randall (1951). Pyruvate was measured by the method of Friedemann & Haugen (1943). From one portion the 2,4-dinitrophenylhydrazones of the keto acids were extracted for 5 min. with benzene and from another for 25 min. with ethyl acetate. The extraction was accomplished with a mechanical mixer (Matburn Ltd.). Regular measurements of *E* at 435 m μ were made with a Unicam SP. 600 spectrophotometer.

The colorimetric determination of pyruvate in the samples was checked by the enzymic method of Segal, Blair & Wyngaarden (1956). From 18 determinations the quantity of pyruvate measured by the enzymic method was 98.16 $\pm$ 1.01% of that determined by the colorimetric method. When metabolites were determined by enzymic methods the samples were precipitated with HClO₄ (3%, w/v, final concn.). The pH of the perchloric acid-treated filtrates was adjusted to about 6 with KOH. The filtrates were kept in an ice–water bath for 30 min. and potassium perchlorate was removed by centrifuging in the cold. A portion of the supernatant was used in the assays.

L-Glycerol 1-phosphate and 3-phosphoglycerate were determined by the method of Hohorst, Kreutz & Bücher (1959) with slight modifications. The formation and disappearance respectively of NADH was followed at 340 m μ with a Hilger Uvispek or with the Unicam SP. 700 recording spectrophotometer (light-path 1 cm.). The final volume in each case was 2.8 ml. The molecular extinction coefficient (*e*) of NADH at 340 m μ was assumed to be 6.22 $\times$ 10³. The medium for L-glycerol 1-phosphate determination contained 0.07 M-hydrazine sulphate, 0.18 M-glycine, 0.9 mM-EDTA (pH of a solution containing these three substances was adjusted before use to 9.5) and 1.2 mM-NAD⁺. L-Glycerol 1-phosphate was determined from ΔE obtained after the addition of glycerol phosphate dehydrogenase (100 'Bücher units'; for definition see Beizenherz *et al.* 1953) and incubation at 30° for 40 min.

3-Phosphoglycerate was determined in a medium containing 0.037 M-tris–HCl buffer, pH 8, 5 mM-MgCl₂, 2.5 mM-glutathione, 1 mM-EDTA, 2.5 mM-hydrazine sulphate, 7 mM-ATP, 0.1 mM-NADH. Before use a mixture of tris–HCl, MgCl₂, EDTA and hydrazine sulphate was prepared from stock solutions and adjusted to pH 8. Glutathione was freshly dissolved in 0.1 M-tris–HCl buffer,

Table 1. *Aerobic inhibition of glycolysis in rat-brain-mitochondrial systems*

The basic medium contained: 6 mm-MgCl₂, 10 mm-KH₂PO₄, 20 mm-tris-HCl (pH 7.2), 2 mm-EDTA, 25 mm-nicotinamide; the pH was adjusted with KOH to 7.2 and KCl was added to bring the osmolarity to 0.3. Substrates and cofactors were added to this medium: 4-7 mm-pyruvate, 2-7 mm-glucose, 10 mm-fructose diphosphate, 2 mm-ADP, 0.33 mm-NAD⁺. The volume in the absence and presence of cofactors was 1.5 and 2.0 ml. respectively. All concentrations refer to the final concentrations in the flasks. Gassing with N₂ was for 7 min. Incubation: 40-60 min. at 37.5°. Figures represent the means $\pm$ s.e. with numbers of experiments in parentheses. Oxygen uptakes and substrate changes are given as μ moles/hr./100 mg. of protein. The gas phase in this and the other Tables is indicated by 'Air' or 'N₂' in the column headings.

Substrates added	Cofactors (ADP, NAD ⁺) added	Uptake of O ₂	Lactate formed		Lactate formed (corrected*)		Aerobic inhibition of lactate production† (%)	Pyruvate consumed		Glucose consumed		Alkali-labile phosphate consumed	
			Air	N ₂	Air	N ₂		Air	N ₂	Air	N ₂	Air	N ₂
Pyruvate	-	49.3 $\pm$ 1.35 (9)	1.3 $\pm$ 0.47 (9)	3.3 $\pm$ 0.69 (9)	.	.	.	39.5 $\pm$ 2.48 (9)	12.2 $\pm$ 1.91 (9)	.	.	.	.
Pyruvate + glucose	-	46.8 $\pm$ 0.40 (8)	10.5 $\pm$ 0.73 (8)	44.6 $\pm$ 0.07 (8)	9.2 $\pm$ 0.65 (8)	41.2 $\pm$ 2.18 (8)	78 $\pm$ 1.25 (8)	42.8 $\pm$ 0.07 (8)	13.8 $\pm$ 1.13 (8)	6.6 $\pm$ 1.43 (8)	25.3 $\pm$ 0.91 (8)	.	.
Pyruvate	+	104.2 $\pm$ 6.73 (5)	1.7 $\pm$ 0.81 (4)	5.3 $\pm$ 0.71 (5)	.	.	.	117.3 $\pm$ 11.06 (4)	18.5 $\pm$ 4.40 (5)	.	.	.	.
Pyruvate + glucose	+	110.4 (2)	75.1 $\pm$ 4.35 (3)	118.7 $\pm$ 10.39 (5)	74.0 $\pm$ 4.69 (3)	116.6 $\pm$ 7.87 (3)	37 $\pm$ 4.19 (3)	91.7 (2)	9.9 $\pm$ 2.5 (4)	94.5 (2)	77.7 $\pm$ 8.94 (5)	.	.
Pyruvate + fructose diphosphate	+	102.1 $\pm$ 7.48 (4)	153.9 $\pm$ 20.54 (4)	271.1 $\pm$ 13.34 (4)	152.2 $\pm$ 12.73 (4)	265.8 $\pm$ 20.25 (4)	43 $\pm$ 5.1 (4)	17.2 $\pm$ 2.22 (4)	12.1 $\pm$ 4.0 (4)	.	.	165.6 $\pm$ 14 (3)	314.3 $\pm$ 34 (3)

* Correction made for lactate formation in the absence of glycolytic substrates.

† Percentage inhibition of aerobic lactate production: 100 (lactate_{N₂} - lactate_{air})/lactate_{N₂}.

pH 8; ATP was adjusted to pH 7.5. The reaction was carried out in the Unicam SP. 700 recording instrument and was initiated by the addition of glyceraldehyde phosphate dehydrogenase (150 'Bücher units') and phosphoglycerate kinase (300 units). In both enzymic methods no interference could be detected when the other compounds present in the samples were added at concentrations five times as high as those normally used in the experiments.

RESULTS

Contribution of dismutation of pyruvate to the formation of lactate. Lactate formation by the mitochondrial fraction of rat brain from glucose or fructose diphosphate was measured in the present experiments in air and N_2 . Since the incubation media contained pyruvate it was necessary to find whether the lactate formed was due to glycolysis or to the dismutation of pyruvate (Krebs & Johnson, 1937; Weil-Malherbe, 1937). The magnitude of the dismutation of pyruvate was obtained by measuring the lactate produced and pyruvate consumed in the absence of glycolytic substrates. Furthermore, pyruvate was also determined when glucose or fructose diphosphate was used as substrate. Table 1 summarizes these results. Lactate formation from pyruvate was very small in all the conditions tested when compared with the production of lactate from glycolytic substrates. When measured in N_2 in the absence of cofactors the lactate formed from pyruvate was only 7% of the production of lactate in the presence of glucose; in the presence of cofactors this was about 5%. Moreover the lactate produced from pyruvate alone was only 2% of that formed from pyruvate and fructose diphosphate together in the presence of ADP and NAD^+ . The pyruvate consumption in N_2 was also low, although bigger than the theoretical value (2 moles of pyruvate consumed/mole of lactate formed) for the dismutation reaction. This was found to be due mainly to the oxidation of pyruvate during the period until anoxia was established. In a few experiments controls were incubated and gassed for 7 min. and the value for the pyruvate and lactate content was subtracted from the experimental value. The relation of lactate formed to pyruvate consumed was very near to the theoretical (Table 4). The dismutation of pyruvate in the presence of respiratory inhibitors was found to be similar to that observed in N_2 . The uptake of pyruvate in the presence of glycolytic substrates in N_2 or in the presence of respiratory inhibitors did not indicate an increase in the dismutation of pyruvate to lactate. These results give evidence that the lactate formation from glucose and fructose diphosphate was mainly due to glycolytic reactions.

Rate of glycolysis and the aerobic inhibition of lactate formation under different experimental condi-

tions. The rate of lactate formation was found to be nearly linear for 1 hr. at 37.5° under all experimental conditions tested (Figs. 2, 3). The addition of ADP and NAD^+ increased lactate production from glucose. The lactate formation was even higher when fructose diphosphate was the substrate in the presence of cofactors. The relative rates in N_2 under these conditions were 100, 266 and 608% respectively. The aerobic inhibition of lactate formation was greatest in the absence of cofactors (78%); it was about 40% when ADP and NAD^+ were added to the medium (Table 1).

An aerobic inhibition of glucose consumption was obtained only in the absence of added cofactors. The magnitude of this inhibition agreed closely with the observed inhibition of lactate formation. The uptake of pyruvate in the presence of fructose diphosphate in air was much smaller than in the controls that did not contain fructose diphosphate. It was found that the fructose diphosphate used contained a small amount of impurity (mainly phosphoglycerate), which produced pyruvate during incubation.

Effect of respiratory inhibitors on glycolysis in the absence of ADP and NAD^+ . The nature of the interaction of respiration with glycolysis was further investigated in a system containing glucose without added cofactors. It was found that respiratory inhibitors acting at several different points in the respiratory chain were able to reduce substantially the aerobic inhibition of glycolysis in this system (Table 2). There was some uptake of O_2 that could not be depressed by increasing the concentrations of the inhibitors. However, the aerobic inhibition of glycolysis was at most very slight in the presence of this residual electron transport.

Part of the electron-transport chain concerned in aerobic control of glycolysis. In these experiments the respiratory chain was blocked at certain points and electrons were fed in by suitable electron donors.

(a) Effect of restoration of electron transport between cytochrome *c* and O_2 . Antimycin A inhibits electron transport between cytochrome *b* and *c* (Potter & Reif, 1952). When cytochrome *c* and ascorbic acid are added, electron transport occurs between cytochrome *c* and O_2 . With the amount of mitochondria used in these experiments (3–6 mg. of protein/vessel) maximal inhibition of respiration was obtained with as little as 0.5 μ g. of antimycin A/flask. In these experiments 1–2 μ g. of antimycin A was used/vessel. An equivalent amount of the solvent (ethanol or dimethylformamide) was added to the control vessels.

Antimycin A increased the aerobic lactate formation and glucose consumption to about the anaerobic value. The further addition of cytochrome *c* and ascorbate produced electron transport, but

aerobic glycolysis was not markedly inhibited under these conditions (Table 3).

(b) Effect of the restoration of the electron transport between cytochrome *b* and O_2 . It is possible to block the respiratory chain between NADH and flavoprotein with Amytal (Ernster, Jalling, Löw & Lindberg, 1955), and electron transport can occur between cytochrome *b* and O_2 when succinate is added to the inhibited system.

In preliminary experiments it was found that the half-maximal effect of Amytal in increasing aerobic glycolysis was produced at a concentration approximately 0.6 mM. In the experiments reported here 1.8 mM-Amytal was used: this concentration produced maximal inhibition of NAD^+ -linked respiration and maximal increase in aerobic glycolysis.

Lactate formation from pyruvate in the absence of glucose was increased by Amytal to the anaerobic value (Table 4A). When succinate was added together with pyruvate the aerobic lactate formation was greater than the anaerobic. The lactate formation in air from pyruvate was further increased by Amytal (Table 4B). In the presence of succinate and Amytal the ratio of lactate formed to pyruvate used in air was greater than the theoretical value that would be expected for the dismutation of pyruvate. This indicates that in this case only a relatively small part of the lactate obtained was formed by a dismutation of pyruvate. It is known (Chance & Hollunger, 1957; Birt & Bartley, 1960) that succinate is able to increase the amount of mitochondrial NADH: it is possible that some reduction of pyruvate was produced in this way.

Table 2. *Effect of respiratory inhibitors on the aerobic control of glycolysis in the absence of added cofactors*

Basic medium; 4.5 mM-pyruvate, 2.5 mM-glucose, mitochondria, 4.3–4.8 mg. of protein/vessel. Final volume, 1.5 ml. Incubation: 60 min. at 37.5°. Values for lactate represent the difference between the amount of lactate formed in the presence and absence of glucose. Oxygen uptakes and changes in metabolites are given as μ moles/hr./100 mg. of protein.

Expt. no.	Nature and concn. of respiratory inhibitor added	Uptake of O_2	Lactate formation		Aerobic inhibition of lactate production (%)	Glucose uptake		Aerobic inhibition of glucose uptake (%)
			Air	N_2		Air	N_2	
1	None	41.7	11.8	49.4	76	5.8	26	78
	Antimycin A (0.0023 mM)	26.4	41.3	45.8	10	20.8	24.6	15
2	None	51.7	8.3	37.7	78	5.3	29.1	82
	Amytal (1.8 mM)	12.6	26.6	29.1	9	14.9	15.0	1
3	None	49.0	11.3	40.2	72	6.1	24.8	75
	KCN (mM)	.	34.3	39.7	13	23.8	26.2	9
	NaN_3 (mM)	.	37.9	39.2	3	20.6	20.6	0
4	None	41.4	11.7	47.8	76	.	.	.
	2- <i>n</i> -Heptyl-4-hydroxyquinoline <i>N</i> -oxide (0.004 mM)	37	19.4	47.0	59	.	.	.
	(0.016 mM)	21.6	44.1	47.0	6	.	.	.

Table 3. *Effect of antimycin A and restoration of electron transport on the aerobic inhibition of glycolysis*

Basic medium; 4.35 mM-pyruvate, 2.5 mM-glucose, mitochondria, 4.7 mg. of protein/vessel; 2.3 μ M-antimycin A, 0.1 mM-cytochrome *c* and 50 mM-ascorbic acid, when added. Final volume, 1.5 ml. Incubation: 60 min. at 37.5°. Oxygen uptake and substrate changes are given as μ moles/hr./vessel.

Substrates	Additions (mM)	Uptake of O_2	Lactate formation		Glucose uptake		Pyruvate uptake	
			Air	N_2	Air	N_2	Air	N_2
Pyruvate + glucose	None	2.50	0.56	2.49	0.27	1.23	2.34	0.72
	Antimycin A	1.25	2.04	2.22	0.99	1.17	1.16	0.61
	Cytochrome <i>c</i> + ascorbic acid	5.99	0.45	2.0	.	.	1.84	0.74
	Antimycin A + cytochrome <i>c</i> + ascorbic acid	5.76	1.85	2.26	.	.	1.44	0.56
Pyruvate	None	2.50	0	0.15	.	.	2.46	0.81
	Antimycin A	1.10	0.09	0.10	.	.	1.16	0.51
	Cytochrome <i>c</i> + ascorbic acid	6.03	0.02	0.09	.	.	1.94	0.69
	Antimycin A + cytochrome <i>c</i> + ascorbic acid	5.16	0.11	0.13	.	.	1.56	0.59

Table 4. *Effect of Amytal on the aerobic control of glycolysis in the presence and absence of succinate*

Basic medium; 4 mm-pyruvate, 2.6 mm-glucose, 4.6–4.8 mg. of mitochondrial protein/vessel; 10 mm-succinate and 1.8 mm-Amytal, when added. (A) and (B) are the means of two experiments each. Changes in the concentration of the metabolites were determined from the difference in the concentrations reached between 7 min. and 1 hr. of incubation. Oxygen uptakes and substrate changes are given as $\mu\text{moles/hr./100 mg. of protein.}$

Additions	Uptake of O ₂	Lactate formation		Lactate formation (corrected*)		Pyruvate uptake		Glucose uptake	
		Air	N ₂	Air	N ₂	Air	N ₂	Air	N ₂
(A) Pyruvate + glucose	48.32	11.18	43.97	7.98	37.72	46.22	13.98	5.64	26.53
Pyruvate	49.60	3.20	6.25	.	.	40.23	12.68	.	.
Pyruvate + glucose + Amytal	14.09	32.69	34.40	25.76	27.98	13.41	8.88	19.46	18.90
Pyruvate + Amytal	12.14	6.93	6.42	.	.	11.04	9.39	.	.
(B) Pyruvate + glucose + succinate	199.45	12.18	54.16	6.37	50.08	40.03	11.61	5.99	32.75
Pyruvate + succinate	205.96	5.81	4.08	.	.	44.61	9.40	.	.
Pyruvate + glucose + Amytal + succinate	160.83	50.67	49.33	40.55	44.59	8.94	6.06	28.40	26.81
Pyruvate + Amytal + succinate	154.19	10.12	4.74	.	.	10.85	10.05	.	.

* Correction made for lactate formation in the absence of glucose.

Succinate was able to produce electron transport in the Amytal-blocked system. However, the aerobic inhibition of glycolysis was not restored by succinate.

Effect of uncouplers of oxidative phosphorylation on aerobic glycolysis. It is known that in tissue preparations containing whole cells uncouplers of oxidative phosphorylation abolish the Pasteur reaction (Dodds & Greville, 1934; Lynen & Koenigsberger, 1951). Expts. 1 and 2 of Table 5 show the effect of dinitrophenol and dicoumarol in the absence of cofactors. The rate of respiration was more than doubled and the aerobic inhibition of glycolysis was completely abolished. It was a reproducible finding that in the presence of these uncouplers of phosphorylation the glucose uptake in air was greater than in N₂.

When dinitrophenol or dicoumarol was added to a system containing ADP and NAD⁺ the aerobic lactate formation approached the anaerobic value (Expts. 3 and 4 of Table 5).

Effect of succinate in an Amytal-blocked system on aerobic glycolysis in the presence of added ADP and NAD⁺. In control studies it was confirmed that oxidative phosphorylation with succinate as substrate in the presence of cofactors was hardly affected by Amytal. Although the electron transport and coupled phosphorylation between cytochrome *b* and O₂ proceeded in the presence of succinate and Amytal, the aerobic control of glycolysis was not restored (Table 6). Thus under these conditions the processes of oxidative phosphorylation did not exert any regulatory effect on the rate of glycolysis.

By comparing the data in Tables 4 and 6 it can be seen that the uptake of O₂ with pyruvate and

succinate as substrates was much more inhibited by Amytal in the presence than in the absence of ADP and NAD⁺. This is due to the fact that, in the presence of cofactors, the contribution of NAD⁺-linked pyruvate oxidation to the total uptake of O₂ is much greater than it is in their absence (pyruvate uptake 140 and 40 $\mu\text{moles/hr./100 mg. of protein respectively}$).

Aerobic inhibition of glycolysis with fructose diphosphate as substrate in the presence of ADP and NAD⁺. Under these experimental conditions, the aerobic inhibition of glycolysis was about 40 % and the dismutation of pyruvate to lactate was about 2 % of the total lactate production (cf. Table 1).

It was shown by Balázs (1959) and confirmed in the present experiments that the aerobic inhibition of lactate formation in this system is due to an inhibition of the enzyme glyceraldehyde phosphate dehydrogenase. When enolase was inhibited by fluoride, and the oxidoreductive step of the glycolysis was shifted towards the oxidation of glyceraldehyde phosphate by addition of arsenate, the amount of phosphoglycerate formed accounted for the amount of glyceraldehyde phosphate oxidized. The amount of lactate formed accounted for the amount of NADH that was used for pyruvate reduction. Table 7 shows that, in the presence of fluoride or fluoride and arsenate, less phosphoglycerate is formed in air than in N₂, i.e. the rate of the glyceraldehyde phosphate-dehydrogenase reaction is lower in air than in N₂; the percentage inhibition was similar to that obtained in lactate formation.

Effect of added crystalline glyceraldehyde phosphate dehydrogenase. Direct evidence for the inhibition of glyceraldehyde phosphate-dehydrogenase

Table 5. *Effect of uncouplers of phosphorylation on the aerobic inhibition of lactate formation*

Basic medium; substrates as indicated; 2 mm-ADP and 0.33 mm-NAD⁺, when added. Mitochondria were added as mg. wet wt. of tissue: Expt. 1, 280 (6.3 mg. of protein); Expt. 2, 270; Expt. 3, 250 (6.4 mg. of protein); Expt. 4, 120. Incubation volume, in the presence and absence of cofactors, 1.5 and 2 ml. respectively. Incubation: 60 min., except in Expt. 3 (40 min., results calculated to 60 min.). Oxygen uptakes and changes in metabolites are given as μ moles/hr./vessel.

Expt. no.	Nature and concn. of substrates added	Cofactors (ADP, NAD ⁺) added	Nature and concn. of uncoupler added	Uptake of O ₂	Lactate formation		Pyruvate uptake		Glucose uptake	
					Air	N ₂	Air	N ₂	Air	N ₂
1	Pyruvate (4.5 mm) + glucose (2.6 mm)	-	None	2.9	0.64	2.96	2.52	0.77	0.40	1.70
	Pyruvate (4.5 mm)	-	Dinitrophenol (0.1 mm)	6.85	2.70	2.54	4.27	0.33	1.84	1.29
	Pyruvate (4.5 mm)	-	None	2.9	0.06	0.19	2.45	0.78	.	.
2	Pyruvate (6.6 mm) + glucose (2.5 mm)	-	Dinitrophenol (0.1 mm)	6.7	0.06	0.19	4.01	0.32	.	.
		-	None	2.64	0.89	3.03	.	.	0.49	1.59
		-	Dinitrophenol (0.1 mm)	7.40	2.94	2.76	.	.	1.89	1.36
3	Pyruvate (7.0 mm) + L-malate (3.0 mm) + glucose (2.7 mm)	+	Dicummarol (0.02 mm)	7.98	2.53	2.53	.	.	1.69	1.29
	Pyruvate (7.0 mm) + L-malate (3.0 mm)	+	None	24.24	6.03	9.99	13.39	1.53	7.5	6.54
		+	Dinitrophenol (0.1 mm)	28.88	9.23	9.33	9.22	0.92	7.37	6.33
4	Pyruvate (6.6 mm) + α -oxoglutarate (10 mm) + glucose (3.2 mm)	+	None	18.68	0.20	1.32	12.36	2.80	.	.
		+	Dinitrophenol (0.1 mm)	20.08	0.45	1.04	8.13	1.38	.	.
		+	None	11.40	1.98	3.06	.	.	4.58	2.10
		+	Dinitrophenol (0.05 mm)	14.40	2.88	3.06	.	.	4.48	1.88
		+	Dinitrophenol (0.1 mm)	11.70	2.61	2.61	.	.	2.51	1.79

activity in air was obtained by observing the effect of the addition of an excess of this enzyme to the system (Table 8). The addition of the crystalline enzyme had no effect on the dismutation of pyruvate, but under these conditions the aerobic lactate formation from fructose diphosphate approached the anaerobic value. The difference between lactate formation in air and N₂ was 4.9 μ -moles/vessel in the control, but in the presence of added enzyme the difference was only about 1 μ mole. Glyceraldehyde phosphate dehydrogenase also increased lactate formation in N₂. This was to be expected since in the presence of fructose diphosphate, pyruvate, ADP and NAD⁺ the rate-limiting reaction was the oxidation of glyceraldehyde phosphate (Balázs & Lagnado, 1959). Phosphoglycerate formation and pyruvate consumption were also increased by adding the enzyme. This indicated that rate-limiting steps in the glycolytic chain between these substances were bypassed, and lactate was formed at a rate determined only by the operation of glyceraldehyde phosphate dehydrogenase and lactate dehydrogenase. With glucose as substrate in the absence of cofactors the addition of glyceraldehyde phosphate dehydrogenase did not affect the aerobic control of glycolysis. Thus these experiments indicated that the aerobic inhibition of glycolysis observed in the fortified and unfortified systems were due to different mechanisms. The following experiments give further evidence on this point.

Effect of respiratory inhibitors and uncouplers of phosphorylation. In previous experiments (Tables 2 and 5) it was shown that respiratory inhibitors and uncouplers were able to remove the aerobic control of glycolysis in systems where glucose was the glycolytic substrate. However, these compounds had no effect when fructose diphosphate was the substrate in a fortified system containing added ADP and NAD⁺ (Table 9).

Effect of monothiols and BAL. Glyceraldehyde phosphate dehydrogenase prepared from different sources is known to be a SH enzyme. Hatch & Turner (1959) reported that the aerobic inhibition of lactate formation in pea-seed extracts was due to the inhibition of this enzyme, which could be protected by monothiols. However, glutathione or cysteine had almost no effect in the brain-mitochondrial system (Expt. 2, Table 10). On the other hand, 2 mm-BAL raised the aerobic glycolysis to the value obtained in N₂ (Expt. 1, Table 10). Similar results were obtained when the effect of monothiols and BAL was tested in an assay system for glyceraldehyde phosphate dehydrogenase in the presence of fluoride and arsenate with fructose diphosphate as substrate in a medium fortified with cofactors. The half-maximal protective concentration of BAL was found to be

Table 6. *Effect of Amytal on the aerobic inhibition of lactate formation in the presence of ADP and NAD⁺*

Basic medium; 7 mM-pyruvate, 10 mM-succinate, 2 mM-ADP, 0.33 mM-NAD⁺; mitochondria, 6.5 mg. of protein/vessel. Incubation volume, 2.0 ml. Incubation: 40 min. Glucose 2.7 mM and Amytal 1.8 mM, when added. Oxygen uptakes and changes in metabolites are given as μ moles/vessel.

Additions	Uptake of O ₂	Lactate formation		Lactate formation (corrected*)		Pyruvate uptake		Glucose uptake	
		Air	N ₂	Air	N ₂	Air	N ₂	Air	N ₂
Glucose	20.30	4.0	5.97	3.38	5.51	6.08	0.67	4.89	4.36
Glucose + Amytal	7.90	6.20	6.10	5.39	5.42	0	0.47	4.89	4.32
None	17.5	0.62	0.46	.	.	6.31	1.22	.	.
Amytal	7.85	0.81	0.68	.	.	1.22	0.93	.	.

* Difference in lactate formation in the presence and absence of glucose.

about 0.3 mM (Fig. 1). BAL had no effect on the aerobic inhibition of glycolysis in a medium containing glucose as glycolytic substrate in the absence of added cofactors (Expt. 3, Table 10).

Joint operation of the two inhibitory mechanisms of the aerobic control of glycolysis. The two inhibitory mechanisms could be studied together in a system that contained fructose diphosphate and glucose as glycolytic substrates in the presence of ADP and NAD⁺. Addition of glucose to a system containing fructose diphosphate and cofactors produced an increase in the anaerobic lactate formation, whereas glycolysis was not affected in air. When dinitrophenol and glyceraldehyde phosphate dehydrogenase were added separately the aerobic inhibition of lactate formation was decreased (Table 11). The combined addition of dinitrophenol and the enzyme produced comparable glycolysis in air and in N₂.

Reversibility of aerobic inhibition of glycolysis by changing the gas phase. Fig. 2 shows an experiment in a system containing glucose in the absence of added cofactors when the gas phases were reversed after incubation for 30 min. The rate of glycolysis changed according to the phase, whether from air to N₂ or the reverse. This also applied to the systems where fructose diphosphate was the glycolytic substrate in presence of ADP and NAD⁺. Fig. 3 shows the reversibility of the aerobic control of lactate formation in such a system in the presence of fluoride and arsenate.

DISCUSSION

The heterogeneous composition of the brain-mitochondrial fraction is now generally recognized (Beaufay, Berleur & Doyen, 1957; Petruschkha & Giuditta, 1959; Whittaker, 1959; Aldridge & Johnson, 1959). That glycolytic activity is obtained in this fraction does not necessarily imply that all the enzymes involved in the conversion of glucose into lactate are localized in the mitochondrial particles (Balázs & Lagnado, 1959; Johnson, 1960).

Fig. 1. Effect of BAL on the aerobic inhibition of glycolysis. Results are the means of two experiments. In the absence of BAL, lactate formation in N₂ was 256 μ moles/hr./100 mg. of protein. Incubation mixture contained the basic medium, 10 mM-fructose diphosphate, 6.6 mM-pyruvate, 10 mM- α -oxoglutarate, 2 mM-ADP and 0.33 mM-NAD⁺.

Fig. 2. Demonstration of reversal of the aerobic control of glycolysis by changing the gas phase. Basic medium, 2.53 mM-glucose, 6.6 mM-pyruvate; no cofactors were added. Mitochondria added: 6.5 mg. of protein/vessel. Oxygen consumption 2.78 μ moles/hr. Gas phase: ○—○, air; ●—●, N₂. Change of gas phase was made at arrow. Lactate values were corrected for the formation of lactate in the absence of glucose.

Table 7. *Aerobic inhibition of glycolysis with fructose diphosphate as substrate in the presence of ADP and NAD⁺*

Basic medium; 0.62 mM-3-phosphoglycerate was introduced with the fructose diphosphate added; 2 mM-ADP, 0.33 mM-NAD⁺; where arsenate was added (pH adjusted to 7.2) the concentration of inorganic phosphate was decreased from 10 to 3 mM. Mitochondria added: 2.9 mg. of protein/vessel. Final concentrations: fructose diphosphate 12.5 mM, pyruvate 9.37 mM, NaF 10 mM, Na₂HAsO₄ 8 mM. Changes in metabolites are given as μ moles/hr./vessel.

Substrates	Inhibitor added	(a) Uptake of O ₂	(b) Lactate formed			(c) Lactate formed (corrected*)			(d) Change in 3-phosphoglycerate			Inhibition (%)	(e) Total phosphoglycerate formed (corrected†)			(e-c) (Air - N ₂)			(f) L-Glycerol 1-phosphate formed			(g) Pyruvate uptake		
			Air	N ₂	N ₂	Air	N ₂	N ₂	Air	N ₂	N ₂		Air	N ₂	N ₂	Air	N ₂	N ₂	Air	N ₂	N ₂	Air	N ₂	N ₂
Fructose diphosphate + pyruvate	None	3.08	5.94	8.67	5.91	8.50	30	-0.13	+1.09															
	NaF	2.84	5.50	7.85	5.42	7.72	30	+5.28	+7.59															
	Na ₂ HAsO ₄	2.95	4.85	7.33	4.69	7.16	34	-0.18	+1.10															
	Na ₂ HAsO ₄ + NaF	2.77	4.31	7.32	4.24	7.07	40	+4.22	+6.89															
	None	3.26	0.03	0.17																				
Pyruvate	NaF	2.56	0.08	0.13																				
	Na ₂ HAsO ₄	2.69	0.16	0.17																				
	Na ₂ HAsO ₄ + NaF	2.24	0.07	0.25																				
	None	1.86	4.37	7.94				-0.84	+0.37															

* Correction made for lactate formation in the absence of fructose diphosphate.

† The correction factor (1.1) was based on evidence obtained in experiments in which 2- and 3-phosphoglycerate were measured separately by the chromatographic method of Cowgill (1955) as described by Balázs (1959); the concentration of 2-phosphoglycerate under similar experimental conditions in N₂, as specified in the Table, was found to be 10.1 ± 1.15 (3) % of that of 3-phosphoglycerate.

Table 8. *Effect of crystalline glyceraldehyde phosphate dehydrogenase on the aerobic inhibition of glycolysis with fructose diphosphate as substrate*

Nature and concn. of substrates added		Cofactors (ADP, NAD ⁺) added	Glycer- aldehyde phosphate dehydro- genase (mg. of protein)	Uptake of O ₂	Lactate formed		Lactate formed (corrected*)		Aerobic inhibition of lactate formation (%)	Change in 3-phospho- glycerate		L-Glycerol 1-phosphate formed		Pyruvate uptake		Glucose uptake	
					Air	N ₂	Air	N ₂		Air	N ₂	Air	N ₂	Air	N ₂	Air	N ₂
Fructose diphosphate (10 mm) + Pyruvate (4.5 mm)		++	—	4.8	7.60	12.60	7.51	12.43	40	-0.37	+0.55	0	0.40	0.50	0.14	—	—
		++	0.8	5.4	16.78	18.94	16.69	17.87	7	+2.75	+2.45	0.10	0.42	2.57	2.90	—	—
		++	1.6	5.31	16.87	18.17	16.78	17.99	7	+3.02	+2.82	—	—	2.72	2.46	—	—
Pyruvate (4.5 mm)		++	—	4.95	0.09	0.17	—	—	—	—	—	—	—	3.71	0.50	—	—
		++	0.8	4.95	0.09	0.17	—	—	—	—	—	—	—	3.60	0.38	—	—
		++	1.6	4.99	0.09	0.18	—	—	—	—	—	—	—	3.78	0.39	—	—
Glucose (4.3 mm) + pyruvate (4.5 mm)		—	—	2.71	0.41	1.90	0.41	1.90	78	—	—	0	0.01	2.20	0.64	0.48	1.32
		—	0.8	2.65	0.44	1.90	0.44	1.76	75	—	—	—	—	2.42	0.55	0.44	1.33
		—	1.6	2.63	0.40	1.98	0.40	1.92	79	—	—	—	—	2.35	0.50	0.42	1.30
Pyruvate (4.5 mm)		—	—	2.54	0	0	—	—	—	—	—	—	—	2.04	0.30	—	—
		—	0.8	2.52	0	0.14	—	—	—	—	—	—	—	2.07	0.27	—	—
		—	1.6	2.37	0	0.06	—	—	—	—	—	—	—	1.93	0.29	—	—

* Correction was made for lactate formation in the absence of fructose diphosphate or glucose.

Density-gradient centrifuging, which can to some extent separate the different particles present in the mitochondrial fraction, could not be used in the present work since some of the properties of brain mitochondria that are apparently concerned in the interaction of respiration and glycolysis (e.g. oxidative phosphorylation and respiratory control) are affected by the high concentrations of solutes required for this procedure. It should be emphasized therefore that the brain-mitochondrial system can serve only as a basis of approach to whole-cell studies.

The formation of lactate in the brain-mitochondrial system under the conditions of the present experiments was shown to be mainly due to glycolysis. The amount of lactate formed through the reduction of pyruvate by NADH generated by processes other than the oxidation of glyceraldehyde phosphate was relatively small; the anaerobic dismutation of pyruvate could account for only 2-7 % of the total lactate formed, depending on the experimental conditions (Table 1). In the dismutation reaction the NADH produced by the oxidation of 1 molecule of pyruvate is utilized by lactate dehydrogenase to form lactate from another molecule of pyruvate. The rates of dehydrogenation and reduction of pyruvate in the brain-mitochondrial preparations were much greater than the rate of anaerobic dismutation of pyruvate. Succinate is known to increase the amount of mitochondrial NADH (Chance & Hollunger, 1957; Birt & Bartley, 1960) and in the presence of succinate and Amytal

Fig. 3. Demonstration of the reversibility of the aerobic control of glycolysis. Basic medium; 10 mm-fructose diphosphate, 6.6 mm- α -oxoglutarate, 15 mm-pyruvate, 0.9 mm-ATP, 1.3 mm-AMP, 0.33 mm-NAD⁺, 3 mm-inorganic phosphate, 8 mm-Na₂HAsO₄, 10 mm-NaF; mitochondrial protein 3.1 mg./vessel. Oxygen uptake 4.3 μ -moles/40 min. Gas phase: $\circ$ — $\circ$, air; $\bullet$ — $\bullet$, N₂. Change of gas phase was made at arrow.

Table 9. *Effect of respiratory inhibitors and dinitrophenol on the aerobic inhibition of lactate formation with fructose diphosphate as substrate in the presence of cofactors*

Basic medium; 2 mM-ADP, 0.33 mM-NAD⁺; substrates were as indicated. Potassium cyanide and NaN₃ were injected in suitable concentrations into the vessels through a rubber tube after 6.5 min. of gassing with N₂. In the aerobic vessels with these inhibitors no alkali was added in the centre well and KCN and NaN₃ were present from the beginning of the incubation. In the presence of KCN and NaN₃ the changes in the concentration of the metabolites were determined from the difference in the concentrations reached between 7 and 47 min. of incubation. Oxygen uptakes and substrates changes are given as μ moles/hr./100 mg. of protein.

Expt. no.	Nature and concn. of substrates added	Nature and concn. of inhibitor added	Uptake of O ₂	Lactate formation		Change in 3-phosphoglycerate		Change in L-glycerol 1-phosphate	
				Air	N ₂	Air	N ₂	Air	N ₂
1	Pyruvate (9.1 mM) + fructose diphosphate (12.5 mM)	None	104.8	202.0	294.9	-4.4	+36.9	+5.0	+17.3
		Amytal (1.8 mM)	41.2	221.2	311.6	-4.35	+42.5	+3.2	+15.9
		Dinitrophenol (0.1 mM)	141.5	179.9	273.1	-16.5	+44.6	+2.1	+17.1
2	Pyruvate (9.1 mM)	None	110.9	1.0	5.8	.	.	.	.
		Amytal (1.8 mM)	23.2	3.6	4.8	.	.	.	.
		Dinitrophenol (0.1 mM)	140.0	1.6	3.0	.	.	.	.
3	Pyruvate (6.7 mM) + fructose diphosphate (10 mM) + α -oxoglutarate (8.5 mM)	None	448.0	142.1	226.8	.	.	.	.
		Antimycin A (0.0012 mM)	59.5	142.1	216.7	.	.	.	.
		None	418.6	93.7	213.8	.	.	.	.
4	α -Oxoglutarate (8.5 mM) + fructose diphosphate (10.0 mM)	KCN (1.0 mM)	.	90.7	213.8	.	.	.	.
		KCN (2.0 mM)	.	96.2	201.3	.	.	.	.
		KCN (5.0 mM)	.	87.7	175.4	.	.	.	.
4	α -Oxoglutarate (8.5 mM) + fructose diphosphate (10.0 mM)	None	304.2	110.5	238.0	.	.	.	.
		NaN ₃ (1.0 mM)	.	87.2	203.1	.	.	.	.
		NaN ₃ (2.0 mM)	.	85.1	205.6	.	.	.	.
		NaN ₃ (5.0 mM)	.	87.2	176.5	.	.	.	.
		NaN ₃ (10.0 mM)	.	58.1	150.0	.	.	.	.

Table 10. *Effect of monothiools and BAL on the aerobic inhibition of glycolysis*

Basic medium; substrates were as indicated; 2 mM-ADP, 0.33 mM-NAD⁺ when added; glutathione and cysteine were freshly dissolved and pH was adjusted to 7.2; BAL was dissolved in dimethylformamide and the concentration of the solvent was 0.4% (v/v) in both the experimental and control flasks. Oxygen uptakes and changes in metabolites are given as μ moles/hr./100 mg. of protein.

Expt. no.	Nature and concn. of substrates added	Cofactors (ADP, NAD ⁺) added	Nature and concn. of 'SH' compound added	Uptake of O ₂	Lactate formed		Glucose uptake		Pyruvate uptake	
					Air	N ₂	Air	N ₂	Air	N ₂
1	Fructose diphosphate (10 mM) + pyruvate (4.2 mM) Pyruvate (4.2 mM)	+	None	80.6	136.2	274.6	.	.	.	.
			BAL (2 mM)	119.8	323.7	310.0	.	.	.	.
			None	87.7	1.4	4.1	.	.	.	.
			BAL (2 mM)	120.0	2.8	4.5	.	.	.	.
2	Fructose diphosphate (10 mM) + α -oxoglutarate (10 mM) + pyruvate (6.7 mM)	+	None	387.8	149.3	253.0	.	.	.	.
			Glutathione (1 mM)	393.3	160.2	263.4	.	.	.	.
			Glutathione (5 mM)	386.4	169.5	248.9	.	.	.	.
			Glutathione (10 mM)	314.5	128.6	224.0	.	.	.	.
			Cysteine (1 mM)	279.3	152.9	245.7	.	.	.	.
			Cysteine (5 mM)	252.3	152.9	226.0	.	.	.	.
3	Glucose (2.9 mM) + pyruvate (5.3 mM) Pyruvate (5.3 mM)	-	None	47.4	10.3	37.1	5.0	20.4	35.3	4.9
			BAL (2 mM)	62.0	11.0	34.1	5.0	19.4	44.9	5.9
			None	50.9	0.5	3.0	.	.	31.6	11.4
			BAL (2 mM)	68.6	0.7	1.9	.	.	50.2	14.6

the non-glycolytic reduction of pyruvate was increased (Table 4). Thus these observations indicate the presence of a rate-limiting reaction in the formation of NADH from pyruvate in nitrogen, e.g. a restriction in the removal of acetyl-CoA (suggested by Professor Sir Hans Krebs).

The NADH produced at the oxidoreductive step of glycolysis was oxidized mainly by pyruvate and lactate dehydrogenase in the brain-mitochondrial system. This could be shown when mitochondrial preparations fortified with ADP and NAD⁺ were incubated with fructose diphosphate, pyruvate and fluoride in the presence or absence of arsenate (Table 7). Under these conditions the main reactions are as shown in (1) and (2).

Under these conditions the amount of NADH produced by the oxidation of glyceraldehyde phosphate was given by the amount of phosphoglycerate formed. The amount of NADH oxidized by pyruvate could be determined by the lactate formed (after correcting for the dismutation of pyruvate). The difference between the amount of phosphoglycerate and lactate formed (Table 7, *e-c*) indicated the amount of NADH oxidized by reactions other than (2); this was found to be about 7% of the total NADH produced by reaction (1). Most of the NADH not oxidized by pyruvate was apparently oxidized by dihydroxyacetone phosphate: this was indicated by the amount of L-glycerol 1-phosphate formed in nitrogen (reaction 3).

The concentration of L-glycerol 1-phosphate found after incubation in air was less than in nitrogen, but this may be because this substance is oxidized by brain-mitochondrial preparations. The transfer of reducing equivalents from NADH produced by reaction (1) to the oxidation sites of the electron-transport chain was apparently restricted in the brain-mitochondrial system, since the amount of NADH that did not react by reaction (2) (Table 7, *e-c*) was relatively small and could be mainly accounted for by the L-glycerol 1-phosphate formed. These properties of the brain-mitochondrial fraction make it a convenient system for studying the mechanism of the aerobic control of glycolysis.

The present work shows that, contrary to the previous assumption (Balázs, 1959), the reversible aerobic inhibition of glycolysis associated with brain-mitochondrial preparations depends on more than one mechanism; two separate mechanisms may be demonstrated by selecting suitable experimental conditions. One mechanism was clearly

Table 11. *Effect of dinitrophenol and glyceraldehyde phosphate dehydrogenase on the aerobic inhibition of glycolysis in the presence of cofactors, with fructose diphosphate and glucose as glycolytic substrates*

Basic medium; 10 mM-fructose diphosphate, 7.5 mM-pyruvate, 3 mM-L-malate, 2 mM-ADP, 0.33 mM-NAD⁺. Additions: 0.1 mM-dinitrophenol; glyceraldehyde phosphate dehydrogenase freshly dialysed from stock enzyme suspension; 0.8–1.6 mg. of protein was used/vessel. Values for lactate represent the difference between the amount of lactate formed in the presence and absence of fructose diphosphate and glucose. Values for uptake of O₂ and lactate formation are the mean of three experiments and are given as μ moles/hr./100 mg. of protein.

Additions	Uptake of O ₂	Lactate formed		Aerobic inhibition of lactate formation (%)
		Air	N ₂	
None	359.3	138.8	271.4	49
Dinitrophenol	380.2	213.9	271.8	21
Glyceraldehyde phosphate dehydrogenase	365.5	322.1	418.4	23
Dinitrophenol + glyceraldehyde phosphate dehydrogenase	373.4	389.9	416.5	6

Table 12. *Summary of the effect of different experimental conditions on the aerobic inhibition of glycolysis*

'Glucose-unfortified' system: the mitochondrial preparation was incubated in the basic medium, which contained glucose and pyruvate. 'Fructose diphosphate-fortified' system: the mitochondrial suspension was incubated in the basic medium in the presence of fructose diphosphate, pyruvate, ADP and NAD⁺ (for details cf. Table 1). +, Aerobic inhibition of glycolysis substantially reduced; -, no effect.

Experimental conditions	'Glucose-unfortified' system	'Fructose diphosphate-fortified' system
Presence of respiratory inhibitors	+	-
Presence of uncouplers of oxidative phosphorylation	+	-
Presence of glyceraldehyde phosphate dehydrogenase	-	+
Presence of BAL	-	+
Gas phase changed	+	+

shown by arranging the experimental conditions in such a way that the cofactors (ATP, ADP, NAD⁺) became the pacemakers of glycolysis ('glucose-unfortified system'). Another mechanism was best demonstrated when an enzyme, glyceraldehyde phosphate dehydrogenase, was the rate-limiting factor in glycolysis ('fructose diphosphate-fortified system'). By choosing suitable experimental conditions the two different inhibitory mechanisms can be made to operate together in the same system.

The main differences between the two inhibitory mechanisms are summarized in Table 12. The aerobic inhibition of glycolysis in the unfortified system (about 70 % inhibition both of glucose consumption and lactate formation) is removed by respiratory inhibitors (cyanide, Amytal, antimycin A and 2-n-heptyl-4-hydroxyquinoline N-oxide).

Electron transport does not appear to be the major factor in the control of glycolysis by this mechanism since (1) the residual respiration, which is insensitive to antimycin A, hydroxyquinoline N-oxide and Amytal, cannot maintain the aerobic inhibition and (2) uncouplers of oxidative phosphorylation (dinitrophenol and dicoumarol) remove the aerobic inhibition of glycolysis although they increase the rate of electron transport approximately threefold. The effect of respiratory inhibitors and uncouplers of phosphorylation on the aerobic control of glycolysis was confirmed by Rendina & Hellerman (1961) in the brain-mitochondrial system. Similar results obtained with cell-free preparations of Ehrlich ascites-tumour cells were reported by Tiedemann & Born (1960) while these experiments were in progress.

In the 'fructose diphosphate-fortified system' neither respiratory inhibitors nor the uncouplers of phosphorylation decreased the aerobic inhibition of lactate formation.

The difference in the mechanism of the two inhibitions was clearly shown by the difference in their response to added crystalline glyceraldehyde phosphate dehydrogenase. The addition of the enzyme had no effect on the aerobic inhibition of lactate formation in the 'glucose-unfortified system', but was able to reduce it substantially in the 'fructose diphosphate-fortified system'. Both inhibitions are completely reversible, as is shown by changing the gas phase from air to nitrogen.

The inhibition of glycolysis in the 'fructose diphosphate-fortified system' appears to be a direct, specific inhibition of glyceraldehyde phosphate dehydrogenase in the presence of oxygen (Balázs, 1959). Since glyceraldehyde phosphate dehydrogenase is apparently a thiol enzyme, its activity might be decreased by the oxidation of the essential thiol groups of the protein under aerobic

conditions. Hatch & Turner (1959) found that oxidation of the essential thiol groups of this enzyme was responsible for the aerobic inhibition of glycolysis in a pea-seed extract. These authors have put forward a probable mechanism for the reversible aerobic inhibition of glycolysis in this system: they conclude that there is a non-enzymic metal-catalysed oxidation of the thiol groups of the enzyme and an enzymic reduction of the protein disulphide groups (Hatch & Turner, 1960). It is feasible that a similar mechanism is involved in brain-mitochondrial preparations, but certain differences must be considered. It is unlikely that the oxidation of the protein thiol groups is brought about by metal catalysis in brain-mitochondrial preparations since the chelating agent EDTA (2 mM), which protects the pea-seed enzyme, was regularly included in the reaction medium in the present experiments, but did not prevent the inhibition of glyceraldehyde phosphate dehydrogenase.

The pea-seed glyceraldehyde phosphate dehydrogenase can be protected from oxidation by anaerobic conditions, or by monothiols, such as glutathione; in brain-mitochondrial preparations the enzyme could be reactivated by anaerobic conditions, but glutathione or cysteine was almost ineffective. BAL, on the other hand, removed the inhibition of the brain glyceraldehyde phosphate dehydrogenase completely; a 50% protective effect was observed with about 0.3 mM-BAL.

The protective action of BAL suggests that in the functional part of the brain glyceraldehyde phosphate dehydrogenase vicinal dithiol groups may be involved. The results of Peters, Sinclair & Thompson (1946) give some support to this possibility. They found that the sensitivity of brain glyceraldehyde phosphate dehydrogenase to trivalent arsenicals, which are considered to be fairly specific inhibitors of dithiol groups, was much higher than that of the muscle enzyme. 6 μ M-Lewisite inhibited the brain glyceraldehyde phosphate dehydrogenase by 58%, whereas 300 μ M-lewisite had no effect on the muscle enzyme. Another possible explanation for the protective effect of BAL might be that it inhibits an enzyme that oxidizes the thiol groups of glyceraldehyde phosphate dehydrogenase. Such an effect of BAL is known to occur with enzymes containing heavy-metal prosthetic groups (polyphenol oxidase, tyrosinase, catalase, peroxidase). However, these enzymes are usually inhibited also by cyanide (Hurwitz, 1955; Lerner, 1953), but cyanide was ineffective in the brain-mitochondrial system.

The other controlling mechanism of glycolysis in which phosphorylation is concerned and which is shown most clearly in the 'glucose-unfortified system' behaves very much like the Pasteur re-

action in the whole cell. The main similarities are: (1) both glucose consumption and lactate formation are inhibited under aerobic conditions; (2) the aerobic inhibition of glycolysis is removed by respiratory inhibitors and by uncouplers of oxidative phosphorylation; (3) the inhibition is completely reversible. It was suggested by Aisenberg & Potter (1957) that the Pasteur reaction in mitochondrial systems might operate through the formation of some hypothetical high-energy intermediate of oxidative phosphorylation, which is able to inhibit the phosphofructokinase reaction and perhaps also the hexokinase reaction. Passonneau & Lowry (1962), studying the kinetic properties of purified phosphofructokinase, put forward the idea that this enzyme may be primarily responsible for the Pasteur reaction in the intact cell. They found that at low substrate concentrations comparable with those in the cell, the activity of this enzyme is increased by low concentrations of ATP or high concentrations of AMP, ADP and inorganic phosphate; such conditions occur when oxidative phosphorylation is inhibited in the whole cell. Our results have also given evidence that the aerobic control of glycolysis in the 'glucose-unfortified system' is connected with oxidative phosphorylation. However, evidence was previously given (Balázs & Richter, 1960b) that the aerobic inhibition of glucose uptake under these conditions is due to a restricted availability of mitochondrial ATP for the phosphorylation of glucose. There was no indication of an inhibition of hexokinase by glucose 6-phosphate, since the phosphorylation rate was not stimulated when yeast hexokinase was added, or when glucose was replaced by 2-deoxyglucose (Balázs, Magyar & Richter, 1962). Although the inhibition of phosphofructokinase was not excluded as a controlling mechanism in the present system, the low rate of the hexokinase reaction was apparently sufficient to account for the aerobic inhibition of glucose uptake. When adenine nucleotides (AMP, ADP or ATP) were added to the system an accumulation of fructose diphosphate was observed (Balázs, 1959), but a dinitrophenol-sensitive aerobic inhibition of lactate formation still remained (Table 5). Thus the present results, if they can be applied to the whole cell, indicate that a compartmentation of ATP in the cell may play a major role in the Pasteur reaction; this was previously suggested by Lynen & Koenigsberger (1951), Racker (1956) and Chance & Hess (1959). However, it is probable that also in the intact cell the Pasteur reaction is the result of the operation of more than one mechanism.

Since the observations in the 'glucose-unfortified system' indicated that oxidative phosphorylation is involved in the aerobic control of glycolysis, an attempt was made to determine whether the three

possible sites of oxidative phosphorylation between NADH and oxygen are of equal importance in the interaction of respiration and glycolysis. After excluding the NADH-flavoprotein site of oxidative phosphorylation by using succinate as substrate in the presence of Amytal, the aerobic control of glycolysis was no longer maintained, although oxidative phosphorylation at the two other sites was hardly affected. Similar results were previously obtained with preparations from brain containing whole cells (Webb & Elliott, 1951). This shows that the aerobic control of glycolysis is not dependent on oxidative phosphorylation in general. The observation that the aerobic control of glycolysis was prevented by blocking the NADH-flavoprotein site of oxidative phosphorylation suggests that this is the main site for the interaction with glycolysis. The special significance of this site in relation to two possible intermediate reactions of oxidative phosphorylation (adenosine triphosphatase and ATP- ^{32}P exchange reactions) is emphasized by the work of Löw, Siekevitz, Ernster & Lindberg (1958), Siekevitz, Löw, Ernster & Lindberg (1958) and Löw (1959). However, until more direct experimental evidence is available this finding is also open to other explanations.

SUMMARY

1. Two different mechanisms have been shown to operate in the reversible aerobic inhibition of glycolysis associated with mitochondrial preparations of rat brain. They can be made to operate separately or together by the choice of suitable experimental conditions.

2. One inhibitory mechanism operates in the system with glucose as substrate in the absence of added ADP and NAD^+ . This resembles the Pasteur reaction obtained with whole-cell preparations.

3. It is concluded that this form of aerobic inhibition of glycolysis is connected with some process of oxidative phosphorylation. It appears that the NADH-flavoprotein site of oxidative phosphorylation is more important in this connexion than the other two sites.

4. Another inhibitory mechanism of glycolysis is the aerobic inhibition of glyceraldehyde phosphate dehydrogenase. This was demonstrated in a system containing fructose diphosphate as the glycolytic substrate with added cofactors; the site of the inhibition was verified by the addition of crystalline glyceraldehyde phosphate dehydrogenase.

5. This form of inhibition is not removed by respiratory inhibitors or by uncouplers of phosphorylation, but can be completely reversed by changing to anaerobic conditions.

6. Glyceraldehyde phosphate dehydrogenase can be protected from aerobic inhibition by 2,3-

dimercaptopropanol, but monothiols (glutathione and cysteine) are almost ineffective.

I thank Dr D. Richter for his interest and encouragement, Dr D. M. Needham for her great help in revising the manuscript and Professor Sir Hans Krebs, Dr F. A. Holton and Dr J. R. Lagnado for valuable discussions. The technical assistance of Mr R. G. Fish and Miss W. Clee is acknowledged.

REFERENCES

- Aisenberg, A. C. & Potter, V. R. (1957). *J. biol. Chem.* **224**, 1115.
 Aisenberg, A. C., Reinafarje, B. & Potter, V. R. (1957). *J. biol. Chem.* **224**, 1099.
 Aldridge, W. N. & Johnson, M. K. (1959). *Biochem. J.* **73**, 270.
 Balázs, R. (1959). *Biochem. J.* **72**, 561.
 Balázs, R. & Lagnado, J. R. (1959). *J. Neurochem.* **5**, 1.
 Balázs, R., Magyar, K. & Richter, D. (1962). *Proc. 5th int. Symp. Neurochem.*
 Balázs, R. & Richter, D. (1960a). *Biochem. J.* **74**, 30P.
 Balázs, R. & Richter, D. (1960b). *Biochem. J.* **76**, 67P.
 Barker, S. B. & Summerson, W. H. (1941). *J. biol. Chem.* **138**, 535.
 Beaufay, H., Berleux, A. M. & Doyen, A. (1957). *Biochem. J.* **66**, 32P.
 Beisenherz, G., Boltze, H. J., Bücher, T., Czok, R., Garbade, K. H., Meyer-Arendt, E. & Pfeleiderer, G. (1953). *Z. Naturf.* **8**, 555.
 Birt, L. M. & Bartley, W. (1960). *Biochem. J.* **76**, 427.
 Chance, B. & Hess, B. (1959). *Science*, **129**, 700.
 Chance, B. & Hollunger, G. (1957). *Fed. Proc.* **16**, 163.
 Cori, G. T., Slein, M. W. & Cori, C. F. (1948). *J. biol. Chem.* **173**, 605.
 Cornforth, J. W. & James, A. T. (1956). *Biochem. J.* **63**, 124.
 Cowgill, R. W. (1955). *Biochim. biophys. Acta*, **14**, 614.
 Cremer, J. E. (1960). *Biochim. biophys. Acta*, **41**, 155.
 Dodds, E. C. & Greville, G. D. (1934). *Lancet*, **i**, 398.
 Ernster, L., Jalling, O., Löw, H. & Lindberg, O. (1955). *Exp. Cell Res. suppl.* **3**, 124.
 Friedemann, L. E. & Haugen, G. E. (1943). *J. biol. Chem.* **147**, 415.
 Gatt, S. & Racker, E. (1959). *J. biol. Chem.* **234**, 1024.
 Graffi, A. & Schneider, E. J. (1956). *Naturwissenschaften*, **43**, 376.
 Hatch, M. D. & Turner, J. F. (1959). *Biochem. J.* **72**, 524.
 Hatch, M. D. & Turner, J. F. (1960). *Biochem. J.* **75**, 66.
 Hohorst, H. J., Kreutz, F. H. & Bücher, T. (1959). *Biochem. Z.* **332**, 18.
 Hurwitz, J. (1955). *J. biol. Chem.* **212**, 757.
 Johnson, M. K. (1960). *Biochem. J.* **77**, 610.
 Kiessling, W. (1938). *Biochem. Z.* **298**, 421.
 Krebs, H. A. & Johnson, W. A. (1937). *Biochem. J.* **31**, 645.
 Lerner, A. B. (1953). *Advanc. Enzymol.* **14**, 73.
 Lohmann, K. & Jendrassik, L. (1926). *Biochem. Z.* **178**, 419.
 Löw, H. (1959). *Biochim. biophys. Acta*, **32**, 1.
 Löw, H., Siekevitz, P., Ernster, L. & Lindberg, O. (1958). *Biochim. biophys. Acta*, **29**, 392.
 Lowry, O. H., Rosebrough, N. J., Farr, A. L. & Randall, R. J. (1951). *J. biol. Chem.* **193**, 265.

- Lynen, F. & Koenigsberger, R. (1951). *Leibigs Ann.* **573**, 60.
- Nelson, N. (1944). *J. biol. Chem.* **153**, 375.
- Passonneau, J. V. & Lowry, O. H. (1962). *Biochem. biophys. Res. Commun.* **7**, 10.
- Peters, R. A., Sinclair, H. M. & Thompson, R. H. S. (1946). *Biochem. J.* **40**, 516.
- Petruschka, E. & Giuditta, A. (1959). *J. biophys. biochem. Cytol.* **6**, 129.
- Potter, V. R. & Reif, A. E. (1952). *J. biol. Chem.* **194**, 287.
- Price, V. E. & Levintow, L. (1952). *Biochemical Preparations*, vol. 2, p. 22. Ed. by Ball, E. G. London: Chapman and Hall.
- Racker, E. (1956). *Ann. N.Y. Acad. Sci.* **63**, 1017.
- Rendina, G. & Hellerman, L. (1961). *Fed. Proc.* **20**, 83.
- Schwartz, A. & Lee, K. S. (1960). *Biochim. biophys. Acta*, **44**, 590.
- Segal, S., Blair, A. E. & Wyngaarden, J. B. (1956). *J. Lab. clin. Med.* **48**, 137.
- Siekevitz, P., Löw, H., Ernster, L. & Lindberg, O. (1958). *Biochim. biophys. Acta*, **29**, 378.
- Sisakian, N. M. & Pinus, E. A. (1958). *Biokhimiya*, **23**, 904.
- Somogyi, M. (1945). *J. biol. Chem.* **160**, 69.
- Somogyi, M. (1952). *J. biol. Chem.* **195**, 19.
- Terner, C. (1954). *Biochem. J.* **56**, 471.
- Tiedemann, H. & Born, J. (1958). *Biochim. biophys. Acta*, **27**, 656.
- Tiedemann, H. & Born, J. (1960). *Hoppe-Seyl. Z.* **321**, 205.
- Umbreit, W. W., Burris, R. H. & Stauffer, J. F. (1945). (editors). *Manometric Techniques and Related Methods for the Study of Tissue Metabolism*, 3rd ed. Minneapolis: Burgess Publishing Co.
- Von Korff, R. W. (1959). *Biochim. biophys. Acta*, **31**, 467.
- Von Korff, R. W. & Twedt, R. M. (1957). *Biochim. biophys. Acta*, **23**, 143.
- Webb, J. L. & Elliott, K. A. C. (1951). *J. Pharmacol.* **103**, 24.
- Weil-Malherbe (1937). *Biochem. Z.* **31**, 2202.
- Whittaker, V. P. (1959). *Biochem. J.* **72**, 694.

Biochem. J. (1963) **86**, 509

The Intracellular α -Galactosidase of a Rumen Strain of *Streptococcus bovis*

By R. W. BAILEY

Plant Chemistry Division, Department of Scientific and Industrial Research, Palmerston North, New Zealand

(Received 7 August 1962)

The α -galactosidases have not been studied as extensively as β -galactosidases. Thus a review of these enzymes by Wallenfels & Malhotra (1961) devotes 50 pages to an account of β -galactosidases but only 8 pages to the corresponding α -enzymes. In addition almost all of the α -galactosidases that have been examined in any detail have come from yeasts, moulds or higher plants. Bacterial enzymes of this class appear to be rare. Carrere, Lambin & Courtois (1960), for example, investigated the intestinal flora of many animal species, including ruminants, and obtained only one bacterial strain, from guinea-pig intestine, which produced an α -galactosidase.

Many pasture plants contain α -linked galactose present in oligosaccharides such as raffinose and the galactosylgalactosyl glycerides of the lipid fraction (Weenink, 1961). Cattle and sheep feeding on pasture metabolize these compounds in their rumens, the microflora of which should, therefore, contain organisms producing α -galactosidases. A preliminary study (Bailey, 1962) suggested that this was the case and cell extracts from mixed rumen bacteria were found to exhibit α -galactosidase activity. The enzyme was also found in the cells of one rumen bacterial species, *Streptococcus*

bovis. Extracts prepared from the cells of this organism grown in a glucose medium contained scarcely any other simple oligosaccharidases. A detailed study of the *S. bovis* α -galactosidase has now been made and the results obtained form the subject of the present paper.

EXPERIMENTAL

Buffers. The following buffers were used: sodium citrate (0.1M, pH 6.0), sodium acetate (0.1M, pH 6.0), sodium phosphate (0.1M, pH 6.0) and citric acid (0.1M)-disodium hydrogen phosphate (0.02M) (McIlvaine, 1921; pH 6.0 except in studies on the effect of pH on enzyme activity).

Organism. *S. bovis*, strain I (Bailey & Oxford, 1958), was revived from a freeze-dried culture prepared from cells that had been grown in the glucose medium described below. The organism was grown in a medium containing glucose (1%, w/v), sodium acetate (anhydrous) (1.2%, w/v), Bacto-Yeast Extract (0.5%) and Bacto-Tryptose (1.4%, w/v). Cultures were incubated at 37° under aerobic conditions with the atmosphere in the flasks enriched with CO₂ gas.

Cell extracts. Medium was inoculated and incubated for 18–20 hr., by which time good growth had occurred and the cultures contained 2–3 g. wet wt. of cells/l. Cells were harvested by centrifuging at 25 000g for 30 min. and washed by suspending in either acetate or citrate buffer and