of the cyclohexyl esters of the acids produced showed cyclohexyl propionate, dicyclohexyl malonate and dicyclohexyl succinate to be present, the last two in a 5:2:1 molar ratio (calculated from the areas under the peaks on the chromatogram). Six structures are therefore possible for the docosahexaenoic acid: the double bonds could be at positions 4,7,10,13,16,19, or 3,7,10,13,16,19, or 3,6,10,13,16,19, or 3,6,9,13,16,19, or 3,6,9,12,16,19, or 3,6,9,12,15,19. Reduction of the methoxycarbonyl group of the methyl docosahexaenoate (I) to hydroxymethyl with lithium aluminium hydride to give the docosahexaenol (II), conversion of this into the toluene-*p*-sulphonyl ester (III) and further reduction with lithium aluminium hydride gave the docosahexaene (IV). Gas chromatography of the cyclohexyl esters of the oxidation products of this compound showed cyclohexyl propionate, cyclohexyl *n*-butyrate, dicyclohexyl malonate and (in small amounts) dicyclohexyl succinate to be present. The formation of *n*-butyric acid on oxidation of the docosahexaene formed by reduction of the methoxycarbonyl group of the methyl docosahexaenoate to a methyl group is possible only with the first of the six structures given above. The succinic acid present in the oxidation mixture is presumably due to the incomplete reduction of the docosahexaenyl toluene-*p*-sulphonate by lithium aluminium hy-

dride. The double bonds are therefore in the 4,7,10,13,16,19 positions in the docosahexaenoic acid. The presence of a peak at 710 cm.⁻¹ in the infrared-absorption spectrum and the absence of a peak in the 960–970 cm.⁻¹ region indicates *cis*-double bonds.

SUMMARY

1. all-*cis*-Docosa-4,7,10,13,16,19-hexaenoic acid has been isolated as the methyl ester from cattle retinae and its structure determined by degradation.

We thank Professor Sir Hans Krebs, F.R.S., for advice and interest in the work, which was aided by the Rockefeller Foundation and the National Institutes of Health, United States Public Health Service, Grant no. 3369 to the Department of Biochemistry, University of Oxford. A.R.H. thanks the Medical Research Council for financial support.

REFERENCES

- Bartley, W., van Heyningen, R., Notton, B. M. & Renshaw, A. (1962). *Biochem. J.* **85**, 332.
Hörhammer, L., Wagner, H. & Wolff, P. (1960). *Hoppe-Seyl. Z.* **319**, 7.
James, A. T. (1960). *Meth. biochem. Anal.* **8**, 1.
Lemieux, R. U. & Rudloff, E. von (1955). *Canad. J. Chem.* **33**, 1701.
Metcalfe, L. D. & Schmitz, A. A. (1961). *Analyt. Chem.* **33**, 363.
Renshaw, A. & Biran, L. A. (1962). *J. Chromat.* **8**, 343.

Biochem. J. (1963) **87**, 265

Carbohydrates in Protein

5. PROCEDURES FOR THE ISOLATION OF GLYCOPEPTIDES FROM HEN'S-EGG ALBUMIN AND THEIR OXIDATION BY PERIODATE*

By A. P. FLETCHER, G. S. MARKS,† R. D. MARSHALL AND A. NEUBERGER
Department of Chemical Pathology, St Mary's Hospital Medical School, London, W. 2

(Received 2 October 1962)

It has been suggested that the covalent bond linking carbohydrate to protein in hen's-egg albumin is of the *N*-glycosidic type, linking the amide nitrogen of asparagine directly to the reducing carbon of a sugar molecule (Johansen, Marshall & Neuberger, 1961; Marks & Neuberger, 1961; Nuenke & Cunningham, 1961). Yamashina & Makino (1962) suggested that the linkage may be that of an imino ether, the oxygen of the amide group of asparagine being directly linked to the reducing carbon of a sugar moiety.

* Part 4: François, Marshall & Neuberger (1962).

† Present address: Department of Pharmacology, University of Alberta, Edmonton, Alberta, Canada.

To carry out further studies on the carbohydrate-protein linkage, we wished to obtain a glycopeptide containing aspartic acid as the only amino acid constituent, i.e. containing, we believe, 5 atoms of nitrogen/5 molecules of mannose (Johansen, Marshall & Neuberger, 1960, 1961; François, Marshall & Neuberger, 1962), but this has proved difficult. Most proteolytic procedures gave glycopeptides containing other amino acids in addition to asparagine or aspartic acid, and such peptides are insensitive to further attack by enzymic procedures (Jevons, 1958; Johansen, Marshall & Neuberger, 1958, 1961; Nuenke & Cunningham, 1961; Lee & Montgomery, 1962). We have there-

fore compared, with respect to both degree of degradation of the protein and the amino acid composition of the purified glycopeptide, the value of various proteolytic enzymic procedures for the preparation of purified glycopeptides.

We report also the results of the periodate oxidation of a purified glycopeptide. Two groups of workers (Bragg & Hough, 1961; Lee & Montgomery, 1961; Chatterjee, Durant, Lee & Montgomery, 1961) have applied this technique to an investigation of the structure of the carbohydrate by using whole egg albumin. However, since the protein is oxidized by periodate, these studies have been of limited value. We have eliminated such interference by working with a purified glycopeptide rather than with the whole protein. This enabled us to measure the periodate uptake and the formic acid liberation, and also to determine quantitatively which monosaccharide units were attacked by periodate.

Short reports of these findings have been given (Marks, Marshall, Neuberger & Papkoff, 1962; Marks, Marshall & Neuberger, 1962).

EXPERIMENTAL

Analytical methods

Carbohydrate (mannose) analyses were performed by an orcinol-sulphuric acid method (François *et al.* 1962) and the nitrogen was determined by a Kjeldahl procedure (Chibnall, Rees & Williams, 1943).

Amino acid analyses were carried out by Moore & Stein chromatography with the Technicon Autoanalyser (Technicon Instruments Co. Ltd., Hanworth Lane, Chertsey, Surrey), an adaptation of the cyanide-ninhydrin procedure (Yemm & Cocking, 1955; Rosen, 1957) being used for the analyses. Hydrolyses were carried out *in vacuo* in sealed tubes for 18 hr. at 110°.

Ammonia determinations were carried out with phosphotungstic acid as described by Marshall & Neuberger (1960).

Preparation of glycopeptide: method 1

(a) *Enzymic digestion.* This was similar to that described by Johansen *et al.* (1961). An aqueous 5% solution of egg albumin (186 g. of protein, equivalent to 3.71 g. of 'mannose', in 3.6 l.) was denatured by heating in a boiling-water bath for about 45 min. After equilibration of the mixture at 37°, the pH was adjusted to 2.8 with hydrochloric acid. Crystalline pepsin (Armour Laboratories, Hampden Park, Eastbourne) (3 g.), dissolved in 10 ml. of water, was added. Digestion was allowed to proceed, the pH being kept at 2.8 by the addition of 4N-hydrochloric acid. A further 3 g. of pepsin was added after 24 hr. and 1.5 g. after 48 hr. The total digestion period was 265 hr. The pH of the digest was then adjusted to 7.8 and 0.5 g. each of trypsin and chymotrypsin (Armour Laboratories) were added. Digestion was continued for 48 hr., further quantities (0.5 g.) of trypsin and chymotrypsin being added after 24 hr. The extent of hydrolysis was followed by the ninhydrin reaction (Moore & Stein, 1954). At all times the digest was covered with a layer of toluene.

(b) *Purification of the glycopeptide.* The digest was neutralized (to pH 5-6) and concentrated to about 700 ml., and a small amount of precipitate, which was discarded, was removed by centrifuging. To the supernatant was added 3 vol. of ethanol, when almost all the carbohydrate (3.28 g., equivalent to 89%) was precipitated. After being kept in the cold room overnight, the precipitate was separated by centrifuging, redissolved in water and reprecipitated in the presence of 75% (v/v) ethanol. This precipitate contained 136 atoms of nitrogen/5 molecules of mannose compared with 508 in the original protein. It was dissolved in about 450 ml. of water and the pH adjusted to 2 with 3N-hydrochloric acid. Phosphotungstic acid [130 g., purified by the procedure of Van Slyke & Rieben (1944)] was added as an aqueous 25% solution to precipitate basic amino acids. The supernatant, which contained practically all the carbohydrate (3.25 g.), was shaken with 3 × 500 ml. volumes of pentanol-ether (1:1, v/v) to remove the phosphotungstic acid. This supernatant contained 38 atoms of nitrogen/5 molecules of mannose, 93% of the nitrogen of the whole protein having been removed by these procedures. The pH of this solution was brought to 5.5 with lithium hydroxide and the carbohydrate separated by precipitation in the presence of 75% (v/v) ethanol.

Further purification of the glycopeptide was achieved with Sephadex G-25 (lot no. To-4925, Pharmacia, Uppsala, Sweden). A column (34.5 cm. × 4 cm.) was prepared with 100 g. of Sephadex. An aqueous solution (20 ml.) of the above carbohydrate-containing material (2.7 g. of carbohydrate) was placed on the column and eluted with 15 mM-lithium bromide at a rate of 70 ml./hr. The eluate was collected in 10 ml. fractions. Carbohydrate and nitrogen analyses were carried out (Fig. 2a). Fraction 33 (166 mg. of mannose; 8.65 atoms of nitrogen/5 molecules of mannose) was used in the periodate-oxidation studies. Fractions 28-32 inclusive (1.11 g. of mannose) were combined, concentrated and rechromatographed on Sephadex G-25. Glycopeptide was recovered (0.53 g. of mannose; 7.23 atoms of nitrogen/5 molecules of mannose). This is termed 'glycopeptide fraction A'. The nitrogen content of this fraction was unchanged after electrophoresis at pH 2.3 (in 2N-acetic acid) on a Porath (1956) column as described by Johansen *et al.* (1961).

Reaction of the glycopeptide with benzyloxycarbonyl chloride. This was carried out by the modified Schotten-Baumann procedure of Greenstein & Winitz (1961). Typically, an amount of the glycopeptide equivalent to 140 mg. of mannose (156 μ moles of glycopeptide) was dissolved in 1.5 ml. of water containing a suspension of 300 mg. of sodium hydrogen carbonate. Benzyloxycarbonyl chloride (624 μ moles as two 0.33 ml. portions with 1 hr. between the additions), prepared by the procedure of Farthing (1950), was allowed to react with the glycopeptide solution by shaking it at 37° for about 24 hr. The excess of sodium hydrogen carbonate was removed by filtration and the residue washed once with water. The solution and washings were diluted to 50 ml. with water and the excess of benzyloxycarbonyl chloride was extracted with ether. The benzyloxycarbonyl derivative of the glycopeptide was treated with carboxypeptidase as follows.

Digestion with carboxypeptidase. The solution was reduced to a volume of 5 ml. and adjusted to pH 7.5. Carboxypeptidase (Worthington Biochemical Corp., Freehold, N.J., U.S.A.) (15 mg.) in 5 ml. of solution at pH 7.5 was added.

After digestion for 24 hr. at 37°, the carbohydrate-containing material was purified on a Porath (1956) column (250 g. of Munktell cellulose; 110 cm. $\times$ 3.4 cm.) with 2*N*-acetic acid as eluent as described by Johansen *et al.* (1961).

The benzyloxycarbonyl groups were removed from the resulting material by hydrogenation in the presence of palladium black. Typically, an amount equivalent to 5.68 mg. of mannose in 20 ml. of 2*N*-acetic acid was shaken for 2 hr. in the presence of 15 mg. of palladium black in an atmosphere of hydrogen. After filtration the solution was evaporated to dryness under reduced pressure, and the residue was stored *in vacuo* overnight over potassium hydroxide and P₂O₅, dissolved in water and precipitated by ethanol twice to give 'glycopeptide fraction B'.

Preparation of glycopeptide: method 2

A 5 g. sample of egg albumin and 0.1 g. of Pronase-P [Kingsley & Keith (Chemicals) Ltd., Rex House, 38 King William Street, London, E.C. 4] were dissolved in 100 ml. of 15 mM-calcium chloride solution; the pH was adjusted to 8.5 and the solution was incubated at 37°. The pH was adjusted to 8.5 when necessary. Proteolysis was followed by means of the ninhydrin reagent of Moore & Stein (1948); reaction was complete after 24 hr. The solution was concentrated *in vacuo* to 30 ml., and 120 ml. of ethanol was added. After 48 hr. at 4°, the precipitate that had formed was dissolved in 14 ml. of 0.1*N*-acetic acid and applied to a column of Sephadex G-25 (131 cm. $\times$ 2 cm.) previously equilibrated with 0.1*N*-acetic acid. The column was developed with 0.1*N*-acetic acid and 10 ml. fractions collected. Repeated evaporation of the solution containing the mannose *in vacuo* or drying from the frozen state yielded 'glycopeptide fraction C'. Substitution of the free amino group with the benzyloxycarbonyl radical, treatment with carboxypeptidase and purification on Sephadex G-25 followed by removal of the benzyloxycarbonyl group gave 'glycopeptide fraction D'.

Preparation of glycopeptide: method 3

A solution of egg albumin (5 g.) in water (200 ml.) was brought to pH 7.5 with dilute sodium hydroxide, denatured by heating at 80–82° for 15 min., and then cooled to room temperature. To the solution 2-mercaptoethanol, EDTA and phosphate buffer, pH 7.5, were added to give final concentrations of 10 mM, 10 mM and 50 mM respectively, and the pH was adjusted to 7.5. The solution of denatured egg albumin was digested for 6 hr. with 15 mg. of crystalline papain at 60°, a further 10 mg. being added after 3 hr. (cf. Lee & Montgomery, 1962). Proteolysis was followed by using the ninhydrin reagent of Moore & Stein (1948).

To denature the papain the solution was adjusted to pH 3.5 and heated at 75° for 30 min. (Lineweaver & Schwimmer, 1941). It was then cooled and adjusted to pH 8.5 with dilute sodium hydroxide, and insoluble material was removed by centrifuging. To the supernatant was added calcium chloride to give a final concentration of 15 mM, after correction for the binding of Ca²⁺ ions by the EDTA. Pronase (100 mg.) was added and the solution incubated at 37°; the pH was adjusted to 8.5 when necessary. After 24 hr. a further 100 mg. of Pronase was added and incubation was continued for a further 24 hr. at 37°. After 96 hr. at room temperature 'glycopeptide fraction E' was isolated by precipitation with ethanol and chromatographed on Sephadex G-25 as described under 'method 2' above.

Periodate oxidation

(a) *Measurement of periodate uptake.* The glycopeptide preparation (containing 13.1 mg. of mannose) was dissolved in water (17 ml.), and 10 mM-sodium metaperiodate solution (17 ml.) was added. A blank was prepared by omission of the glycopeptide. At intervals, three 2 ml. samples of the solution were withdrawn and added to Erlenmeyer flasks which contained 96.9 mM-sodium arsenite solution (0.5 ml.) and 0.9% (w/v) sodium hydrogen carbonate solution (0.13 ml.). The walls of the flask were washed down with 1 ml. of water, and 20% (w/v) potassium iodide solution (0.1 ml.) was immediately added. The mixture was left at room temperature for 15 min. and then the excess of arsenite not oxidized by the periodate was titrated with 10 mM-iodine solution (Jackson, 1944; Aminoff & Morgan, 1951).

(b) *Measurement of formic acid liberation.* The glycopeptide preparation (containing 9.9 mg. of mannose) was dissolved in 10 mM-sodium metaperiodate solution (16 ml.) and kept at 5–6°. Formic acid was determined by an iodometric method (Kolthoff & Sandell, 1952) after the removal of periodate with ethylene glycol. To 2 ml. of the solution ethylene glycol (0.25 ml.) was added, followed after 20 min. by 20% (w/v) potassium iodide solution (0.1 ml.). An excess of 10 mM-sodium thiosulphate solution (0.5 ml.) was then added and the excess back-titrated with 10 mM-iodine solution. An appropriate blank solution was prepared by adding ethylene glycol (0.25 ml.) to 20 mM-sodium metaperiodate solution (1 ml.) and after 20 min. adding the corresponding amount of glycopeptide in water (1 ml.). The blank solution was titrated and its formic acid content in terms of 10 mM-sodium thiosulphate subtracted from that shown by the reaction mixture.

(c) *Measurement of the mannose lost on periodate oxidation.* Generally labelled [¹⁴C]mannose (The Radiochemical Centre, Amersham, Bucks.) (0.01 mc) and 0.3 g. of mannose, once-recrystallized, were combined and recrystallized by the procedure of Isbell, Frush & Schaffer (1955).

The glycopeptide (containing 27.75 mg. of mannose) was dissolved in 80 ml. of 5 mM-sodium metaperiodate and after 6 hr. was passed through a column (7 cm. $\times$ 1 cm.) of Dowex 1 (200–400 mesh; acetate form) to remove IO₃⁻ and IO₄⁻ ions (Smith & Willeford, 1954). The column was washed with water (200 ml.) and the combined eluate evaporated to dryness in a rotatory evaporator at less than 40°. Radioactive mannose (35 mg.) was added to the residue and after the addition of 2*N*-hydrochloric acid the mixture was heated at 100° for 3 hr. After dilution of the hydrolysate with water to 100 ml. it was neutralized with Amberlite IR-4B. The resin was washed with water to remove any adhering glycopeptide and the aqueous solution reduced in volume in a rotatory evaporator at 40° to about 5–10 ml. Freshly distilled phenylhydrazine (0.3 ml.) was added followed by acetic acid to give a pH of 5.5. After being kept at room temperature for 2 hr. and at 4° overnight, the precipitate was filtered and washed successively with 50% (v/v) acetic acid, water, ethanol and ether. It was recrystallized by dissolving in hot pyridine (1.5 ml.), filtering at 50° and adding 5 vol. of water at the same temperature. The precipitate was filtered, washed as before and the crystallization repeated, yielding the phenylhydrazone, m.p. 187–188.5° (phenylhydrazone of mannose, m.p. 187–188°). After being dried, the radioactive sample was counted by mounting the dry powder in 0.3 cm.² polythene planchets and by using at least 10 mg. for counting at

infinite thickness in a Geiger-Müller end-window counter. The mannose phenylhydrazone had an activity of 10 163 counts/3000 sec. (s.d., ± 2.0 counts/min.). The mannose phenylhydrazone prepared from the radioactive mannose used in this experiment had an activity of 13 970 counts/3000 sec. (s.d., ± 2.4 counts/min.), and from these values it was calculated that 13.04 mg. (47%) of the original 27.75 mg. of mannose in the glycopeptide had survived periodate oxidation.

(d) *Measurement of the glucosamine lost on periodate oxidation.* The glycopeptide (containing 5.55 mg. of mannose) was dissolved in 5 mm-sodium metaperiodate solution (15.9 ml.) and, after 2 hr., 6 hr., and 24 hr. periods, 5.3 ml. samples were withdrawn and the IO_3^- and IO_4^- ions removed by passing these samples through columns of Dowex 1 (acetate form). The columns were washed with water and the eluates made up to 25 ml. A blank was prepared by dissolving the glycopeptide (containing 1.85 mg. of mannose) in water (25 ml.). Samples (2 ml.) of these solutions were made 4N with respect to hydrochloric acid in stoppered tubes and heated on a boiling-water bath for 4 hr. After being cooled, the contents of the tubes were evaporated to dryness, the residues were dissolved in water (1 ml.) and glucosamine was determined by the method of Rondle & Morgan (1955).

Action of almond emulsin on the glycopeptide preparation. Almond emulsin (100 mg.) was dissolved in water (5 ml.) and dialysed overnight against water (3 l.). A precipitate settled in the dialysis bag. The suspension (2 ml.) was added to 0.15 ml. of a glycopeptide solution (containing 2.8 mg. of mannose), the pH was adjusted to 5 and the mixture was incubated at 35° for 20 hr. An appropriate blank solution was prepared by omitting the glycopeptide. After dilution of the mixture with water (1 ml.), 20% (w/v) trichloroacetic acid solution (0.5 ml.) was added, the precipitate removed by centrifuging and the supernatant extracted several times with ether. The supernatant was chromatographed on Whatman no. 4 paper with a solvent prepared by adding ethyl acetate to the monophasic system, ethyl acetate-water-pyridine-acetone (10:10:5:2, by vol.), until two phases separated; the upper phase was used (Malpress & Hytten, 1958). Mannose was detected by spraying the paper with aniline hydrogen phthalate (Partridge, 1949).

RESULTS AND DISCUSSION

The isolation of glycopeptides from enzymic digests of glycoproteins involves the separation of the carbohydrate-containing component from amino acids and peptides. The procedure may be simplified if all the non-carbohydrate-containing material is as dissimilar as possible in some property to the glycopeptide. The property that we hoped to utilize was that of size. Our previous studies had indicated that a glycopeptide could be obtained whose molecular weight was about 1600–1700; if the other peptide material could be obtained with molecular weights of 500 or less, then the use of gel-filtration would seem possible for the separation. Consequently several enzymic procedures were investigated.

Enzymic digestion procedures. We have compared the rate and extent of proteolysis brought about by: (a) pepsin followed by trypsin and chymotrypsin together, as described by Johansen *et al.* (1961) except that the pepsin digestion was allowed to continue until hydrolysis was complete; (b) Pronase-P, a proteolytic enzyme from *Streptomyces griseus* (Nomoto & Narahashi, 1959a–d; Nomoto, Narahashi & Murakami, 1960a–c); and (c) papain, by a similar procedure to that described by Lee & Montgomery (1962), followed by Pronase-P. The extent of hydrolysis after various times is shown in Fig. 1, where values have been calculated by assuming that each peptide gives the same molar colour yield as leucine in the ninhydrin assay.

Although pepsin can eventually cause a greater degree of hydrolysis of egg albumin (cf. Calvery, 1933) than Pronase-P, this is achieved only after a very prolonged hydrolysis. Indeed the initial rapid reaction with pepsin occurs in about the first 24 hr., when 18% of the peptide bonds are split, whereas Pronase-P brings about a rupture of 33% of these bonds. The latter value agrees approximately with the 42% found by Mr M. Melamed of this Laboratory by using essentially similar conditions for digestion, but by determining the number of α -amino groups produced by formol titration. Nomoto *et al.* (1960c) reported the splitting of about 80% of the peptide bonds in egg albumin by using Pronase-P under different conditions of digestion, and by assaying by the Van Slyke technique. Fig. 1 shows that prior digestion of the protein with papain offers no advantage in this respect over the use of Pronase-P alone. Kakiyama & Schmid (1962) and Eylar (1962) observed that Pronase-P may cause a partial splitting of the carbohydrate-protein bond in orosomucoid, but we have not observed such a cleavage with egg albumin.

Fig. 1. Rate and extent of hydrolysis of egg albumin by various enzymic procedures. ▲, Pepsin followed by trypsin plus chymotrypsin (the latter enzymes were added at point A); ●, Pronase-P; ■, papain followed by Pronase-P (the latter enzyme was added at point B). For the sake of clarity, only a few of the experimentally determined values have been plotted.

Purification of glycopeptides. The initial stage in the purification of the enzymic digest involved a precipitation of the glycopeptides in the presence of 75–80 % (v/v) ethanol. Approximately 70 % of the total nitrogen remained in the supernatant, whereas the glycopeptide was precipitated almost quantitatively. In method 1 further purification of the digest was achieved by precipitation of basic amino acids and large peptides with phosphotungstic acid. The two steps removed 92 % of the nitrogen in the original digest.

Application of the partially purified glycopeptides in method 1 to Sephadex G-25 (Porath & Flodin, 1959) and elution with 15 mM-lithium bromide gave a further purification of glycopeptide (see Fig. 2a). Thus the material in fractions 28–33 had a nitrogen content of about 8.5 atoms of nitrogen/5 molecules of mannose and contained a total of about 1.2 g. of mannose. Rechromatography of this material on Sephadex G-25 gave glycopeptide containing 7.23 atoms of nitrogen/5 molecules of mannose (0.53 g. of mannose). The nitrogen:mannose ratio of this material was unchanged on further chromatography on Sephadex G-25 or on column electrophoresis in 2N-acetic acid. Since the glycopeptide containing aspartic acid as the only amino acid constituent would have a nitrogen:mannose molar ratio of 5:5, it was inferred that at least two other amino acids were present. This was confirmed by quantitative amino acid analysis (Table 1: column A) which showed that one glycopeptide unit contained 3.63 equiv. of amino acid (or nitrogen). This leaves 3.60 atoms of nitrogen in the glycopeptide molecule not present as amino acids. This glycopeptide gives almost exactly 1 equiv. of ammonia on mild acid hydrolysis (Marks *et al.* 1963). The possibility of the presence of a fourth glucosamine residue (Johansen *et al.* 1960) is now excluded. The glycopeptide penetrates the pores of the Sephadex since it is eluted much later than carboxyhaemoglobin (Fig. 2a).

In the purification of glycopeptide from the Pronase digestion (method 2) the phosphotungstic acid used in method 1 was omitted and the material from the ethanol digest placed directly on a Sephadex G-25 column previously equilibrated with 0.1N-acetic acid. The results of the elution are shown in Fig. 2b, and it is clear that a very effective purification of glycopeptide is achieved by this procedure. Moreover, an essentially quantitative recovery of the glycopeptide was achieved. Quantitative amino acid analysis (Table 1: column C) showed that, in addition to aspartic acid, leucine was the only amino acid present in appreciable amount.

The papain–Pronase digest was purified in a manner identical with that used for the Pronase digest. Recoveries of purified glycopeptide were about 90 %. The quantitative amino acid analyses (Table 1: column E) showed that tyrosine, glutamic

acid and lysine were present in glycopeptides prepared by this procedure.

Sequence of amino acids in the glycopeptide. It is clear from the results of various workers that the amino acid sequence in the vicinity of the carbohydrate may be represented as shown (I),

Fig. 2. Purification of glycopeptide from the enzymic digests by using Sephadex G-25. (a) Digest obtained by method 1 (pepsin followed by trypsin plus chymotrypsin): ●, mannose values; ▲, atoms of nitrogen/5 molecules of mannose; ×, carboxyhaemoglobin elution in a prior experiment. Some of the experimentally determined values have been omitted for the sake of clarity. (b) Digest obtained by method 2 (Pronase-P): ●, mannose values; ○, Moore & Stein (1948) ninhydrin values.

Table 1. *Amino acid content of purified egg-albumin glycopeptide*

Experimental details are given in the text. The results are expressed as moles of amino acid/5 moles of mannose; *A*, *C* and *E* were obtained without, and *B* and *D* after, carboxypeptidase treatment of the benzyloxycarbonyl-substituted glycopeptide.

Amino acid Protein digested with ...	Pepsin followed by trypsin + chymotrypsin		Pronase-P		Papain followed by Pronase-P
	<i>A</i>	<i>B</i>	<i>C</i>	<i>D</i>	<i>E</i>
Aspartic acid	1.02	1.01	0.96	0.94	1.0
Leucine	1.01	0.07	0.56	0.20	0.13
Threonine	0.67	0.11	0.22	0.085	0.15
Serine	0.73			0.02	0.14
Glutamic acid	0.07	0.04	0.03	0.02	0.33
Glycine	0.07	—	—	0.07	0.12
Valine	0.06	0.04	0.06	—	—
Tyrosine	—	—	0.05	0.05	0.53
Alanine	—	—	—	—	0.14
Isoleucine	—	—	—	—	0.04
Lysine	—	—	—	—	Present*

* Qualitative identification only.

where R represents the carbohydrate moiety. Clearly the aim of our experiments is to cleave the Tyr-Asp and the Asp-Leu bonds. Method 1 (pepsin followed by trypsin and chymotrypsin) gives glycopeptide in which there is complete splitting of the Tyr-Asp bond but not of the Asp-Leu bond. Neither papain nor a *Streptomyces* protease preparation (Tai & van Heyningen, 1951) effected further hydrolysis of this glycopeptide, i.e. the Asp-Leu, Leu-Thr and Thr-Ser bonds were resistant to these enzymes while present in the glycopeptide. In contrast, Pronase-P, when acting on the whole protein, split almost completely the Leu-Thr and Thr-Ser bonds and partially the Asp-Leu bond, in addition to completely severing the Tyr-Asp bond. Further, papain cleaved the Asp-Leu bond when acting on the whole protein, confirming the observation of Lee & Montgomery (1962). It seemed therefore that digestion of the whole protein by papain followed by Pronase might split both the Asp-Leu (by papain) and the Tyr-Asp (by Pronase) bonds yielding glycopeptide with aspartic acid as the only amino acid. However, cleavage of the Asp-Leu bond in the protein by papain rendered the Tyr-Asp bond less susceptible to the action of Pronase. The latter bond, however, was obviously partly hydrolysed. It was estimated that material prepared by this procedure (method 3) contains about 40–50 % of glycopeptide containing aspartic acid as the only amino acid constituent. No attempt was made to fractionate this mixture as easier methods became available for isolating glycopeptide containing only aspartic acid.

One is tempted to suggest that the carbohydrate moiety causes steric hindrance to the attack of further enzymes. However, it may be that the presence of an undesirable charged group in the

neighbourhood of the bond where rupture is desired (see, e.g., Léonis, Li & Chung, 1959) may be responsible for the non-susceptibility of the glycopeptides to further enzymic action. Thus, e.g., papain has a lower activity on substrates with a carboxyl group in a highly ionized form (Kimmel & Smith, 1954).

Carbohydrate-aspartic acid complex. In general, it has proved difficult to prepare a glycopeptide containing aspartic acid as the only amino acid. Glycopeptides containing several amino acids in which the α -amino group of aspartic acid was free (Johansen *et al.* 1961) were hydrolysed by carboxypeptidase, releasing serine, threonine and leucine. However, after purification, the resultant glycopeptide almost invariably still contained significant amounts of serine, threonine and leucine. The glycopeptide at this stage was not susceptible to the further action of carboxypeptidase. Kaverzneva & Bogdanov (1961) and Nuenke & Cunningham (1961) have reported similar findings.

Bogdanov, Kaverzneva & Tsi De-Fan (1961) indicated that combination of such a glycopeptide with the benzyloxycarbonyl group would render it susceptible to the action of carboxypeptidase. We tried this procedure on preparations of glycopeptide prepared by methods 1 and 2. Quantitative amino acid analyses (Table 1: columns *B* and *D*) indicate that the amino acids other than aspartic acid, although not completely removed, were very substantially reduced in quantity. These analyses were carried out after the removal of the benzyloxycarbonyl group. It is clear that the best method of preparing the glycopeptide with only aspartic acid is by digestion with Pronase-P and purification with Sephadex G-25 (method 2). This is followed by substitution with the benzyloxycarbonyl group,

digestion with carboxypeptidase and the removal of the benzyloxycarbonyl group.

It seems probable that the insusceptibility of the non-acylated glycopeptide to the action of carboxypeptidase is due to the presence of a partly positively charged amino group. Thus carboxypeptidase does not hydrolyse *C*-terminal arginine or lysine residues (Smith, 1954) or, in general, dipeptides (Hofmann & Bergmann, 1940). The latter are resistant owing to the presence of a positively charged amino group (Elkins-Kaufmann & Neurath, 1949; Yanari & Mitz, 1957).

The benzyloxycarbonyl-glycopeptide containing aspartic acid as the only amino acid was obtained as a solid which exhibited birefringence. However, the actual crystal form could not be recognized.

After the removal of the benzyloxycarbonyl group, the glycopeptide had $[\alpha_D^{25}] + 70.0^\circ$ (*c* 1.0, with respect to mannose, in water). Under the particular conditions that we have employed, this glycopeptide gave a brown spot with ninhydrin on paper. This probably indicates a β -aspartamido linkage. Thus, although β -aspartylpeptides have been reported as giving blue spots with ninhydrin on paper (Liwschitz & Zilkha, 1954, 1955, 1957; Liwschitz, Edlitz-Pfeffermann & Lapidoth, 1957; Le Quesne & Young, 1952; John & Young, 1954; Bryant, Moore, Pimlott & Young, 1959; Buchanan, Haley & Markiw, 1962), this appears to be only under very specific conditions (Bryant *et al.* 1959). *L*- β -Aspartyl-*L*-histidine has been reported as giving a yellow-brown colour with ninhydrin (Kakimoto & Armstrong, 1961), but under conditions where amino acids and peptides show up in a variety of different colours (Levy & Chung, 1953). However, β -aspartyl-*L*- and -*D*-valine give brown colours on heating in water with ninhydrin in the presence of acetate (Le Quesne & Young, 1952), as also do 1- β -(*L*-aspartamido)-1-deoxy-*D*-glucose [*N*-(*L*- β -aspartyl)- β -*D*-glycosylamine] and 2-(*L*- β -aspartamido)-2-deoxy-*D*-glucose [*N*-(*L*- β -aspartyl)-*D*-glucosamine] (Marks & Neuberger, 1961).

We have always dried our chromatograms before dipping them in 1% (w/v) ninhydrin in acetone containing 2% (v/v) of pyridine. The papers were then developed by heating at 80° for about 5 min. Under these conditions brown spots are given by asparagine, 1-(*L*- β -aspartamido)-1-deoxy-*D*-glucose, 2-acetamido-1- β -(*L*- β -aspartamido)-1,2-dideoxy-*D*-glucose (Marks, Marshall & Neuberger, 1963), 2-(*L*- β -aspartamido)-2-deoxy-*D*-glucose and the highly purified glycopeptide. All other commonly occurring amino acids (except proline) give a purple colour under these conditions. After 2-3 days, especially when warm, even the colour produced by these β -aspartyl compounds becomes bluish-brown.

Periodate oxidation. Fig. 3, obtained by plotting the periodate reduced against the time of oxidation,

and Fig. 4, obtained by plotting the formic acid release at 5-6° against the time of oxidation by periodate, show that the reaction takes place in two distinct phases. An initial rapid reaction is followed by a second phase in which the reaction still continues but at a much lower rate. A correction for the over-oxidation can be obtained by a simple extrapolation to zero time. One glycopeptide unit (containing 5 mannose and 3 glucosamine residues) reduced 5.05 molecules of sodium metaperiodate (Fig. 3) and yielded 0.7 molecule of formic acid (Fig. 4).

The presence of *N*-acetyl groups on the glucosamine residues in the glycopeptide limits the consumption of periodate to, at most, 1 molecule of periodate/amino sugar residue and prevents the liberation of formic acid on periodate oxidation (Hough & Taha, 1956). Consequently the formic acid liberated from the glycopeptide must be derived from a (1 $\rightarrow$ 6)-linked mannose residue or a non-reducing mannose end group. To decide between these two alternatives the glycopeptide

Fig. 3. Oxidation of egg-albumin glycopeptide by periodate at 5-6°, showing the amount of periodate used by 14.6 μmoles of glycopeptide after various times.

Fig. 4. Amount of formic acid produced/μmole of glycopeptide after various times of oxidation of the glycopeptide by periodate at 5-6°.

was treated with a preparation of almond emulsin, kindly given to us by Dr W. J. Whelan, who prepared it by the procedure of Tauber (1932-33). In this experiment mannose was detected by paper chromatography, showing that a mannose residue occurs as a non-reducing end group and is the source of the formic acid liberated. This is in accordance with the identification of 2,3,4,6-tetra-*O*-methyl-D-mannose among the products of exhaustive methylation of egg albumin in the experiments of Bragg & Hough (1961). Further, since less than 1 molecule of formic acid was liberated from 1 glycopeptide unit, the possibility of any (1→6)-linked mannose residues occurring in the glycopeptide can be excluded.

Further experiments were directed to determining the number of mannose residues in the glycopeptide which had survived oxidation by periodate. The specific method for mannose (François *et al.* 1962) was used for unambiguous determination. The method involves the hydrolysis of the mannose-containing polymer in the presence of radioactive mannose, isolation of the sugar as the phenylhydrazone and determination of the radioisotope dilution of the radioactive mannose. By this method it was shown that 53% of the mannose contained in the glycopeptide was lost during periodate oxidation. In independent studies by Lee & Montgomery (1961) and Bragg & Hough (1961) losses of 51 and 69% respectively of the original mannose were found after periodate oxidation of egg albumin.

The colorimetric method of Rondle & Morgan (1955) was used on an acid hydrolysate (4*N*-hydrochloric acid for 3 hr.) to determine the amount of glucosamine remaining intact in the glycopeptide after periodate oxidation for 2, 6 and 24 hr. After these respective time-intervals, 85, 77.8 and 70.4% respectively of the glucosamine remained intact. Since 1 glycopeptide unit contains 3 *N*-acetylglucosamine molecules it is clear that only one of these molecules is attacked by periodate. Lee & Montgomery (1961) and Bragg & Hough (1961) have also observed the oxidation of only one of the three *N*-acetylglucosamine residues in egg albumin by periodate.

Since 2.7 of the 5 mannose residues in a glycopeptide unit are attacked by periodate and one mannose is a non-reducing end group, 3.7 molecules of periodate will be reduced by the mannose of the glycopeptide unit. The *N*-acetylglucosamine would account for the reduction of 1 molecule of periodate so that 1 glycopeptide unit would be expected to reduce 4.7 molecules of periodate. This value is in fair agreement with the finding that 1 glycopeptide unit reduced 5.05 molecules of periodate.

Since only one mannose residue can be in a non-reducing terminal position and the oxidized *N*-

acetylglucosamine residue must be either in a non-reducing terminal position or be (1→6)-linked, the carbohydrate moiety must be either linear or have a maximum of one branch point.

SUMMARY

1. The rate and extent of enzymic hydrolysis of egg albumin was studied for three systems: (a) pepsin followed by trypsin and chymotrypsin together, (b) Pronase-P and (c) papain followed by Pronase-P.

2. Techniques involving the use of Sephadex G-25 are described for the isolation of glycopeptides from these enzymic digests. Yields of glycopeptide by methods (b) and (c) are very high.

3. The amino acid constituents in these purified glycopeptides were compared. These were used to point out the changes in specificity of the enzyme depending on whether the whole protein or the glycopeptide is the substrate.

4. A method suggested by Bogdanov *et al.* (1961) for obtaining glycopeptide with aspartic acid as the only amino acid constituent has been confirmed and the usefulness of the procedure was demonstrated.

5. The carbohydrate-aspartic acid complex obtained gave a brown colour when treated on paper with ninhydrin under the conditions described. The significance of this is discussed.

6. Quantitative studies of periodate uptake by a purified glycopeptide preparation were made. The formic acid liberated and the quantities of mannose and glucosamine oxidized were also measured. These studies make it probable that the carbohydrate is either linear or has a maximum of one branch point.

We thank Dr N. M. Green for his gift of the *Streptomyces* protease preparation. The technical assistance of Mr R. Cooper is gratefully acknowledged. Acknowledgement is made for grants from the Medical Research Council (Junior Research Fellowship) (to A.P.F.), The British Empire Cancer Campaign (to G.S.M.) and the U.S. Public Health Service (to R.D.M.).

REFERENCES

- Aminoff, D. & Morgan, W. T. J. (1951). *Biochem. J.* **48**, 74.
- Bogdanov, V. P., Kaverzneva, E. D. & Tsi De-Fan (1961). *Abstr. 5th int. Congr. Biochem., Moscow*, p. 25.
- Bragg, P. D. & Hough, L. (1961). *Biochem. J.* **78**, 11.
- Bryant, P. M., Moore, R. H., Pimlott, P. J. & Young, G. T. (1959). *J. chem. Soc.* p. 3868.
- Buchanan, D. L., Haley, E. H. & Markiw, R. T. (1962). *Biochemistry*, **1**, 612.
- Calvery, H. O. (1933). *J. biol. Chem.* **102**, 73.
- Chatterjee, A. K., Durant, G. J., Lee, Y. C. & Montgomery, R. (1961). *Biochem. biophys. Res. Commun.* **4**, 425.
- Chibnall, A. C., Rees, M. W. & Williams, E. F. (1943). *Biochem. J.* **37**, 354.

- Elkins-Kaufmann, E. & Neurath, H. (1949). *J. biol. Chem.* **178**, 645.
- Eylar, E. H. (1962). *Biochem. biophys. Res. Commun.* **8**, 195.
- Farthing, A. C. (1950). *J. chem. Soc.* p. 3213.
- François, C., Marshall, R. D. & Neuberger, A. (1962). *Biochem. J.* **83**, 335.
- Greenstein, J. P. & Winitz, M. (1961). *Chemistry of the Amino Acids*, vol. 2, p. 891. New York: John Wiley and Sons Inc.
- Hofmann, K. & Bergmann, M. (1940). *J. biol. Chem.* **134**, 225.
- Hough, L. & Taha, M. I. (1956). *J. chem. Soc.* p. 2042.
- Isbell, H. S., Frush, H. L. & Schaffer, R. (1955). *J. Res. nat. Bur. Stand.* **54**, 201.
- Jackson, E. L. (1944). In *Organic Reactions*, vol. 2, p. 341. Ed. by Adams, R., Bachmann, W. E., Fieser, L. F., Johnson, J. R. & Snyder, H. R. New York: John Wiley and Sons Inc.
- Jevons, F. R. (1958). *Nature, Lond.*, **181**, 1346.
- Johansen, P. G., Marshall, R. D. & Neuberger, A. (1958). *Nature, Lond.*, **181**, 1345.
- Johansen, P. G., Marshall, R. D. & Neuberger, A. (1960). *Biochem. J.* **77**, 239.
- Johansen, P. G., Marshall, R. D. & Neuberger, A. (1961). *Biochem. J.* **78**, 518.
- John, W. D. & Young, G. T. (1954). *J. chem. Soc.* p. 2870.
- Kakimoto, Y. & Armstrong, M. D. (1961). *J. biol. Chem.* **236**, 3280.
- Kakiyama, S. & Schmid, K. (1962). *Biochim. biophys. Acta*, **58**, 80.
- Kaverzneva, V. P. & Bogdanov, E. D. (1961). *Biokhimiya*, **26**, 105.
- Kimmel, J. R. & Smith, E. L. (1954). *J. biol. Chem.* **207**, 515.
- Kolthoff, I. M. & Sandell, E. B. (1952). *Textbook of Quantitative Inorganic Analysis*, 3rd ed., p. 587. New York: The Macmillan Co.
- Lee, Y. C. & Montgomery, R. (1961). *Arch. Biochem. Biophys.* **95**, 263.
- Lee, Y. C. & Montgomery, R. (1962). *Arch. Biochem. Biophys.* **97**, 9.
- Léonis, J., Li, C. H. & Chung, D. (1959). *J. Amer. chem. Soc.* **81**, 419.
- Le Quesne, W. J. & Young, G. T. (1952). *J. chem. Soc.* p. 24.
- Levy, A. L. & Chung, D. (1953). *Analyt. Chem.* **25**, 396.
- Lineweaver, H. & Schummer, S. (1941). *Enzymologia*, **10**, 81.
- Liwschitz, Y., Edlitz-Pfeffermann, Y. & Lapidot, Y. (1957). *J. Amer. chem. Soc.* **79**, 3069.
- Liwschitz, Y. & Zilkha, A. (1954). *J. Amer. chem. Soc.* **76**, 3698.
- Liwschitz, Y. & Zilkha, A. (1955). *J. Amer. chem. Soc.* **77**, 1265.
- Liwschitz, Y. & Zilkha, A. (1957). *J. chem. Soc.* p. 4394.
- Malpress, F. H. & Hytten, F. E. (1958). *Biochem. J.* **68**, 708.
- Marks, G. S., Marshall, R. D. & Neuberger, A. (1962). *Biochem. J.* **84**, 30 p.
- Marks, G. S., Marshall, R. D. & Neuberger, A. (1963). *Biochem. J.* **87**, 274.
- Marks, G. S., Marshall, R. D., Neuberger, A. & Papkoff, H. (1962). *Biochim. biophys. Acta*, **63**, 340.
- Marks, G. S. & Neuberger, A. (1961). *J. chem. Soc.* p. 4872.
- Marshall, R. D. & Neuberger, A. (1960). *Nature, Lond.*, **186**, 311.
- Moore, S. & Stein, W. H. (1948). *J. biol. Chem.* **176**, 367.
- Moore, S. & Stein, W. H. (1954). *J. biol. Chem.* **211**, 907.
- Nomoto, M. & Narahashi, Y. (1959a). *J. Biochem., Tokyo*, **46**, 653.
- Nomoto, M. & Narahashi, Y. (1959b). *J. Biochem., Tokyo*, **46**, 839.
- Nomoto, M. & Narahashi, Y. (1959c). *J. Biochem., Tokyo*, **46**, 1481.
- Nomoto, M. & Narahashi, Y. (1959d). *J. Biochem., Tokyo*, **46**, 1645.
- Nomoto, M., Narahashi, Y. & Murakami, M. (1960a). *J. Biochem., Tokyo*, **48**, 453.
- Nomoto, M., Narahashi, Y. & Murakami, M. (1960b). *J. Biochem., Tokyo*, **48**, 593.
- Nomoto, M., Narahashi, Y. & Murakami, M. (1960c). *J. Biochem., Tokyo*, **48**, 906.
- Nuenke, R. H. & Cunningham, L. W. (1961). *J. biol. Chem.* **236**, 2452.
- Partridge, S. M. (1949). *Nature, Lond.*, **164**, 443.
- Porath, J. (1956). *Biochim. biophys. Acta*, **22**, 151.
- Porath, J. & Flodin, P. (1959). *Nature, Lond.*, **183**, 1657.
- Rondle, C. J. M. & Morgan, W. T. J. (1955). *Biochem. J.* **61**, 586.
- Rosen, H. (1957). *Arch. Biochem. Biophys.* **67**, 10.
- Smith, E. L. (1954). In *The Chemical Structure of Proteins*, p. 109. Ed. by Wolstenholme, G. E. W. & Cameron, M. P. London: J. and A. Churchill Ltd.
- Smith, M. A. & Willeford, B. R., jun. (1954). *Analyt. Chem.* **26**, 751.
- Tai, T. Y. & van Heyningen, W. E. (1951). *J. gen. Microbiol.* **5**, 110.
- Tauber, H. (1932-33). *J. biol. Chem.* **99**, 257.
- Van Slyke, D. D. & Rieben, W. K. (1944). *J. biol. Chem.* **156**, 743.
- Yamashina, I. & Makino, M. (1962). *J. Biochem., Tokyo*, **51**, 359.
- Yanari, S. & Mitz, M. A. (1957). *J. Amer. chem. Soc.* **79**, 1154.
- Yemm, E. W. & Cocking, E. C. (1955). *Analyst*, **80**, 209.