deacylation of the amino-acyl complex (Nathans & Lipmann, 1961). In the crude rat-liver 15s preparations, however, the retention of counts on the transfer RNA resulting from inhibition by chlortetracycline is less striking and in the analogous *E. coli* preparations there is no increase in the radioactivity of the transfer RNA over the control values. The enzymic deacylation of the transfer RNA-amino acid complex described by Nathans & Lipmann (1961) may be more rapid in the cruder preparations. It is not clear at present whether chlortetracycline prevents part or all of the transfer RNA from becoming attached to the ribosomal RNA since Bloemendal & Bosch (1962) have shown that this reaction is not dependent on amino acid transfer.

It appears that the rat-liver preparations may be less sensitive to the tetracyclines than those from *E. coli*. In this context, Rendi & Ochoa (1961) have reported briefly that oxytetracycline inhibited the transfer of amino acid from transfer RNA to the ribosomal protein of a bacterial preparation but was without effect on the same reaction in a corresponding rat-liver system. Since 400 μ M-oxytetracycline is only 27% inhibitory in the rat-liver system described here, as compared with 86% inhibitory action in the bacterial preparation, it may be that the drug concentration used by Rendi & Ochoa (1961) was too low for an inhibitory effect to be observed in their rat-liver system. The reason for the apparent difference in sensitivity to the tetracyclines of the animal and bacterial systems is not clear.

SUMMARY

1. Chlortetracycline has been shown to reduce the incorporation of [14 C]leucine into the ribosomal protein of both rat-liver and *Escherichia coli* cell-free preparations *in vitro*.

2. In neither the animal nor bacterial system did the drug interfere with the attachment of the amino acid to the transfer RNA.

3. In both preparations the drug prevented the amino acid of the transfer RNA-amino acid complex from transferring to the ribosomal protein. In highly purified systems the amino acid not transferred to the ribosome was largely retained on the transfer RNA.

4. Oxytetracycline and tetracycline also inhibited the transfer of amino acid to the ribosomal protein in both animal and bacterial systems. The rat-liver preparations were apparently less sensitive to the tetracyclines than those from *E. coli*. Chloramphenicol, inhibitory to the *E. coli* system, was without effect on the rat-liver system.

I am grateful to Mr A. Godfrey for valuable technical assistance.

REFERENCES

- Bollum, F. J. (1959). *J. biol. Chem.* **234**, 2733.
Bloemendal, H. & Bosch, L. (1962). *Biochem. J.* **84**, 92r.
Gale, E. F. & Folkes, J. P. (1953). *Biochem. J.* **53**, 493.
Hahn, F. E. (1958). *Proc. 4th int. Congr. Biochem., Vienna, Symp.* no. 5, p. 104.
Hoagland, M. B., Stephenson, M. L., Scott, J. F., Hecht, L. I. & Zamecnik, P. C. (1958). *J. biol. Chem.* **231**, 241.
Korner, A. (1959). *Biochem. J.* **73**, 61.
Korner, A. (1961). *Biochem. J.* **81**, 168.
Layne, E. (1957). In *Methods in Enzymology*, vol. 3, p. 450. Ed. by Colowick, S. P. & Kaplan, N. O. New York: Academic Press Inc.
Mans, R. J. & Novelli, G. D. (1961). *Arch. Biochem. Biophys.* **94**, 48.
Nathans, D. & Lipmann, F. (1961). *Proc. nat. Acad. Sci., Wash.*, **47**, 497.
Nikolov, T. K. & Ilkov, A. T. (1961). *Abstr. Commun. 5th int. Congr. Biochem., Moscow*, p. 44.
Nirenberg, M. W. & Matthaei, J. H. (1961). *Proc. nat. Acad. Sci., Wash.*, **47**, 1580.
Rendi, R. & Ochoa, S. (1961). *Science*, **133**, 1367.

Biochem. J. (1963) **87**, 453

Adenosine Triphosphate-Creatine Phosphotransferase from Ox Brain: Purification and Isolation

By T. WOOD

*Department of Biochemistry, Institute of Psychiatry (British Postgraduate Medical Federation,
University of London), Maudsley Hospital, London, S.E. 5*

(Received 16 November 1962)

Adenosine triphosphate-creatine phosphotransferase (EC 2.7.3.2) plays an important part in cerebral metabolism by maintaining concentrations of ATP under resting conditions and, after activity

such as that after electrical stimulation, restoring these concentrations at the expense of the phosphocreatine of the tissue (McIlwain, 1959). Although the enzyme from skeletal muscle has

been intensively studied (Kuby, Noda & Lardy, 1954*a, b, c*; Ennor & Rosenberg, 1954; DaCosta & Friedburg, 1960), little attention has been directed to that of other tissues, although the enzyme is usually assumed to be present in those tissues that contain phosphocreatine (Ennor & Morrison, 1960). The enzyme from fowl visceral muscle was studied by Geiger (1956), and a preliminary investigation of the guinea-pig-brain enzyme was carried out by Narayanaswami (1952). The present paper describes the isolation of the enzyme from ox brain before study of its properties, and amplifies the preliminary account given by Wood (1962).

MATERIALS

Ox brain. Brains were removed from animals within 30 min. of slaughter and packed in ice for transport to the laboratory. On arrival, the dura was removed, any superficial blood washed away, and each brain placed in a polythene bag and refrigerated at -20° . Brains were used for preparation of the enzyme after storage for up to 8 weeks and no loss of activity attributable to storage was ever detected. Extracts from a frozen brain had the same activity as those prepared from the fresh brain.

Reagents for preparation of the enzyme. Water was distilled once from metal and once from glass. Acetone, potassium chloride, EDTA (dipotassium salt), magnesium acetate, magnesium sulphate, ammonium sulphate, sodium acetate and acetic acid were all Hopkin and Williams Ltd. A.R. grade.

Other reagents. Creatine and sodium thioglycollate were recrystallized from a commercial preparation. ATP (disodium salt, tetrahydrate) was obtained from Schwarz BioResearch Inc., Orangeburg, N.Y., U.S.A. Glycine was Hopkin and Williams Ltd. A.R. grade. Glutathione was from The Distillers Co. Ltd. (Biochemicals), Speke, Liverpool 19. L-Cysteine hydrochloride was from British Drug Houses Ltd., Poole, Dorset. Crystalline bovine serum albumin was from Armour Laboratories, Eastbourne. Ponceau S and Amidochwarz 10B were obtained from G. T. Gurr Ltd., London, S.W. 6.

Media for electrophoresis and chromatography. Hydrolysed starch was the product of the Connaught Medical Laboratories, Toronto, Canada. Oxoid cellulose acetate electrophoresis strips were from Oxo Ltd., London, S.E. 1. Diethylaminoethyl-Sephadex (DEAE-Sephadex) was grade A50-Medium supplied by Pharmacia, Uppsala, Sweden.

METHODS

Determinations

Measurement of enzyme activity. Enzyme activity was measured as described by Noda, Kuby & Lardy (1955), except that sodium thioglycollate was added from a 20 mM stock solution, pH 7.0, to give a final concentration in the incubation mixture of 1 mM. Samples were withdrawn from the reaction mixture at 0, 5 and 15 min. by a 1 ml. all-glass tuberculin syringe.

The activity was calculated by plotting $x/(1-x)$ against t , where x is the number of μ moles of phosphocreatine

appearing/ml. of reaction mixture in time t min. The gradient of the straight line obtained gave the apparent second-order velocity constant, K' , in ml./ μ mole/min. One unit of enzyme is defined as that amount of enzyme/ml. of reaction mixture, containing initially ATP (1 mM), creatine (24 mM) and magnesium acetate (6 mM), at 30° and pH 9.0 (96 mM-glycine-NaOH buffer), that will catalyse the phosphotransferase reaction with an apparent second-order velocity constant, K' , of 1.0.

For measurement of the activities of fractions eluted from the chromatographic column, a 'rapid assay' procedure was developed in which the amount of enzyme present was calculated from the amount of phosphate transferred from ATP to creatine during incubation for 5 min. A standard curve for the 'rapid assay' was constructed from the results of a number of previous full-length assays on various preparations, by plotting the amount of phosphate transferred in 5 min. against the number of enzyme units/ml. of incubation mixture (Fig. 1). Apart from being only an approximate measure of the enzyme present, the 'rapid assay' had the disadvantage that there was no criterion of the validity of the assay comparable with the requirement in the full-length assay that a plot of the parameter $x/(1-x)$ against the times of 0, 5 and 15 min. should lead to a straight line through the origin (Kuby *et al.* 1954*a*).

Activation by thiols. Stock solutions of L-cysteine, glutathione and sodium thioglycollate were prepared, adjusted to pH 7.0 with N-NaOH and stored at 5° . Suitable dilutions of these were added to the assay mixture before use. Caution was necessary in the use of thioglycollate, since a 2-month-old solution, which smelled perceptibly of mercaptans, caused 85% inhibition.

Protein determinations. The biuret procedure described by Layne (1957) was used with crystalline bovine serum albumin as a standard. Protein was measured in effluents from chromatographic columns by determining the extinction against water at 280 m μ . For the pure enzyme, in a solution containing 1.15 mg./ml. by the biuret procedure, the extinction measured at 280 m μ in a cell of light-path 1 cm. was taken as unity.

Fig. 1. Standard curve for the 'rapid assay' procedure for the determination of adenosine triphosphate-creatine phosphotransferase. Experimental details are given in the text.

Measurement of pH. Measurements were made on a model 23A pH-meter manufactured by Electronic Instruments Ltd., Richmond, Surrey.

Electrophoresis

Electrophoresis on cellulose acetate. Cellulose acetate strips (20 cm. $\times$ 5 cm.) were marked with a line parallel to the short edge of the strip, 6 cm. from one end and extending to within 1 cm. of each edge. After being soaked overnight at 2° in barbiturate buffer, pH 8.6 and *I* 0.05 (Kohn, 1960), the strips were blotted and the sample was applied along the pencilled line from a capillary pipette. The strip was placed in the Shandon Universal electrophoresis tank (Shandon Scientific Co., London, S.W. 7) with the starting line nearest the cathode and electrophoresis carried out at 2° for 2½ hr. at 200V, as described by Kohn (1960). On completion of the run, the strips were blotted and stained with 0.2% Ponceau S in 10% (w/v) trichloroacetic acid and washed in 5% (v/v) acetic acid.

Electrophoresis in starch gel. Electrophoresis was carried out at room temperature in an EEL electrophoresis tank (Evans Electroselenium Ltd., Halstead, Essex), by using the discontinuous buffer system of Poulik (1957) and following the general procedure recommended by Bodman (1960). Starch gels (11.8%, w/v) were prepared in 76 mm. tris-50 mm. citrate, pH 9.0, and cast in Perspex trays. The bridge buffer was 0.30M-boric acid-0.06N-NaOH, pH 9.0. A current of 3 mA/cm. width of tray (fully occupied by sample inserts) was passed for 3 hr. Gels were stained with 1% (w/v) Amidoschwarz 10B in methanol-acetic acid-water (15:5:2, by vol.) and washed in the same solvent.

Purification of the enzyme

Extraction and preliminary purification of the enzyme. The procedure was based on that of Noda *et al.* (1955) for the preparation of the enzyme from rabbit muscle, but extensive modifications were necessary before the method could be applied to brain tissue.

One whole ox brain was removed from the deep-freeze, weighed and allowed to warm at room temperature until it was possible to cut it with a knife. It was then cut into slices 1 cm. thick and allowed to warm to 0°. All further operations were conducted in the cold room at 2°. The sliced brain was homogenized with ice-cold 10 mM-KCl-1 mM-EDTA (2 ml./g. of tissue) in a top-drive macerator (Townson and Mercer Ltd., Croydon, Surrey) in short bursts to produce an even homogenate. The homogenate was stirred for 15 min. and allowed to warm to 0° if necessary. It was then centrifuged for 90 min. at 14 000g (all *g* values are average *g*) in polypropylene bottles. The supernatant was poured off, its volume and pH (6.4-6.7) were measured, and the pH was adjusted to 9.0 with 0.5N-NaOH. The solution was then transferred to 1 l. glass-stoppered bottles in the cold room, stirred magnetically, and an equal volume of acetone was added from a fine glass jet dipping below the surface of the liquid (Askonas, 1951). The 50% (v/v) acetone mixture was stirred at 2° for 3-4 days, by which time 80-90% of the protein had flocculated, leaving a water-clear supernatant. The mixture was centrifuged for 30 min. at 10 000g in polypropylene bottles, and the clear supernatant was placed in 1 l. glass-stoppered bottles, stirred magnetically, and a volume of acetone was added equal to that of the

magnesium sulphate solution to be added subsequently. 2.0M-Magnesium sulphate solution was then added slowly from a pipette to give a final concentration of 20 mM. The solution was stirred for 4-5 hr., by which time an abundant flocculent precipitate had appeared in suspension and a crystalline deposit consisting mainly of inorganic salts had formed on the bottom of the bottles. The protein suspension was poured off without disturbing the deposit and centrifuged for 10 min. at 500g. No attempt was made to dissolve the small amount of enzyme (less than 5% of the total) adhering to the crystalline deposit since the high salt content of the resulting solution interfered with the acetone fractionation at the next stage. The sedimented protein was extracted with 70 mM-magnesium acetate, pH 9.0 (6% of the volume of the original extract), and the suspension was centrifuged for 30 min. at 800g. The resulting sediment was again extracted with 70 mM-magnesium acetate (4% of the volume of the original extract) and centrifuged as above. The two supernatants were combined and the pH was adjusted to 9.0 with 0.2N-NaOH.

Acetone fractionation. Acetone was added slowly, with stirring, to give a final concentration of 30% (v/v), and the solution, in a stoppered flask, was stirred for 1 hr. The precipitate (0-30% fraction) was centrifuged down at 5000g for 10 min., the supernatant collected and acetone added to give a final concentration of 40% (v/v). The mixture was stirred for 1 hr., and the precipitate (30-40% fraction) and supernatant (> 40% fraction) were collected as above. In some instances acetone was added to the supernatant to give a final concentration of 50% (v/v). The mixture was stirred for 1 hr., the solution was centrifuged and the precipitate (40-50% fraction) was collected as above. On recovery, each precipitate was redissolved immediately in 50 mM-ammonium citrate, pH 9.0, to give a final concentration of 5-10 mg. of protein/ml. and centrifuged until the solution was free of insoluble matter.

Ammonium sulphate fractionation. The 30-40% acetone fraction was diluted with 50 mM-ammonium citrate, pH 9.0, to give a concentration of 5 mg./ml. Solid ammonium sulphate was added gradually, with stirring, to give a concentration of 1.75M (all concentrations calculated at 20°; Dixon, 1953) and the solution stirred for 1 hr. The mixture was centrifuged for 10 min. at 5000g, and the supernatant was poured off and ammonium sulphate added to it to give a concentration of 2.4M. The mixture was stirred for 1 hr., and the precipitate (1.75-2.4M-ammonium sulphate fraction) was centrifuged down and dissolved in 50 mM-ammonium citrate, pH 9.0.

Column chromatography

Column chromatography was carried out at 2° on a column (6 cm. $\times$ 2 cm.) of DEAE-Sephadex and fractions were collected by a Towers automatic fraction collector (J. W. Towers and Co. Ltd., Widnes, Lancs.). A salt gradient was applied to the column from a mixing device consisting of a separating funnel, *B*, inserted through a rubber bung into the neck of a conical flask, *A*, that had a side arm fused to the bottom and contained a magnetic stirrer. The gradient was calculated from the expression:

$$c = C_B - (C_B - C_A) e^{-V/V_A}$$

where *c* is the concentration of solution emerging from the flask when a volume *V* has been delivered, *C_A* is the initial concentration of the solution in the conical flask, *C_B* is the

concentration of the solution in the funnel, V_A is the constant volume of solution in the conical flask, and V is the volume of the solution delivered from the time the tap of the separating funnel was opened. This expression is identical with that obtained by Cherkin, Martinez & Dunn (1953) and further discussed by Lakshmanan & Lieberman (1953).

Procedure. DEAE-Sephadex was prepared in the acetate form, equilibrated with 0.1M-sodium acetate, pH 6.8, and suspended in 50 mM-sodium acetate-1 mM-EDTA, pH 6.8. Just before use, the pH was adjusted to 6.8 and the adsorbent packed into a glass column under gravity. The column was washed with the 'starting buffer' (Peterson & Sober, 1956), consisting of 2 mM-sodium acetate-1 mM-EDTA, pH 7.0, until the pH of the effluent was 6.8-7.0. The enzyme sample (20-100 mg. of the 1.75-2.4M-ammonium sulphate fraction in 10 ml.), previously dialysed overnight against 50 vol. of 'starting buffer', was run on to the column and washed through with 'starting buffer'. After the passage of 30 ml. of this buffer, 0.10M-sodium acetate-1 mM-EDTA, pH 5.5, was run on to the column and allowed to flow through until the pH of the effluent had fallen to 5.5 (about 60 ml. was required). At this point the gradient was started (100 ml. of 0.10M-sodium acetate-1 mM-EDTA, pH 5.5, in the conical flask, 200 ml. of 0.40M-sodium acetate-1 mM-EDTA, pH 5.5, in the funnel).

The average flow rate during the run was 50 ml./hr. Fractions of 5 ml. each were collected in tubes containing 0.6 ml. of M-NaHCO₃, which was sufficient to raise the pH of the fractions above 7.0. Each fraction was mixed and the extinction was measured at 280 m μ against water. Enzyme activity was measured in selected tubes by the 'rapid assay' procedure described above. Samples for electrophoresis were concentrated, when necessary, under negative pressure, by 'ultrafiltration-dialysis', as described by Everall & Wright (1958) against 1.7 mM-ammonia-1 mM-EDTA (pH about 9-10). Material from the column was prepared for rechromatography by combining the contents of a number of tubes, concentrating by ultrafiltration-dialysis and dilution of the concentrate with 2 vol. of 'starting buffer'.

Investigation of fractions from the column. Samples of enzyme from the column were taken and, without further treatment, initial reaction velocities were measured as described by Kuby *et al.* (1954*b*) over a range of ATP concentrations. Michaelis constants and maximum velocities were determined from a Lineweaver & Burk (1934) plot. Measurements of sedimentation coefficients were carried out in the ultracentrifuge on the unconcentrated material. Assays for D-2-phosphoglycerate hydro-lyase (EC 4.2.1.11) were performed on a number of fractions by the method of Malmström (1957).

RESULTS

Measurement of enzyme activities. Following the procedure of Noda *et al.* (1955) a plot of $x/(1-x)$ versus time (where x is the number of μ moles of phosphate transferred to creatine) gave good straight lines through the origin, provided that the enzyme concentration in the reaction mixture was less than 0.04 unit/ml. A correction was made for adenosine-triphosphatase activity by running a sample, in parallel, with creatine omitted from the

reaction mixture. Adenosine-triphosphatase activity was negligible at all stages of the purification procedure after the preparation of the initial extract. Part of the small adenosine-triphosphatase activity of the initial extract may have been due to small amounts of creatine introduced into the assay mixture with the enzyme, since traces of creatine in the diluted enzyme solution taken for the assay were detectable by paper chromatography. Preliminary experiments showed no loss of enzyme activity occurring when the enzyme, at a dilution of 0.03 unit/ml., was incubated in the absence of thioglycollate, either alone or in the presence of creatine or ATP, for 30 min. at 30°.

Activation by thiols. Cysteine, glutathione and sodium thioglycollate, at concentrations of 1 mM, activated the enzyme to the same extent, whereas 1 mM-EDTA produced only slight activation (Table 1).

Stability of the enzyme. The brain enzyme was not as stable towards heat as the rabbit-muscle enzyme, and heat treatment of a crude brain extract (pH 6.8) for 30 min. at 40° as in the original procedure of Kuby *et al.* (1954*a*) caused a loss of 50% of the activity. Treatment of the initial extract with purified ethanol or propan-1-ol at 0°, reagents to which the muscle enzyme is stable (Kuby *et al.* 1954*a*; Askonas, 1951), resulted in the loss of a large fraction of the enzyme activity. No activity loss could be detected at the stage of the treatment of the initial extract with acetone, other than that due to removal of a small proportion of the enzyme brought down in association with the precipitated protein.

Preparations of the enzyme at a number of stages of purification lost activity on storage at 5° over a period of 1-3 weeks. Thus a preparation of specific activity 15 units/mg. lost activity at the constant rate of 2% per day over a period of 12 days. Loss of activity could not be arrested by storage at -10° in the frozen state, or by storage at 5° in the presence of the equilibrium mixture formed in a solution containing initially creatine (10 mM), ATP (1 mM) and magnesium acetate (8 mM) at pH 9.0 (96 mM-glycine-sodium hydroxide buffer).

The enzyme was stable when exposed to the solution used to elute it from the DEAE-Sephadex column, since no loss of activity was observed when the enzyme was kept for 2 hr. at 0° in 0.2M-sodium acetate buffer, pH 5.5.

Extraction procedure. After homogenization of the tissue and centrifuging at 14 000g, the volume of the supernatant was only 70-90% of the volume of potassium chloride solution added. A second extraction performed with the same volume of potassium chloride solution and blending for the minimum time required to give a homogeneous

dispersion yielded a similar result. After the centrifuging, the residue was again suspended, and samples of the suspension and of the first and second extracts were assayed. The results of a typical experiment are shown in Table 2. In a series of 12 enzyme preparations, involving 200–500 g. of brain, volumes of liquid ranging from 18 to 30 % of the volume of extractant used were absorbed at the first extraction; the same phenomenon occurred when 0.5–2 g. portions of tissue were treated in a similar manner.

A value for the enzyme content of the tissue calculated from the volume and activity of the initial extract would be considerably in error, owing to the unknown amount of enzyme associated with the insoluble residue. It was therefore assumed that the enzyme activity was distributed uniformly through the whole volume of the extractant taken, the fact that only a portion of the latter could be recovered on centrifuging being disregarded. On this basis the values obtained were in good agreement with those determined by two

successive extractions and assay of the suspended residue. Thus, in the example in Table 2, the total activity calculated from 284 ml. of extractant with an activity of 7.6 units/ml. was 2160 units, which compared favourably with a total of 2030 units determined directly.

Enzyme content of the tissue. By using the above basis for calculation, the enzyme content of ox brain was found to be 24 ± 11 units/g. wet wt. (12 preparations).

Purification of the enzyme. The progress of the purification is summarized in Table 3.

Electrophoresis on cellulose acetate and in starch gel. The progress of the purification procedure was followed by electrophoresis of the various fractions on cellulose acetate and in starch gel (Fig. 2). Both methods gave somewhat similar patterns but those in starch gel were more reproducible and all the components moved towards the anode. However, electrophoresis on cellulose acetate was preferred since the procedure was more rapid and the strips could be stained and washed within 5 min. of

Table 1. *Activation of cerebral adenosine triphosphate-creatine phosphotransferase by thiols*

The initial extract was made by extracting frozen ox brain with 10 mM-KCl–1 mM-EDTA (2 ml./g. of tissue); other fractions were those obtained during the purification procedure. The compounds tested were included in the incubation medium, which contained creatine (24 mM), ATP (1 mM), magnesium acetate (8 mM) and glycine, pH 9.0 (100 mM); the temperature was 30°. The reaction was started by the addition of the enzyme. Experimental details are given in the text. The percentage activation was calculated from the rates in two mixtures run in parallel, one containing the activator and one without.

	Specific activity (units/mg. of protein)		Activation (%)
	Before activation	After activation	
(a)			
Initial extract + thioglycollate (1 mM)	0.19	1.7	800
Initial extract + cysteine (1 mM)	0.19	1.7	800
Initial extract + glutathione (1 mM)	0.19	1.7	800
Initial extract + EDTA (1 mM)	0.19	0.25	30
(b) Activation by 1 mM-thioglycollate			
Initial extract	0.5	1.6	220
70 mM-Magnesium acetate extract	8.7	11.2	30
35–40 % (v/v) Acetone fraction	18	22	22
40–45 % (v/v) Acetone fraction	1.5	2.3	54
1.75–2.4 M-(NH ₄) ₂ SO ₄ fraction	16	29	80
Purified enzyme from DEAE-Sephadex column	52	52	0

Table 2. *Distribution of liquid, protein and adenosine triphosphate-creatine phosphotransferase activity after extraction of brain tissue with potassium chloride solution*

Half an ox brain (wt. 142 g.) was homogenized with 284 ml. of 10 mM-KCl–1 mM-EDTA. The homogenate was centrifuged (for 90 min. at 14 000g) and the residue was resuspended in 284 ml. of extractant. After a second centrifuging (for 90 min. at 14 000g) the residue was again resuspended in 284 ml. of extractant. Experimental details are given in the text. N.D., Not determined.

	Volume (ml.)	Activity (units/ml.)	Total activity (units)	Protein (mg./ml.)	Specific activity (unit/mg. of protein)
First extract	225	7.6	1710	8.0	0.95
Second extract	230	1.0	230	2.9	0.35
Residue suspended in extractant	460	0.2	92	N.D.	N.D.

Table 3. *Purification of adenosine triphosphate-creatine phosphotransferase from one ox brain*

The weight of the brain was 453 g. The purification procedure was as described in the text. Fractions precipitated by acetone or $(\text{NH}_4)_2\text{SO}_4$ were redissolved in 50 mM-ammonium citrate, pH 9.0. Experimental details are given in the text.

	Volume (ml.)	Total enzyme (units)	Total protein (mg.)	Specific activity (units/mg. of protein)	Yield (%)
10 mM-KCl-1 mM-EDTA extract	765	5730	4050	1.42	100
70 mM-Magnesium acetate extract	70	4620	455	10.1	80
Solution of crystalline deposit	24	226	17	13.4	4
0-30% (v/v) Acetone fraction	22	123	74	1.7	2
30-40% (v/v) Acetone fraction	32	3160	177	19.7	55
> 40% (v/v) Acetone fraction	112	336	117	2.9	6
0-1.75M- $(\text{NH}_4)_2\text{SO}_4$ fraction	7	39	18	2.2	1
1.75-2.4M- $(\text{NH}_4)_2\text{SO}_4$ fraction	7	3220	119	27.0	56
> 2.4M- $(\text{NH}_4)_2\text{SO}_4$ fraction	36	88	112	0.8	2
Enzyme from column	—	—	—	49-73	~56

Fig. 2. Electrophoresis patterns on cellulose acetate of fractions obtained during the purification of adenosine triphosphate-creatine phosphotransferase (barbiturate buffer, pH 8.6 and I 0.05). (A), Fractions obtained by salt and acetone precipitation procedure: (a) initial KCl extract; (b) 70 mM-magnesium acetate extract; (c) 0-30% (v/v) acetone fraction; (d) 30-40% (v/v) acetone fraction; (e) 40-50% (v/v) acetone fraction; (f) 0-1.75M- $(\text{NH}_4)_2\text{SO}_4$ fraction; (g) 1.75-2.4M- $(\text{NH}_4)_2\text{SO}_4$ fraction; (h) 2.4M- $(\text{NH}_4)_2\text{SO}_4$ fraction. O-O', Origin; Hb, position of haemoglobin; E, position of a sample of the purified enzyme. (B), Fractions obtained by elution of the 1.75-2.4M- $(\text{NH}_4)_2\text{SO}_4$ fraction from a DEAE-Sephadex column by acetate at pH 5.5 (tube nos. refer to those given in Fig. 3a): (a) tubes 2+3; (b) tubes 8-11; (c) tube 30; (d) tubes 40+41, 42+43, 46+47, 48+49, all gave identically this pattern; (e) tubes 52-54. O-O', Origin; E, position of adenosine triphosphate-creatine phosphotransferase; P, position of D-2-phosphoglycerate-hydro-lyase activity. ■, Heavy staining; ▨, medium staining; ▤, light staining.

completing the run, compared with a time of 5-12 hr. required for starch gels. Cellulose acetate had the disadvantage that some bands migrated toward the cathode and there were small variations in pattern from one run to the next. These variations could be limited, however, by ensuring that precisely the same procedure was followed for each run when soaking the strips in buffer, applying the sample, and positioning the strips and supports in the tank. The presence of large amounts of inorganic salts interfered with the separation on cellulose acetate and it was necessary to dialyse some samples before applying them to the strip.

The initial potassium chloride extracts of brain gave patterns of rather poorly defined bands with hazy edges and a considerable amount of background, both on cellulose acetate and in starch gel, whereas more purified fractions, although still containing a large number of components, gave sharp bands with much less background colour. A similar observation was made by Robertson (1957), who attributed the poor resolution of the crude extracts to interference by proteolipids. The interpretation of the patterns obtained on cellulose acetate (Fig. 2) proved difficult, which was not unexpected, since each band visible on the electrophoretic pattern probably represents a mixture of several proteins of similar charge and size, each of which on treatment with a particular reagent may behave in a dissimilar way to other proteins in the same band and similarly to some proteins present in neighbouring bands. However, it did appear that the slower-moving proteins were principally removed by 30% (v/v) acetone or 1.75M-ammonium sulphate, and a number of faster-moving proteins, in a position similar to that of serum albumin, remained unprecipitated by 40% (v/v) acetone or 2.4M-ammonium sulphate. The enrichment of certain fractions with the enzyme was paralleled by an increase in the relative intensity of one of the 'albumin-like' fast-moving bands

(marked E in Fig. 2). The pure enzyme ran to the same position as this band both on cellulose acetate and in starch gel.

Column chromatography. Electrophoresis on cellulose acetate of fractions from the DEAE-Sephadex column showed that proteins generally tended to be eluted in the order of increasing electrophoretic mobility, as reported by Peterson & Sober (1956). In every preparation so far submitted to fractionation on the column, enzyme activity was located in two major peaks and a minor peak (I, II and III in Fig. 3a). The recovery of the enzyme from the column was almost 100%. After concentration by ultrafiltration-dialysis the material from peak I and from peak II ran as a single band, of the same mobility, both on cellulose acetate and in starch gel, this mobility being con-

siderably higher than that of a sample of the purified rabbit-muscle enzyme (kindly provided by Dr D. C. Watts).

Samples of enzyme from the summits of peaks I and II were taken, and Michaelis constants, maximum velocities, sedimentation coefficients and activities were determined. These quantities are summarized in Table 4, together with the corresponding values for the rabbit-muscle enzyme (Noda, Kuby & Lardy, 1954; Kuby *et al.* 1954*b*). The materials from peaks I and II were devoid of D-2-phosphoglycerate-hydro-lyase activity but considerable activity was found in that from peak III, which, on cellulose acetate electrophoresis, gave a major band in the same position as components I and II and two weaker bands, one of which had D-2-phosphoglycerate-hydro-lyase activity (Fig. 2).

Fig. 3. Gradient elution of adenosine triphosphate-creatine-phosphotransferase fractions on DEAE-Sephadex by sodium acetate-1 mM-EDTA, pH 5.5. (a) The 1.75-2.4M-(NH₄)₂SO₄ fraction; (b) material from peak I of Fig. 3 (a), rechromatographed after concentration by ultrafiltration-dialysis; (c) material from peak II of Fig. 3 (a), rechromatographed after concentration by ultrafiltration-dialysis. —, $E_{280\text{ m}\mu}$; - - -, enzyme activity;, acetate gradient.

Table 4. Comparison of the properties of components I and II of the cerebral adenosine triphosphate-creatine phosphotransferase from the diethylaminoethyl-Sephadex column with those of the pure muscle enzyme

Protein was measured by the biuret method. Sedimentation coefficients were determined in a Spinco analytical ultracentrifuge. K_m and V_{max} were calculated from a Lineweaver-Burk plot of initial velocities measured at 30° in incubation mixtures containing creatine (24 mM), sodium thioglycollate (1 mM), glycine, pH 9.0 (100 mM), and variable but equal concentrations of ATP and magnesium acetate.

	Component I	Component II	Muscle enzyme*
Specific activity (units/mg. of protein)	73	49	52-72
K_m (variable substrate, ATP) (mM)	0.54	0.57	0.50
V_{max} (μ mole/min./ μ g. of protein)	0.059	0.064	0.18
K_m/V_{max} (arbitrary unit)	0.023	0.021	—
Sedimentation coefficient, $S_{20,w}$ (s)†	4.73	4.77	5.0
Concentration by biuret (mg./ml.)			
Extinction at 280 m μ in 1 cm. cell	1.13	1.17	1.14

* Data of Kuby, Noda & Lardy (1954*a, b*), Noda, Kuby & Lardy (1954) and Mahowald, Noltmann & Kuby (1962).

† Measured at concentrations of 0.5 mg. of brain enzyme/ml., and 8.0 mg. of muscle enzyme/ml.

Rechromatography of the material from peaks I and II yielded two very similar patterns (Figs. 3b and 3c), which appeared to indicate interconversion of these two forms. In addition, a peak had appeared of material elutable with 0.10 M-acetate and absorbing light at 280 m μ . This material did not have an ultraviolet-absorption curve typical of a protein ($E_{280}/E_{260} = 0.89$) and appeared to arise from the dialysis sac, since material with a similar absorption curve was obtained when 20 ml. of 0.40 M-acetate, pH 5.5, was concentrated by ultrafiltration-dialysis. The liquid from inside the sac, placed on the column and eluted in the same manner as was the enzyme, gave rise to an ultraviolet-absorbing peak in the same position as that observed in Figs. 3 (b) and 3 (c).

DISCUSSION

Content of the enzyme in the brain. About 80–90 % of the total enzyme activity was removed from the homogenized tissue by the first extraction, in agreement with the results of Narayanaswami (1952). After a second extraction the residue was almost free of activity and it was concluded that, as in muscle, adenosine triphosphate-creatine phosphotransferase is located in the cytoplasm, or, if present in particle-bound form, is easily removable under these conditions. Considerable imbibition of liquid by the cell debris occurred, presumably owing to swelling and disruption of sub-cellular components in the hypo-osmotic medium, and this complicated calculations of enzyme content from the results obtained from a number of enzyme preparations. More reliable results could have been obtained by dispersing the brain in a larger volume of extractant but the resulting solution would then have been unsuitable for preparation of the enzyme. The value of 24 ± 11 units/g. of brain can be compared with the value of 207 units/g. of tissue quoted for rabbit skeletal muscle (Kuby *et al.* 1954a).

The amount of enzyme in the potassium chloride extract of brain was one-eighth of that in the extract of muscle described by Kuby *et al.* (1954a), whereas the amount of protein was approximately one-half, making the specific activity of the enzyme in the brain extract about one-quarter of that in the corresponding muscle extract. Assuming that assays of the enzyme in the pure state and in the crude extract may be compared, Czok & Bücher (1960) have calculated from the data of Kuby *et al.* (1954a) that the muscle enzyme constitutes 5.7 % of the total protein in the extract. If the specific activity of the pure brain enzyme is taken as 72 units/mg. the corresponding value for the brain extract is about 2 %.

Purification and properties. Enzymic activity

was recovered from the DEAE-Sephadex column in two major peaks and one minor peak. The enzyme from either peak I or II sedimented as a single component in the ultracentrifuge and migrated as a single band on cellulose acetate and in starch gel. In these media both components had the same mobility, which was approximately five times that of a sample of the muscle enzyme, indicating that the molecule of the brain enzyme had a more negative charge at pH 8.6 than the muscle enzyme. The specific activities of component I (72 units/mg.) and component II (49 units/mg.) were of the same magnitude as the values reported by Kuby and his co-workers for the muscle enzyme [52 units/mg. when the enzyme was first crystallized by Kuby *et al.* (1954a), and 75 units/mg. reported by Mahowald, Noltmann & Kuby (1962) for their latest preparation]. The only differences so far observed between components I and II were in the specific activities and, in one preparation, that II gave a broad diffuse band on cellulose acetate whereas I gave a narrow sharply defined band. However, both components from a second preparation gave sharp bands and were indistinguishable from each other.

Recovery of the enzyme in three peaks would appear to indicate the existence of three 'isoenzymes' (Markert & Møller, 1959). However, cellulose acetate electrophoresis of material from peak III revealed the presence of four protein bands and, since considerable D-2-phosphoglycerate-hydro-lyase activity was also detected and was localized in one of the bands, it was concluded that the peak was due to association of adenosine triphosphate-creatine phosphotransferase with this enzyme and other foreign proteins. Since rechromatography of components I and II yielded two very similar patterns (Figs. 3b and 3c) with a number of peaks, it was concluded that the two species of the enzyme were interconvertible. From this evidence and the results of kinetic measurements and studies in the ultracentrifuge, it was concluded that I and II represented structurally different forms of the same enzyme. Thus Hartley, Peterson & Sober (1962) isolated on diethylaminoethylcellulose a number of isomers of crystalline bovine serum albumin differing in the number and degree of oxidation of their thiol groups and were able to produce such isomers by treatment of the solid protein with ethanol, and Rosenberg, Rogers, Haebig & Steck (1962) have shown that, when aqueous solutions of albumin are treated with ethanol, hydrophobic bonds may be broken and the molecule expanded. Since the enzyme being studied has similar solubility properties, a similar mobility on electrophoresis, and most probably a molecular weight similar to that of serum albumin, it seems likely that the low pH, exposure to air and

exposure to the conditions on the column might have been sufficient to bring about mild structural changes in the 'albumin-like' molecule of the enzyme without loss of its catalytic activity. That the brain enzyme has a labile tertiary structure is suggested by its ready inactivation by heat, alcohols and on storage, and by the spontaneous activation of some samples that has been observed.

Activation of the muscle enzyme by thiols was reported by Lehmann & Pollak (1942) and by Oliver (1955) in crude extracts, and in purified preparations by Cho, Haslett & Jenden (1960), who found that the addition of 1 mM-thioglycollate prevented the reversible inactivation of the highly diluted enzyme during titrimetric assay. In their experiments, Ennor & Rosenberg (1954) were able to observe an effect only at pH 7, whereas Kuby *et al.* (1954*b*) did not observe any effect of cysteine on the reaction velocity or on the heat stability of their purified enzyme. However, Mahowald *et al.* (1962) have confirmed the findings of Cho *et al.* (1960). In the present investigation, up to 800 % activation of the brain enzyme by 1 mM-thioglycollate was consistently observed at various stages of purification. Since EDTA at the same concentration gave only slight activation, it seems unlikely that the effect could be due to the removal of inhibitory metal impurities.

These observations of activation by thiols are of considerable interest since it is known that of the six thiol groups present in the muscle-enzyme molecule, two are located at the active site and participate in the formation of the enzyme-substrate complex (Rabin & Watts, 1960; Watts, Rabin & Crook, 1962; Mahowald *et al.* 1962). Hotta (1962) has suggested that the hexose monophosphate pathway in brain plays a major physiological role by maintaining glutathione and the cellular milieu in the reduced state. The above observations suggest that brain adenosine triphosphate-creatine phosphotransferase may be one enzyme where the maintenance of its thiol groups in a reduced state may well be of importance to the efficient functioning of the enzyme *in vivo*.

SUMMARY

1. Adenosine triphosphate-creatine phosphotransferase was extracted with potassium chloride solution from ox brains; it was present in amounts averaging 24 ± 11 units/g. of tissue and represented approx. 2 % of the protein in the extract.

2. A 50-fold purification was achieved and each step of the procedure was followed by cellulose acetate and starch-gel electrophoresis. The pure enzyme had a specific activity of 73 units/mg. (component I).

3. The enzyme was obtained in three peaks from a diethylaminoethyl-Sephadex column. Two of these peaks appeared to consist of interconvertible forms of the pure enzyme, whereas the third peak contained the enzyme in association with D-2-phosphoglycerate hydro-lyase and other foreign proteins.

4. Marked activation was obtained with thiols, and the enzyme appeared to be less stable to heat, alcohols and storage than the corresponding enzyme from muscle.

The author is indebted to Professor H. McIlwain for helpful discussion and to Dr D. B. Gammack for much helpful advice on the techniques of starch-gel and cellulose acetate electrophoresis. Thanks are also due to Mr H. Russell for able technical assistance and to Mr J. Mills for the assays of D-2-phosphoglycerate hydro-lyase. Runs in the ultracentrifuge were carried out by Dr D. B. Gammack, in an instrument provided by the Medical Research Council.

REFERENCES

- Askonas, B. A. (1951). *Biochem. J.* **48**, 42.
- Bodman, J. (1960). In *Chromatographic and Electrophoretic Techniques*, vol. 2, p. 91. Ed. by Smith, I. London: William Heinemann Ltd.
- Cherkin, A., Martinez, F. E. & Dunn, M. S. (1953). *J. Amer. chem. Soc.* **75**, 1244.
- Cho, A. K., Haslett, W. L. & Jenden, D. J. (1960). *Biochem. J.* **75**, 115.
- Czok, R. & Bücher, T. (1960). *Advanc. Protein Chem.* **15**, 315.
- DaCosta, W. A. & Friedburg, F. (1960). *J. biol. Chem.* **235**, 3134.
- Dixon, M. (1953). *Biochem. J.* **54**, 457.
- Ennor, A. H. & Morrison, J. F. (1960). In *The Enzymes*, vol. 2, p. 108. Ed. by Boyer, P. D., Lardy, H. & Myrback, K. New York: Academic Press Inc.
- Ennor, A. H. & Rosenberg, H. (1954). *Biochem. J.* **57**, 203.
- Everall, P. H. & Wright, G. H. (1958). *J. med. Lab. Tech.* **15**, 209.
- Geiger, W. B. (1956). *J. biol. Chem.* **220**, 871.
- Hartley, R. W., Peterson, E. A. & Sober, H. A. (1962). *Biochemistry*, **1**, 60.
- Hotta, S. S. (1962). *J. Neurochem.* **9**, 43.
- Kohn, J. (1960). In *Chromatographic and Electrophoretic Techniques*, vol. 2, p. 56. Ed. by Smith, I. London: William Heinemann Ltd.
- Kuby, S. A., Noda, L. & Lardy, H. A. (1954*a*). *J. biol. Chem.* **209**, 191.
- Kuby, S. A., Noda, L. & Lardy, H. A. (1954*b*). *J. biol. Chem.* **210**, 65.
- Kuby, S. A., Noda, L. & Lardy, H. A. (1954*c*). *J. biol. Chem.* **210**, 83.
- Lakshmanan, T. K. & Lieberman, S. (1953). *Arch. Biochem. Biophys.* **45**, 235.
- Layne, E. (1957). In *Methods in Enzymology*, vol. 3, p. 450. Ed. by Colowick, S. P. & Kaplan, N. O. New York: Academic Press Inc.

- Lehmann, H. & Pollak, L. (1942). *Biochem. J.* **36**, 672.
- Lineweaver, H. & Burk, D. (1934). *J. Amer. chem. Soc.* **56**, 658.
- Mollwain, H. (1959). *Biochemistry of the Central Nervous System*, 2nd ed., p. 58. London: J. and A. Churchill Ltd.
- Mahowald, T. A., Noltmann, E. A. & Kuby, S. A. (1962). *J. biol. Chem.* **237**, 1535.
- Malmström, B. G. (1957). *Arch. Biochem. Biophys.* **70**, 58.
- Markert, C. L. & Møller, F. (1959). *Proc. nat. Acad. Sci., Wash.*, **45**, 753.
- Narayanaswami, A. (1952). *Biochem. J.* **52**, 295.
- Noda, L., Kuby, S. A. & Lardy, H. A. (1954). *J. biol. Chem.* **209**, 203.
- Noda, L., Kuby, S. A. & Lardy, H. (1955). In *Methods in Enzymology*, vol. 2, p. 605. Ed. by Colowick, S. P. & Kaplan, N. O. New York: Academic Press Inc.
- Oliver, I. T. (1955). *Biochem. J.* **61**, 116.
- Peterson, E. A. & Sober, H. A. (1956). *J. Amer. chem. Soc.* **78**, 756.
- Poulik, M. D. (1957). *Nature, Lond.*, **180**, 1477.
- Rabin, B. R. & Watts, D. C. (1960). *Nature, Lond.*, **188**, 1163.
- Robertson, D. M. (1957). *J. Neurochem.* **1**, 358.
- Rosenberg, R. M., Rogers, D. W., Haebig, J. E. & Steck, T. L. (1962). *Arch. Biochem. Biophys.* **97**, 433.
- Watts, D. C., Rabin, B. R. & Crook, E. M. (1962). *Biochem. J.* **82**, 412.
- Wood, T. (1962). *Biochem. J.* **85**, 31r.

Biochem. J. (1963) **87**, 462

Control of Porphyrin Biosynthesis through a Negative-Feedback Mechanism

STUDIES WITH PREPARATIONS OF δ -AMINOLAEVULATE SYNTHETASE AND δ -AMINOLAEVULATE DEHYDRATASE FROM *RHODOPSEUDOMONAS SPHEROIDES*

BY B. F. BURNHAM* AND JUNE LASCELLES

Microbiology Unit, Department of Biochemistry, University of Oxford

(Received 5 November 1962)

The photosynthetic bacterium *Rhodospseudomonas spheroides*, like many other micro-organisms, accumulates porphyrins in the medium when cultured under iron-deficient conditions (Lascelles, 1961). The addition of small amounts of iron ($2 \mu\text{m-moles/ml.}$) to suspensions of iron-deficient *R. spheroides* causes a large decrease (up to $100 \mu\text{m-moles/ml.}$) in the quantity of porphyrin produced by the organism. Iron cannot therefore be acting solely by diverting porphyrins (or precursors) to iron porphyrins since the quantity of porphyrin that fails to accumulate in the presence of iron is about 100 times more than the amount of iron required to suppress the excretion (Lascelles, 1956).

Iron may act catalytically on porphyrin excretion by promoting the synthesis of a haem compound that in turn controls the activity of an enzyme concerned in porphyrin synthesis through a negative-feedback mechanism. In other negative-feedback systems the final product in a metabolic sequence inhibits the first enzymic step that leads directly to that end product (Umbarger, 1961). In porphyrin synthesis the enzyme that fills this position is δ -aminolaevulate synthetase, which catalyses the formation of δ -aminolaevulate from glycine and succinyl-CoA (Kikuchi, Kumar, Talmage & Shemin, 1958; Gibson, 1958).

* Present address: C.F. Kettering Research Laboratory, Yellow Springs, Ohio, U.S.A.

Previous work has shown that haemin represses the formation of δ -aminolaevulate synthetase in *R. spheroides* (Lascelles, 1960) and that haemin inhibits the action of the enzyme in crude cell extracts (Gibson, Matthew, Neuberger & Tait, 1961). In the present investigation the possibility that haemin exerts a negative-feedback type of control on δ -aminolaevulate synthetase has been examined by studying its action on partially purified preparations of the enzyme from *R. spheroides*. Evidence consistent with such a controlling function has also been shown with suspensions of the organism that produce porphyrins. Preliminary accounts of this work have appeared (Burnham, 1962a, b). The purification of succinyl-coenzyme A thiokinase (used in the assay for δ -aminolaevulate synthetase) and of δ -aminolaevulate dehydratase from *R. spheroides* is also described.

EXPERIMENTAL

Chemicals

CoA (75%), ATP, pyridoxal phosphate, horse-heart cytochrome c and twice-recrystallized bovine haemoglobin were obtained from the Sigma Chemical Co., St Louis, Mo., U.S.A.

Succinyl-CoA was prepared by the method of Simon & Shemin (1953) and was used immediately. The values quoted for succinyl-CoA are based on the amount of CoA