6. Stimulation markedly decreased the specific radioactivity of the aspartate formed in the presence of the ^{14}C -labelled glutamic acid.

7. The approximate rate of conversion of glutamic acid into γ -aminobutyric acid in rat-brain-cortex slices was estimated. It was concluded that the increase of γ -aminobutyric acid in the presence of ethanol *in vivo* and in stimulated slices is due to an inhibition of its breakdown. This effect is not direct, since it was not observed in non-stimulated slices.

We acknowledge our gratitude to our Chief, Dr H. Suomalainen, for encouragement and criticism, to Dr O. Forsander and other colleagues for discussions and valuable comments, and to Mrs Kaija Salmela and Miss Gunnell Sievers for skilful technical assistance.

REFERENCES

- Baxter, C. F. & Roberts, E. (1960). In *Inhibition in the Nervous System and γ -Aminobutyric Acid*, p. 358. Ed. by Roberts, E. London: Pergamon Press Ltd.
- Calvin, M., Heidelberger, C., Reid, J. C., Tolbert, B. M. & Yankwich, P. E. (1949). *Isotopic Carbon*, p. 318. New York: John Wiley and Sons Inc.
- Cohen, M. M., Simon, G. R., Berry, J. F. & Chain, E. B. (1962). *Biochem. J.* **84**, 43p.
- Davison, A. N. (1961). *Biochem. J.* **78**, 272.
- Eidelberg, E., Baxter, C. F., Roberts, E. & Saldias, C. A. (1960). In *Inhibition in the Nervous System and γ -Aminobutyric Acid*, p. 365. Ed. by Roberts, E. London: Pergamon Press Ltd.
- Elliott, K. A. C. & Florey, E. (1957). *J. Neurochem.* **1**, 181.
- Häkkinen, H.-M. & Kulonen, E. (1961). *Biochem. J.* **78**, 588.
- Killam, K. F. & Bain, J. S. (1957). *J. Pharmacol.* **119**, 255.
- Killam, K. F., Dasgupta, S. R. & Killam, E. K. (1960). In *Inhibition in the Nervous System and γ -Aminobutyric Acid*, p. 302. Ed. by Roberts, E. London: Pergamon Press Ltd.
- Kini, M. M. & Quastel, J. H. (1959). *Nature, Lond.*, **184**, 252.
- Kulonen, E., Häkkinen, H.-M. & Wallgren, H. (1960). *Proc. 4th int. Congr. clin. Chem.* p. 135. Edinburgh and London: E. and S. Livingstone Ltd.
- Lovern, J. A. (1955). *The Chemistry of Lipids of Biochemical Significance*, pp. 37–40. London: Methuen and Co. Ltd.
- McIlwain, H. (1951). *Biochem. J.* **49**, 382.
- McIlwain, H. (1959). In *Biochemistry of the Central Nervous System*, p. 46. Ed. by Brücke, F. London: Pergamon Press Ltd.
- McKhann, G. M., Albers, R. W., Sokoloff, L., Mickelsen, O. & Tower, D. B. (1960). In *Inhibition in the Nervous System and γ -Aminobutyric Acid*, p. 169. Ed. by Roberts, E. London: Pergamon Press Ltd.
- McKhann, G. M. & Tower, D. B. (1959). *Amer. J. Physiol.* **196**, 36.
- Purpura, D. P., Berl, S., Gonzalez-Monteagudo, O. & Wyatt A. (1960). In *Inhibition in the Nervous System and γ -Aminobutyric Acid*, p. 331. Ed. by Roberts, E. London: Pergamon Press Ltd.
- Quastel, J. H. (1959). In *Biochemistry of the Central Nervous System*, p. 90. Ed. by Brücke, F. London: Pergamon Press Ltd.
- Richter, D. & Dawson, R. M. C. (1948). *J. biol. Chem.* **176**, 1199.
- Roberts, E. (ed.) (1960). *Inhibition in the Nervous System and γ -Aminobutyric Acid*. London: Pergamon Press Ltd.
- Sammalisto, L. (1962). *Nature, Lond.*, **195**, 185.
- Turský, T. (1960). *Nature, Lond.*, **187**, 322.
- Wallgren, H. (1960). *Acta physiol. scand.* **49**, 216.
- Wallgren, H. (1961). *Biochem. Pharmacol.* **6**, 195.
- Wallgren, H. & Kulonen, E. (1960). *Biochem. J.* **75**, 150.
- Zilversmit, D. B., Entenman, C. & Fishler, M. C. (1943). *J. gen. Physiol.* **26**, 325.

Biochem. J. (1963) **88**, 498

Purine and Pyrimidine Derivatives in Mature Pea Seeds

By E. G. BROWN

Department of Botany, University College, Swansea, S. Wales

(Received 5 April 1963)

In a recent investigation, the acid-soluble nucleotides of mature pea seeds were described and determined (Brown, 1962). However, except for a few reports describing the isolation of individual compounds (e.g. Charbuliez & Bernhard, 1932; Ritthausen, 1876, 1881; Fosse, deGraeve & Thomas, 1932), little is known of other purine and pyrimidine derivatives in seeds. To obtain a more complete picture, mature pea seeds have been examined with the main aim of identifying and determining the relative concentrations of such compounds. The analytical technique used was

based on ion-exchange chromatography; anion-exchange rather than cation-exchange chromatography was employed in order to take advantage of the absence of glycosidic lability under alkaline conditions (Cohn, 1955).

MATERIALS AND METHODS

Preparation of seed extract. Mature seeds of *Pisum sativum* L. var. Meteor obtained from Suttons Seeds Ltd., Reading, and showing 88% germination, were used. Purines, pyrimidines, and their derivatives, were extracted

from 10 g. samples of finely milled seed with ice-cold 0.3N-perchloric acid as described by Brown (1962). After adjustment of the pH to 7.2 with 20% (w/v) KOH, the extract was kept at 0° for 30 min. and then centrifuged to remove potassium perchlorate. The clear pale-yellow supernatant was passed slowly through a column (30 mm. $\times$ 10 mm.) of Dowex 1 (formate form; X 8; 200–400 mesh), and the material passing straight through the column (fraction A) was collected and stored at -15° . The first 30 ml. fraction on elution with 0.03M-formic acid was evaporated to dryness *in vacuo* in a rotary-film evaporator and then dissolved in fraction A. This procedure removed the bulk of interfering pigments and nucleotides, at the same time permitting complete recovery of the purine and pyrimidine bases and their ribosides.

Anion-exchange chromatography. After the above treatment, the extract was adjusted to pH 10.8 with aq. ammonia (sp.gr. 0.88) and percolated through a column (19 mm. $\times$ 0.9 mm.) of Dowex 1 (Cl^{-} form; X 10; 200–400 mesh) at a rate of 0.4 ml./min. After the column had been washed with 10 ml. of water adjusted to pH 10.8 with aq. ammonia, elution by charge adjustment was begun by using ammonia–ammonium chloride buffers (Cohn, 1949). Fractions (10 ml.) were collected at a rate of 0.5 ml./min. After elution with the two buffers described by Cohn, a third solution containing 0.2M-ammonium chloride adjusted to pH 9.4 with aq. ammonia was used to elute uric acid. The column was calibrated with the more commonly occurring purine and pyrimidine bases and ribosides with respect to both their position in the elution sequence (cf. Cohn, 1949, 1955) and their quantitative recovery. Uric acid, in contrast with all the other substances examined, was not quantitatively recovered (about 2% recovery). This was attributed to its rapid oxidation in alkaline solution (Griffiths, 1952). Quantitative recovery was achieved when the pH was not allowed to exceed 8.3. Uric acid was therefore isolated in separate experiments by adsorption of the seed extract, prepared as described above, on the column at pH 8.3, followed by elution with 0.06M-ammonium chloride solution adjusted to pH 7.8 with aq. ammonia. The separation of uric acid from a known mixture of purine and pyrimidine derivatives by this method is shown in Fig. 2 (a).

Examination of fractions. Fractions comprising each of the main peaks were pooled and evaporated to dryness *in vacuo* in a rotary-film evaporator. The dry residues were dissolved in water and used directly for preliminary spectrophotometric examination in a Unicam SP. 500 instrument. Before further resolution by paper chromatography and more detailed examination of the materials, they were desalted by adsorption on a charcoal column as described by Brown (1962) and subsequently eluted with 20 ml. of aq. pyridine (10%, v/v). The eluate was evaporated *in vacuo* and the dry residue dissolved in about 1 ml. of water for paper chromatography on Whatman no. 1 paper with one or more of the following solvent systems: (1) butan-1-ol–acetic acid–water (4:1:5, by vol.) (Partridge, 1948); (2) water at pH 10 (Levenbook, 1955); (3) aq. 5% (w/v) disodium hydrogen phosphate saturated with 3-methylbutan-1-ol (Carter, 1950); (4) M-ammonium acetate (containing sodium borate and EDTA)–ethanol (3:7, v/v) (Plesner, 1955); (5) phenol saturated with water (Fink & Fink, 1949); (6) butanol saturated with water (Stadtman & Barker, 1950); (7) acetate buffer (pH 4.5)–ethanol (1:1, v/v) (Stadtman, 1957); (8) pyridine–propan-2-ol–water (1:1:1,

by vol.) (Stadtman, 1957). All chromatograms were ascending except those run in solvent 3, which were descending. Substances on the chromatograms were located by viewing in ultraviolet light from a Hanovia Chromatolite lamp. For very weakly absorbing spots, the more sensitive method of contact printing on to photographic paper in ultraviolet light (Markham & Smith, 1949) was adopted. Suitable areas were eluted for further examination by the capillary-elution technique of Dent (1947).

Phosphate and pentose were detected and determined as described by Brown (1962). Adenine and its derivatives were detected by the spot test of Gerlach & Döring (1955).

Preparation and determination of hydroxamates. The method of Stadtman & Barker (1950) was employed to prepare acylhydroxamic acids from seed fractions. Aceto-hydroxamic acid, succinohydroxamic acid and malono-hydroxamic acid were prepared from the corresponding ethyl esters by the method of Thompson (1951). Hydroxamates were determined colorimetrically after the addition of ferric chloride solution (Lipmann & Tuttle, 1945).

Preparation of 3'-nucleotidase. Seeds of Dunns Perennial Rye Grass (*Lolium perenne*), supplied by Dunns Farm Seeds Ltd., Salisbury, were soaked in water at 27° in the dark for 30 hr. and then extracted with a total of 2 l. of deionized water in a Waring Blender. The extract was cleared by centrifuging and then fractionated by ammonium sulphate precipitation. The fraction precipitating at 70–90% saturation was collected by centrifuging, redissolved in 5 ml. of deionized water and dialysed overnight against deionized water at 4°. This preparation was used directly without further purification.

Preparation of acetyl-3'-dephosphocoenzyme A. Coenzyme A (5 mg.) was acetylated at pH 7.5 with acetic anhydride by appropriate modification of the method described by Simon & Shemin (1953) for the preparation of succinyl-CoA. The resulting acetyl-CoA was separated as a band by paper chromatography in solvent 7 and, after elution with deionized water (2 ml.), incubated for 30 min. at 37° with 1 ml. of 3'-nucleotidase at pH 7.4. The reaction was stopped by the addition of 60% (v/v) perchloric acid (0.13 ml.) and the mixture centrifuged to remove precipitated protein. The supernatant was adjusted to pH 7.2 with potassium hydroxide, allowed to stand at 0° for 60 min. and then centrifuged to remove potassium perchlorate. The solution was desalted by adsorption on charcoal and subsequent elution with aq. 10% (v/v) pyridine, the latter being removed immediately after elution by evaporation *in vacuo*. The product was subjected to electrophoresis on paper, with a formate buffer at pH 3.6 and a voltage gradient of approx. 20 v/cm. After 2 hr. the paper was examined in ultraviolet light, and the main band (7.3 cm. migration) was eluted with deionized water and freeze-dried. Two, much fainter, bands (most probably unchanged acetyl-CoA and free CoA) were also observed at 15.4 and 21.7 cm.

RESULTS

Separation and identification of purine and pyrimidine derivatives

A typical anion-exchange chromatogram, obtained from 10 g. samples of mature pea seeds, is shown in Fig. 1. The major peaks in the chromatogram were examined, and where possible identified,

as indicated below, with the help of the known elution sequence of the more common purine and pyrimidine bases and nucleosides.

Peak 1. The ultraviolet-absorption spectra of the material composing this peak and its extinction ratios $E_{250\text{ m}\mu}:E_{260\text{ m}\mu}$ and $E_{280\text{ m}\mu}:E_{260\text{ m}\mu}$ of 0.84 and 0.22 respectively at pH 2, and 0.78 and 0.15 respectively at pH 12, indicated adenosine. The material gave a positive reaction in the spot test of Gerlach & Döring (1955) for an adenine derivative, a positive orcinol test for pentose and a negative test for phosphate. Paper-chromatographic comparison with authentic samples of adenosine, in solvents 1 and 2, confirmed identification of this material as adenosine, and showed that it was the only ultraviolet-light-absorbing compound present in peak 1.

Peak 2. The material of peak 2 gave a positive test for pentose but not for phosphate. Its extinction ratios $E_{250\text{ m}\mu}:E_{260\text{ m}\mu}$ and $E_{280\text{ m}\mu}:E_{260\text{ m}\mu}$ were 0.74 and 0.35 respectively at pH 12. These observations were consistent with the properties of uridine. Detailed ultraviolet-absorption spectra, and comparison with authentic samples of uridine by paper chromatography in solvents 1 and 2, confirmed this identification and, as with the preceding peak, showed that only one ultraviolet-light-absorbing compound was present.

Peak 3. Peaks from three separations were combined, fractions being stored at -15° during this accumulation of material. The material gave extinction ratios $E_{250\text{ m}\mu}:E_{260\text{ m}\mu}$ and $E_{280\text{ m}\mu}:E_{260\text{ m}\mu}$ of 0.95 and 0.70 respectively at pH 2, and 0.90 and 0.61 respectively at pH 12. These values, the

elution position and the ultraviolet-absorption maxima and minima supported identification as guanosine. Confirmation was obtained by paper chromatography in solvents 1 and 2, in both of which the material corresponded to an authentic sample of guanosine.

Peak 4. Spectrophotometric examination indicated that this peak was not homogeneous. After treatment on charcoal (see the Materials and Methods section) to desalt the material, it was subjected to paper electrophoresis for 2 hr. in a formate buffer at pH 3.6 under a potential gradient of 20 v/cm. This separated the material into two ultraviolet-light-absorbing zones that migrated 1.6 cm. and 7.2 cm. respectively. The slower-running band was eluted from the paper and a sample chromatographed on paper in solvent 1. This produced a single spot (R_f 0.31) which was eluted in 0.01 N-hydrochloric acid and examined in the spectrophotometer. The material exhibited extinction ratios $E_{250\text{ m}\mu}:E_{260\text{ m}\mu}$ and $E_{280\text{ m}\mu}:E_{260\text{ m}\mu}$ of 0.75 and 0.28 respectively at pH 2, and 1.30 and 1.14 respectively at pH 12. These ratios and the ultraviolet-absorption maxima and minima, at the same pH values, were typical of xanthosine or a xanthosine derivative. The substance gave a positive orcinol test for pentose but no phosphate could be detected. Hydrolysis in N-hydrochloric acid for 1 hr. at 100° yielded xanthine which was identified both by its characteristic absorption spectra at pH 2 and 12, and its behaviour on paper chromatography in solvents 1 and 2. Co-chromatography on paper with authentic samples in solvents 1 and 2 showed the material to correspond to xanthosine.

The material migrating 7.2 cm. on paper electrophoresis was also eluted in 0.01 N-hydrochloric acid and chromatographed as a band on paper in solvent 1. This produced a single ultraviolet-light-absorbing zone (R_f 0.16) which was eluted in 0.01 N-hydrochloric acid and had ultraviolet-absorption ratios $E_{250\text{ m}\mu}:E_{260\text{ m}\mu}$ and $E_{280\text{ m}\mu}:E_{260\text{ m}\mu}$ of 0.84 and 0.23 respectively at pH 2, and 0.81 and 0.15 respectively at pH 11. These ratios, together with the ultraviolet-absorption maxima and minima (λ_{max} . 259 m μ and λ_{min} . 235 m μ at pH 2; λ_{max} . 260 m μ and λ_{min} . 231 m μ at pH 11) are typical of an adenosine derivative. The presence of bound adenine was confirmed by the spot test of Gerlach & Döring (1955) and by the liberation on hydrolysis in 2 N-hydrochloric acid for 1 hr. at 100° of free adenine, identified by its characteristic ultraviolet-absorption spectra at pH 2 and 12 and by paper chromatography in solvents 1 and 2. The material was also found to contain phosphate and pentose (see the Materials and Methods section) and gave a delayed nitroprusside reaction (Stadtman, 1957), indicating the presence of a thiol ester. It reacted with hydroxylamine at neutral pH to yield a hydroxamic

Fig. 1. Anion-exchange chromatogram of the purine and pyrimidine derivatives in mature pea seeds. Separation of the extract, equivalent to 10 g. of seeds, was on Dowex 1 (Cl^- form). Experimental details are given in the text. The number of 10 ml. fractions collected and the point at which the composition of the eluting solvent was changed are indicated on the abscissa.

acid detected by the deep-red colour it produced in the presence of ferric chloride. This hydroxamate was identified as acetohydroxamate by co-chromatography on paper with an authentic sample in solvents 1, 5 and 6. All the foregoing observations suggested that the original material was related to acetyl-CoA. On electrophoresis at pH 3.6 the material moved 7.2 cm. towards the anode, whereas acetyl-CoA moved 20 cm. The peak 4 material has, therefore, a smaller net negative charge on the molecule than has acetyl-CoA. Since analysis for the adenine:total phosphate ratio gave 1.0:2.04, the material was tentatively identified as the acetyl derivative of 3'-dephosphocoenzyme A, a sample of which was prepared, as described in the Materials and Methods section, and compared by paper chromatography, by thin-layer chromatography on cellulose (Stahl, 1961) and by paper electrophoresis with the peak 4 material. Solvents 1, 7 and 8 were employed for the paper chromatography and solvents 1 and 7 for the thin-layer chromatography; paper electrophoresis was at pH 3.6 in a formate buffer with a voltage gradient of 20 v/cm. for 2 hr. In all cases the materials behaved similarly and, when applied as a mixed sample, were not resolved from one another by the above procedures. It was therefore concluded that the peak 4 material was acetyl 3'-dephosphocoenzyme A.

Peak 5. Ultraviolet-absorption spectra of this peak, as for the preceding peak, indicated complex composition. After being desalted by treatment with charcoal, it was chromatographed on paper as a band in solvent 1 in which it was resolved into seven zones, five of which were trace amounts of materials, fluorescent in ultraviolet light, having R_f values: 0.00 (white fluorescence), 0.43 (pale-blue fluorescence), 0.47 (blue fluorescence), 0.52 (yellow fluorescence) and 0.81 (pale-blue fluorescence). The main band resolved from peak 5 had R_f 0.14 and in ultraviolet light appeared as a dark zone. It was eluted with 0.01N-hydrochloric acid and had extinction ratios $E_{250\text{ m}\mu}:E_{260\text{ m}\mu}$ and $E_{280\text{ m}\mu}:E_{260\text{ m}\mu}$ of 0.74 and 0.37 respectively at pH 2, and 0.81 and 0.32 at pH 12. These values and the ultraviolet-absorption maxima and minima indicated a uridine derivative. The material gave a positive orcinol test for pentose and a positive test for phosphate. It had the same R_f values in solvents 1 and 3 as authentic uridine 5'-monophosphate. However, uridine 5'-monophosphate would not normally be eluted at this point during ion-exchange chromatography as it is much more firmly held than the nucleosides by virtue of its phosphoric acid grouping. This suggested that the nucleotide was eluted whilst bound through its phosphate group to another unit, and that the free nucleotide isolated was a decomposition product. It seemed likely

that the original compound was of the widely distributed UDP-X type. UDP-glucose and UDP-galactose have been found previously in mature pea seeds (Brown, 1962). Further paper-chromatographic investigation of peak 5 revealed the presence of free sugars (detected by aniline hydrogen phthalate reagent) and inorganic phosphate. As neither sugar nor inorganic phosphate would survive the charcoal-adsorption step of the desalting procedure unless in bound form, it was concluded that they are decomposition products in keeping with the suggested nature of the original uridine nucleotide. Although this substance was the main component of peak 5, it was nevertheless present in too small an amount for more detailed examination.

The second ultraviolet-light-absorbing band (R_f 0.36), after elution and spectrophotometric examination, gave absorption ratios $E_{250\text{ m}\mu}:E_{260\text{ m}\mu}$ and $E_{280\text{ m}\mu}:E_{260\text{ m}\mu}$ of 1.20 and 0.08 respectively at pH 2, and 1.15 and 0.25 respectively at pH 12. The ultraviolet-absorption spectra exhibited fine structure in the region 250–260 m μ typical of a benzenoid nucleus. This material was not identified.

Determination of identified purine and pyrimidine derivatives

Purine and pyrimidine derivatives comprising the peaks of the anion-exchange chromatogram (Fig. 1) were determined spectrophotometrically. For homogeneous peaks, determination was by computation from $E_{265\text{ m}\mu}$ of the constituent fractions; heterogeneous peaks were further resolved by paper chromatography or paper electrophoresis before determination of their constituents. The relative amounts of the various purine and pyrimidine derivatives isolated and identified are shown in Tables 1 and 2. The acetyl 3'-dephosphocoenzyme A of peak 4, besides being determined from spectrophotometric data, was also deter-

Table 1. *Nucleosides of mature pea seeds*

Nucleosides were separated by anion-exchange chromatography as described in Fig. 1 and determined by computation of $E_{265\text{ m}\mu}$ of the fractions comprising the peak. Substances in heterogeneous peaks were determined spectrophotometrically only after they had been resolved by paper chromatography or paper electrophoresis. All values represent the mean of six separations. For comparison, the last column shows the total amounts of the corresponding nucleotides determined by Brown (1962).

Nucleoside	Content in seeds (μ moles/10 g.)	Total corresponding nucleotides in seeds (μ moles/10 g.)
Adenosine	1.52	7.51
Uridine	1.98	1.36
Guanosine	0.67	0.13
Xanthosine	0.91	None detected

Table 2. *Acetyl-3'-dephosphocoenzyme A* content of mature pea seeds

In the spectrophotometric method used, the isolated substance was determined from $E_{285\text{m}\mu}$ at pH 2, the molecular extinction coefficient employed in the calculation being that of CoA at the same pH. The second method involved the formation of acetohydroxamic acid and its subsequent colorimetric assay in the presence of FeCl_3 (Lipmann & Tuttle, 1945). This latter assay was made on unresolved seed extract, the only hydroxamate contributing significantly to the value obtained having been shown to be acetohydroxamate.

Assay method	Content in seeds ($\mu\text{moles}/10\text{ g.}$)
Spectrophotometric determination of the isolated material	1.98
Hydroxamate- FeCl_3 assay of unresolved seed extract	1.89

mined by the formation of acetohydroxamate. The values obtained by the two methods (Table 2) were in good agreement.

Examination for uric acid

Anion-exchange chromatography of 10 g. samples of mature pea seeds, with 0.06 M-ammonium chloride at pH 7.8 as the eluting solvent (see the Materials and Methods section), gave chromatograms as exemplified by Fig. 2 (b). Comparison of this with the corresponding calibration chromatogram (Fig. 2a) suggested that no uric acid was present. The only peak exhibiting significant extinction values at 290 m μ (i.e. in the region of λ_{max} for uric acid at pH 7.8) was that designated D, but, after desalting and concentration, both paper chromatography and the sensitive spectrophotometric method with uricase (Kalckar, 1947) failed to reveal any trace of uric acid in this peak.

Preparation of hydroxamates directly from the initial seed extract

Seeds were extracted with perchloric acid as for ion-exchange chromatography and the extract, after neutralization with potassium hydroxide solution, was examined for hydroxylamine-reacting substances by the technique of Stadtman & Barker (1950). The major product, as expected from the above examination of peak 4, was acetohydroxamate. Traces of two other hydroxamates were found and shown by paper chromatography, in solvents 1, 5 and 6, together with authentic samples, to be succinohydroxamate and malonohydroxamate.

Examination of other seeds

Mature seeds of *Lupinus luteus* L. var. Weiko II and *Phaseolus vulgaris* L. var. Canadian Wonder

Fig. 2. Separation of uric acid by anion-exchange chromatography on Dowex 1 (Cl^- form) with 0.06 M- NH_4Cl at pH 7.8 as the eluting solvent, and collection of 10 ml. fractions. The continuous line indicates $E_{285\text{m}\mu}$ of the fractions and the broken line $E_{290\text{m}\mu}$. (a) Resolution of uric acid from a synthetic mixture containing 0.5–1.0 mg. of each of the following: adenine, guanine, xanthine, hypoxanthine, cytosine, uracil, thymine and uric acid. (b) Chromatogram obtained by the same procedure from an extract of mature pea seeds (10 g. sample).

were examined qualitatively by similar methods to those described and the purine and pyrimidine patterns were similar to that of pea seed.

DISCUSSION

Despite the efficacy of the described ion-exchange methods in separating free purine and pyrimidine bases (see the Materials and Methods section), no compounds of this type were found in the mature seeds of pea, kidney beans or yellow lupin. This is in contrast with the alleged widespread occurrence of these bases in plant tissues (see, e.g., Wehmer & Hadders, 1933; Karrer, 1958), but is in keeping with results obtained in this

Laboratory with extracts of whole seedlings of *Pisum sativum* (E. G. Brown & A. V. Silver, unpublished work). However, as Markham (1955) has pointed out, most reports on the presence of free purine and pyrimidine bases in higher plants were made before the development of modern chromatographic and electrophoretic techniques and in many cases probably describe artifacts arising during drastic isolation procedures from the enzymic or chemical degradation of nucleic acids, nucleotides etc.

The major nucleosides of mature pea seeds, in descending order of quantitative importance, were uridine, adenosine, xanthosine and guanosine. The concentration of free uridine is about 1.5 times that of the total uridine nucleotides (Table 1). The concentration of guanosine is about 5 times that of the total guanosine nucleotides. Adenosine, however, is present in a concentration only one-fifth of that of the total adenosine nucleotides (AMP plus ADP plus ATP). This nucleoside pattern might be expected if the nucleosides are considered to be latent forms of the corresponding nucleotides. For, as discussed by Brown (1962), the nucleotide complement of mature pea seeds is more typical of a system predominantly committed to energy production, whereas the nucleotide complement of seedlings appears to be that of a system mainly concerned with biosynthesis and chemical transformation. Transition from the former state to the latter (presumably during germination) would necessarily involve rapid increase in the concentration of those nucleotides more especially associated with biosynthesis and other anabolic processes. Such nucleotides are those of uridine and guanosine. The adenosine nucleotide concentration, on the other hand, is already high in mature pea seeds and would therefore be expected to require little, if any, supplementation from a nucleoside reserve. The free xanthosine of pea seeds is most probably present by virtue of its metabolic relationship to XMP, a known precursor of GMP (Moyed & Magasanik, 1957).

Although Fosse *et al.* (1932) reported that the seeds of several leguminous plants contain uric acid, some in relatively large amounts (e.g. 23 mg./100 g. of seeds of *Vicia faba*), none was detected in the present work even by the extremely sensitive spectrophotometric uricase assay of Kalckar (1947).

The presence of acetyl-3'-dephospho-CoA in extracts of mature pea seeds raised the question whether or not this was an artifact resulting from the breakdown of acetyl-CoA. 3'-Nucleotidase activity is demonstrable in most damaged plant tissue, but as the peas were homogenized in ice-cold 0.3N-perchloric acid it is unlikely that this enzyme was active during extraction. Further, comparison of the amount of acetyl-3'-dephospho-CoA determined by spectrophotometry after resolution and the amount determined in the initial seed extract by the formation of acetohydroxamate (see Table 2) implies that no acetyl-CoA itself is present, as this substance would be registered by the latter determination but not by the former.

Seifter (1954) reported the CoA content of pea meal to be 440 units/10 g. (i.e. $1.42 \mu\text{moles}/10 \text{ g.}$). As this value is of the same order as that obtained for acetyl-3'-dephospho-CoA in the present investigation (see Table 2), it is possible that Seifter's methods did not distinguish between these two substances. One of these methods, acetylation of sulphanilamide, employed an extract of acetone-dried pigeon-liver powder as an enzyme source. This is, however, also a good source of the enzyme 3'-dephosphocoenzyme A kinase (Wang & Kaplan, 1954), which under the assay conditions would convert 3'-dephospho-CoA into CoA. 3'-Dephospho-CoA is generally said to be inactive in the phosphotransacetylase assay, the second method used by Seifter, but Wang & Kaplan (1954) found this not to be the case in the presence of large amounts of enzyme. It remains to be seen, therefore, whether the CoA previously reported in seeds is in fact acetyl-3'-dephospho-CoA. The probable metabolic origin of the latter is indicated in Scheme 1.

Scheme 1. Probable metabolic origin of acetyl-3'-dephospho-CoA. The reaction marked with an asterisk proceeds only with oxidized CoA (Shuster & Kaplan, 1953).

As with the nucleosides, it is likely that the acetyl-3'-dephospho-CoA of pea seeds is a storage form of the corresponding metabolically active nucleotide, in this instance acetyl-CoA. Thus, as the seed germinates and ATP becomes more readily available, phosphorylation occurs and acetyl-CoA is produced. The possibility should also be considered that acetyl-3'-dephospho-CoA is a reserve of energy in the seed, for acyl thiols are known to have high group-transfer potential and can be used in ATP production. According to Swann (1957), there is evidence that a substance of this type may be the immediate source of energy for cell division. However, as Wang & Kaplan (1954) pointed out, 'the possibility that DPCoA [3'-dephospho-CoA] can function in as yet undescribed systems in which CoA does not react, remains to be investigated'.

SUMMARY

1. Purine and pyrimidine derivatives in mature pea seeds were identified and determined by methods based on anion-exchange chromatography in a chloride system, elution being by charge adjustment.

2. In contrast with some older reports, neither uric acid nor any other free purine or pyrimidine base could be detected in the seeds.

3. The nucleoside constituents of mature pea seeds were, in descending order of quantitative importance, uridine, adenosine, xanthosine and guanosine.

4. An unusual nucleotide isolated from the seeds was identified as acetyl-3'-dephosphocoenzyme A. This was determined both spectrophotometrically and by hydroxamate formation.

5. The crude seed extract showed traces of two other hydroxylamine-reacting substances, one of which gave rise to succinohydroxamate and the other to malonohydroxamate.

6. Mature seeds of *Lupinus luteus* and *Phaseolus vulgaris* exhibited qualitatively similar chromatographic patterns to that of pea seeds.

7. The relationship of the various substances isolated to the metabolic state of dry mature seeds is discussed.

The author acknowledges the able technical assistance of Mrs M. Hughes.

REFERENCES

- Brown, E. G. (1962). *Biochem. J.* **85**, 633.
 Carter, C. E. (1950). *J. Amer. chem. Soc.* **72**, 1466.
 Charbuliez, E. & Bernhard, K. (1932). *Helv. chim. acta*, **15**, 464.
 Cohn, W. E. (1949). *Science*, **109**, 377.
 Cohn, W. E. (1955). In *The Nucleic Acids*, vol. 1, p. 211. Ed. by Chargaff, E. & Davidson, J. N. New York: Academic Press Inc.
 Dent, C. (1947). *Biochem. J.* **41**, 245.
 Fink, K. & Fink, R. M. (1949). *Proc. Soc. exp. Biol., N.Y.*, **70**, 654.
 Fosse, R., deGraeve, P. & Thomas, P. E. (1932). *C.R. Acad. Sci., Paris*, **195**, 1198.
 Gerlach, E. & Döring, H. J. (1955). *Naturwissenschaften*, **42**, 344.
 Griffiths, M. (1952). *J. biol. Chem.* **197**, 399.
 Kalekar, H. M. (1947). *J. biol. Chem.* **167**, 429.
 Karrer, W. (1958). *Konstitution und Vorkommen der organischen Pflanzenstoffe*, p. 1047. Basel: Birkhauser Verlag.
 Levenbook, L. (1955). Cited by G. R. Wyatt in *The Nucleic Acids*, vol. 1, p. 252. Ed. by Chargaff, E. & Davidson, J. N. New York: Academic Press Inc.
 Lipmann, F. & Tuttle, L. C. (1945). *J. biol. Chem.* **159**, 21.
 Markham, R. (1955). In *Modern Methods of Plant Analysis*, vol. 4, p. 246. Ed. by Peach, K. and Tracey, M. V. Berlin: Springer-Verlag.
 Markham, R. & Smith, J. D. (1949). *Biochem. J.* **45**, 294.
 Moyed, H. S. & Magasanik, B. (1957). *J. biol. Chem.* **226**, 351.
 Partridge, S. M. (1948). *Biochem. J.* **42**, 238.
 Plesner, P. (1955). *Acta chem. scand.* **9**, 197.
 Ritthausen, H. (1876). *Ber. dtsh. chem. Ges.* **9**, 301.
 Ritthausen, H. (1881). *J. prakt. Chem.* **24**, 202.
 Seifter, E. (1954). *Plant Physiol.* **29**, 403.
 Shuster, L. & Kaplan, N. O. (1953). *J. biol. Chem.* **201**, 535.
 Simon, E. J. & Shemin, D. (1953). *J. Amer. chem. Soc.* **75**, 2520.
 Stadtman, E. R. (1957). In *Methods in Enzymology*, vol. 3, p. 931. Ed. by Colowick, S. P. and Kaplan, N. O. New York: Academic Press Inc.
 Stadtman, E. R. & Barker, H. A. (1950). *J. biol. Chem.* **184**, 769.
 Stahl, E. (1961). *Angew. Chem.* **73**, 646.
 Swann, M. M. (1957). *Cancer Res.* **17**, 727.
 Thompson, A. R. (1951). *Aust. J. sci. Res. B*, **4**, 180.
 Wang, T. P. & Kaplan, N. O. (1954). *J. biol. chem.* **206**, 311.
 Wehmer, C. & Hadders, M. (1933). In *Handbuch der Pflanzenanalyse*, vol. 4, p. 405. Ed. by Klein, G. Vienna: Springer-Verlag.