two different solvent systems followed by partition chromatography, and by methylation, purification of the methyl ethers and chromatography in two successive systems.

3. The biosynthesis of at least five radioactive compounds in the phenolic fraction was indicated.

4. The lack of evidence for this biosynthesis as a significant biochemical process *in vivo* is emphasized.

The authors are indebted to the following for their gifts of steroids and other biochemicals: Ciba Ltd., Dr R. I. Dorfman, Professor W. Klyne, Paines and Byrne Ltd. and Schering A.G. The skilled technical assistance of Mrs Jean Tilbrook is gratefully acknowledged.

REFERENCES

- Baggett, B., Engel, L. L., Savard, K. & Dorfman, R. I. (1956). *J. biol. Chem.* **221**, 931.
- Barton, G. M., Evans, R. S. & Gardner, J. A. F. (1952). *Nature, Lond.*, **170**, 249.
- Bauld, W. S. (1955). *Biochem. J.* **59**, 294.
- Breuer, H. & Pangels, G. (1960). *Acta endocr., Kbh.*, **33**, 532.
- Brown, J. B. (1955). *Biochem. J.* **60**, 185.
- Cronkite, E. P., Shellabarger, C. J., Bond, V. P. & Lippincott, S. W. (1960). *Radiat. Res.* **12**, 81.
- Engel, L. L., Slaunwhite, W. R., Carter, P. & Nathanson, I. T. (1950). *J. biol. Chem.* **185**, 255.
- Falck, B. (1959). *Acta physiol. scand.* **47** (Suppl.), 163.
- Fishman, J., Bradlow, H. L. & Gallagher, T. F. (1960). *J. biol. Chem.* **235**, 3104.
- Friedgood, H. B., Garst, J. B. & Haagen-Smit, A. J. (1948). *J. biol. Chem.* **175**, 523.
- Hollander, N. & Hollander, V. P. (1958). *J. biol. Chem.* **233**, 1097.
- Huang, W. Y. & Pearlman, W. H. (1962). *J. biol. Chem.* **237**, 1060.
- Ketz, H. A., Witt, H. & Mitzner, M. (1961). *Biochem. Z.* **334**, 73.
- Oakey, R. E. (1961). *Biochem. J.* **81**, 13P.
- Oakey, R. E., Stitch, S. R. & Eccles, S. S. (1963). *J. Chromat.* (in the Press).
- Pearlman, W. H., Pearlman, M. R. J. & Rakoff, A. E. (1954). *J. biol. Chem.* **209**, 803.
- Savard, K. (1953). *J. biol. Chem.* **202**, 457.
- Smith, O. W. & Ryan, K. J. (1961). *Endocrinology*, **69**, 869.
- Stitch, S. R. (1959). *Biochem. J.* **73**, 287.
- Stitch, S. R. & Oakey, R. E. (1961). *Biochem. J.* **81**, 12P.
- Stitch, S. R., Oakey, R. E. & Eccles, S. S. (1962). *Biochem. J.* **82**, 48P.
- Varangot, J. & Cedard, L. (1960). *C.R. Soc. Biol., Paris*, **153**, 1701.
- Wotiz, H. H., Davis, J. W., Lemon, H. M. & Gut, M. (1956). *J. biol. Chem.* **222**, 487.

Biochem. J. (1963) **88**, 76

The Biosynthesis and Metabolism of Oestrogens in the Rat

2. THE CONVERSION OF [4-¹⁴C]TESTOSTERONE INTO [¹⁴C]OESTRADIOL-17 β BY THE OVARY AFTER WHOLE-BODY IRRADIATION

By S. R. STITCH, R. E. OAKEY AND SHEILA S. ECCLES

Medical Research Council Radiobiological Research Unit, Harwell, Didcot, Berks.

(Received 19 November 1962)

Oestrous changes have been shown to persist in the vaginal smears of the rat for short periods after sterilizing doses (4400 r.) of X-irradiation and were followed, often, by periods of continuous vaginal cornification (Mandl & Zuckerman, 1956). This may indicate the ability of the X-irradiated ovary to produce oestrogen even in the absence of an organized follicular system. Slate & Bradbury (1962) observed no immediate irregularity in the vaginal cycles after X-irradiation, regardless of the dose. The continuation of cyclic vaginal change was never observed, however, when all follicular structures were destroyed after irradiation. It has been reported by Drips & Ford (1932) that even slight irradiation, one-tenth to one-fifteenth of the human erythema dose (less than 150 rads), caused temporary irregularity of the oestrous cycle in this

species. These authors noted, also, that the greatest disturbance of the subsequent oestrous cycles occurred when irradiation was carried out in the oestrous stage. With high doses of irradiation no cyclic activity was observed after the complete atrophy of the functioning structures of the ovary.

In mice, it was noted by Bischoff, Ulmann & Ingraham (1944) that oestrus was completely suppressed after 400 r. of X-irradiation and that the effect was irreversible during 13½ months of observation. These results are in contrast with those of Parkes (1926, 1927*a, b, c*), who observed regular oestrous cycles and behaviour up to 69 days after all follicles had been destroyed by irradiation. Nuzhdin, Shapiro, Petrova & Kitaeva (1955) reported that irradiation of the white mouse with doses as small as 50r. had a marked effect on

the oestrous rhythm but that this effect was not apparent until $1\frac{1}{2}$ months after the irradiation.

There appear to have been no direct biochemical measurements of the effect of irradiation on the secretion of oestrogens or other hormones in the rat. The ability of the ovary of the rat to synthesize oestrogen, however, after whole-body irradiation is significant with respect to radiation-induced neoplasia of the mammae. In the Sprague-Dawley rat, in addition to direct irradiation of the target tissue, a functioning ovary is necessary for maximum neoplastic response of the breast tissue to irradiation (Cronkite, Shellabarger, Bond & Lippincott, 1960; Shellabarger, Aponte, Cronkite & Bond, 1962). In the preceding paper (Stitch, Oakey & Eccles, 1963), proof was given that the rat ovary can convert $[4-^{14}\text{C}]$ testosterone into $[^{14}\text{C}]$ oestradiol- 17β . It was pointed out, however, that proof of this conversion *in vitro* was of unknown significance as a physiological process *in vivo* in this species.

We have studied the effect of whole-body irradiation on the capacity of the rat ovary to convert $[4-^{14}\text{C}]$ testosterone into $[^{14}\text{C}]$ oestrogens and have measured the production of oestradiol- 17β at times up to 6 months after irradiation. We have attempted to compare the capacity of ovarian tissue from irradiated and non-irradiated animals to synthesize oestrogen. The present investigation has been designed to obtain this information at various times after a set dose (400 rads) of whole-body irradiation.

MATERIALS AND METHODS

Materials

All materials used were obtained from the same sources and purified, where relevant, by the same processes as described by Stitch *et al.* (1963).

Methods

Animals. Forty-nine female albino rats of the Harwell strain of Wistar origin were taken from eleven litters of similar age. Vaginal smears were taken for at least 7 days in order to establish the oestrous state of these animals.

Irradiation. Rats from each litter were selected when in oestrus and given whole-body irradiation (400 rads midline dose, bilateral irradiation, 64 rads/min., half-value layer 1.2 mm. of Cu). The remaining litter mates were selected in oestrus and were sham-irradiated. The age of the animals at irradiation or sham-irradiation varied between 46 and 58 days.

Tissue. Pairs of irradiated animals were selected when in oestrus, at times varying from 3 hr. to 180 days after irradiation, and were killed by stunning and decapitation. The ovaries were excised immediately and were incubated in 40 ml. of Krebs-Ringer phosphate buffer, pH 7.2, for 5 hr. at 37° . Pairs of control animals were killed, when in oestrus, at the same time-intervals after sham-irradiation,

and the ovaries excised and incubated in a manner identical with that used for the irradiated animals. The control animals were selected as litter mates of the irradiated animals for the corresponding time-points wherever possible.

Isolation of radioactive material from the incubation mixture. The radioactive material was isolated from the incubation mixture after adding carrier oestrone, oestradiol- 17β and oestriol (1 mg. of each) by the procedure described by Stitch *et al.* (1963). The fraction containing the oestrone and the oestradiol- 17β carriers was separated from the oestriol by the method of Friedgood, Garst & Haagen-Smit (1948).

The oestrone-oestradiol fraction was then chromatographed on Celite (Bauld, 1955) according to the modified procedure described by Stitch *et al.* (1963). Elution was carried out with benzene-hexane (1:1, v/v), benzene and benzene-ethylene dichloride (1:3, v/v). After correction for the recovery of oestradiol- 17β carrier, which was determined by fluorimetry (Oakey, Stitch & Eccles, 1963), the yield of radioactive material corresponding to this carrier was determined. Fluorimetry on purified extracts of incubated rat ovaries showed that the contribution of endogenous oestradiol to these experiments was not detectable.

RESULTS

Chromatography was carried out on extracts from ovarian incubation mixtures from both irradiated and control animals at times 3 hr., 1 day, 6 days, 30 days, 50 days and 180 days after irradiation or sham-irradiation respectively. There appears to be no consistent difference in the qualitative pattern of radioactivity eluted from irradiated and control animals.

The distribution of radioactivity and carrier, after partition chromatography of an extract of oestrogen from an incubation mixture of the ovaries from non-irradiated control rats, has been shown by Stitch *et al.* (1963). The chromatographic pattern obtained with ovaries from rats 3 hr. after irradiation is shown in Fig. 1. At 50 days after irradiation a substantial peak of radioactivity accompanies the oestradiol- 17β carrier. This ability to produce, from testosterone, material corresponding to oestradiol- 17β is continued until at least 180 days after irradiation, the limit of the present observations.

For each time-interval the yield of radioactivity corresponding to oestradiol- 17β was calculated as a percentage conversion of the precursor $[4-^{14}\text{C}]$ -testosterone for incubation mixtures of the ovaries from irradiated and control animals. This percentage conversion is plotted, uncorrected for ovarian weight, in Fig. 2. There appears to be no substantial difference in the ability to produce $[^{14}\text{C}]$ oestradiol- 17β from $[4-^{14}\text{C}]$ testosterone between the ovaries of the irradiated and the control rats. Because of the small conversions, internal standards were not used for correcting possible

Fig. 1. Partition chromatography of the phenolic fraction from an incubation of the ovaries of rats 3 hr. after irradiation, with $[4-^{14}\text{C}]$ testosterone. Experimental details are given in the text. Peak 1 is oestrone, peak 2 is oestradiol- 17β . ●, Radioactivity; ○, carrier oestrogen. The histogram represents radioactivity eluted from the column with methanol.

Fig. 2. Synthesis of radioactive oestradiol as a percentage conversion of the precursor $[4-^{14}\text{C}]$ testosterone. Experimental details are given in the text. The yield is plotted uncorrected for ovarian weight. ●, Ovaries from irradiated rats; ○, ovaries from non-irradiated controls.

suppression of the fluorimetry during assay of carrier (Oakley *et al.* 1963). The results obtained are thus only semi-quantitative.

DISCUSSION

The experiments described above provide strong evidence that the ovarian enzymes concerned in the transformation of androgen into oestrogen remain functional after 400 rads of whole-body irradiation in the rat. Moreover, we

have followed the characteristics of the vaginal epithelium of irradiated and non-irradiated control rats for up to 6 months after whole-body irradiation. The results indicate that normal oestrous cycles continue after irradiation (cf. Slate & Bradbury, 1962) but that the amount of cell debris relative to leucocytes in the dioestrus phase is higher in the irradiated animals than in the controls.

Although the material measured and designated oestradiol- 17β has been subjected to a biased chemical and partition-chromatographic purification it has not been subjected to those further serial purifications that were shown by Stitch *et al.* (1963) to be capable of removing small amounts of impurity. Thus the fraction measured is not of 100% radiochemical purity and the influence of irradiation on the synthesis of the contaminating components is unknown. We are, in these experiments, dealing with a limited portion of the biosynthetic pathway only, namely the conversion of androgen into oestrogen. Because this pathway remains intact after irradiation we cannot infer that enzyme systems involved in the complex biosynthesis of androgen from acetate have similar resistance to irradiation. However, some workers (e.g. Hanel, Hjort, Johansen & Purser, 1957; Gould, 1956; Popják, 1956) have shown increased hepatic cholesterol synthesis from acetate after whole-body irradiation. Moreover, the production of androgenic hormone does not appear to diminish after the exposure of rat testes to 5000 and 10 000 r. of X-rays (Abbot, 1959).

Mandl (1959) showed that high doses (about 4400 r.) are necessary to destroy all follicular structures in the rat ovary. Thus the dose that we have used (400 rads of whole-body irradiation) was insufficient to destroy the follicles of the ovary. Histological examination of the ovaries of our rats after 400 rads of whole-body irradiation (D. Lucas, unpublished work) has shown the presence of intact follicles which were smaller than those of the non-irradiated controls.

The source of the ovarian factor controlling continuous vaginal cornification in the rat after all follicles were destroyed by irradiation (Mandl & Zuckerman, 1956) is obscure. Geist, Gaines & Escher (1941) suggested that, in the white mouse, the lutein-like changes that develop in the residual ovarian parenchyma after ovarian irradiation result in the elaboration of sufficient oestrogenic hormone to establish vaginal oestrus in 100% of the animals. Cycles are, however, variable with periods of continuous oestrus of from 1 to 63 days. However, Brambell, Parkes & Fielding (1927) and Parkes & Brambell (1927) found that the development of accessory sexual structures was inhibited in those mice that developed luteal-like ovaries after irradiation, and Huang & Pearlman (1962)

were unable to demonstrate the biosynthesis of oestrogen by the fully-luteinized rat ovary.

Investigations in the rat (Drips & Ford, 1932; Levine & Witschi, 1933; Mandel, 1935) indicate that cyclic oestrous activity ceases when all follicular structures of the ovary are destroyed by irradiation. Genther (1931) found irregular cycles in the female guinea pig after X-irradiation and, often, a prolonged period of oestrus. Whenever oestrus occurred, however, at least one large follicle was found in the ovary.

Thus oestrous cycles may persist after irradiation of the ovary in both the mouse and the rat, but it appears controversial whether regular cycles may continue in the absence of the maturing follicles. In the absence of follicles after heavy doses of X-irradiation persistent vaginal cornification may occur and this could indicate the continued production of oestrogen by the ovary of the irradiated animal. Such ovaries do not respond to gonadotrophin, so the production of oestrogen by the ovaries might be considered autonomous. Another reasonable possibility would be that the adrenal cortex or some other source is responsible for the elaboration of oestrogen. This suggestion is discounted by the observation that the continuous vaginal cornification ceases as soon as the irradiated ovaries are removed (Mandl & Zuckerman, 1956).

In the human, it has been assumed that X-irradiation of the ovary is a rational palliative measure for mammary cancer insofar as it should destroy the capacity of the ovary to secrete oestrogen. This conclusion is open to question. Nathanson, Rice & Meigs (1940) concluded that X-irradiation of the ovary is as efficient as surgery in producing artificial menopause. This conclusion appears to have been true only for the older patients, however, since of six women close to 30 years of age five had one or more menstrual periods after irradiation with 1200–1600r.

Continued oestrogen biosynthetic activity in human ovarian tissue after 2000r. of irradiation has been reported by Lemon, Wotiz, Sommers, Parsons & Davis (1957), using radioactive acetate as a precursor. Smith & Emerson (1954) found, on the basis of the determination of urinary oestrogens, that X-irradiation of the ovary was ineffective for decreasing the excretion of oestrogen and that the output of oestrogen by cancer patients with X-irradiated ovaries was about four times as high as the average output of normal post-menopausal women. This high output was not found from those cancer patients whose ovaries had been surgically removed. In contrast with these results, the findings of Diczfalussy, Notter, Edsmyr & Westman (1959) suggest that irradiation of the human ovary does diminish the excretion of total

oestrogen in patients with breast cancer. The results of these authors, however, show that ovarian irradiation is ineffective in decreasing the urinary output of oestrone or oestradiol-17 β but decreases markedly the excretion of oestriol. Mackay & Garrey (1962) describe the case of a young woman who, after receiving 1050r. to the ovaries, resumed normal menstrual periods after 6 months, conceived and delivered a normal child.

A more critical biochemical appraisal of the effectiveness of ovarian sterilization by radiation to destroy the capacity for oestrogen production may, perhaps, best be effected through an examination of oestrogen concentrations in the arterial and venous blood of the irradiated ovary rather than by biosynthetic studies such as have been described in the present paper.

SUMMARY

1. The effect of whole-body irradiation (400 rads) on the ability of the rat ovary to produce oestrogen *in vitro* from [4-¹⁴C]testosterone at times up to 180 days after irradiation was investigated.

2. After an initial lag, the biosynthesis of oestradiol-17 β , from 6 days after irradiation onwards, appeared to be as high as with tissue from the non-irradiated controls.

3. The biosynthesis of oestrogen by ovarian tissue after the destruction of follicular elements by higher doses of irradiation is discussed.

The authors are grateful to Mrs Jean Tilbrook for technical assistance with this work and to Mr M. Corp for carrying out the irradiations.

REFERENCES

- Abbot, C. R. (1959). *J. Endocrin.* **19**, 33.
- Bauld, W. S. (1955). *Biochem. J.* **59**, 294.
- Bischoff, F., Ulmann, H. J. & Ingraham, L. P. (1944). *Radiology*, **43**, 55.
- Brambell, F. W. R., Parkes, A. S. & Fielding, O. (1927). *Proc. Roy. Soc. B*, **101**, 95.
- Cronkite, E. P., Shellabarger, C. J., Bond, V. P. & Lippincott, S. W. (1960). *Radiat. Res.* **12**, 81.
- Diczfalussy, E., Notter, G., Edsmyr, F. & Westman, A. (1959). *J. clin. Endocrin.* **19**, 1230.
- Drips, D. G. & Ford, F. A. (1932). *Surg. Gynec. Obstet.* **55**, 596.
- Friedgood, H. B., Garst, J. B. & Haagen-Smit, A. J. (1948). *J. biol. Chem.* **174**, 523.
- Geist, S. H., Gaines, J. A. & Escher, G. C. (1941). *Endocrinology*, **29**, 59.
- Genther, I. T. (1931). *Amer. J. Anat.* **48**, 99.
- Gould (1956). In *Progress in Radiobiology: Proc. 4th int. Conf. Radiobiology*, p. 71. Ed. by Mitchell, J. S., Holmes, B. E. & Smith, C. L. Edinburgh and London: Oliver and Boyd.
- Hanel, H. K., Hjort, G., Johansen, G. & Purser, P. R. (1957). *Acta radiol., Stockh.* **48**, 227.

- Huang, W. Y. & Pearlman, W. H. (1962). *J. biol. Chem.* **237**, 1060.
- Lemon, H. M., Wotiz, H. H., Sommers, S. C., Parsons, L. & Davis, J. W. (1957). In *Proc. 2nd int. Symp. Mammary Cancer, University of Perugia*, p. 111. Ed. by Severi, L. Perugia: Division of Cancer Research, University of Perugia.
- Levine, W. T. & Witschi, E. (1933). *Proc. Soc. exp. Biol., N.Y.*, **30**, 1152.
- Mackay, D. J. & Garrey, M. M. (1962). *Lancet*, i, 720.
- Mandel, J. (1935). *Anat. Rec.* **61**, 295.
- Mandl, A. M. (1959). *Proc. Roy. Soc. B*, **150**, 53.
- Mandl, A. M. & Zuckerman, S. (1956). *J. Endocrin.* **13**, 243.
- Nathanson, I. T., Rice, C. & Meigs, J. V. (1940). *Amer. J. Obstet. Gynec.* **40**, 936.
- Nuzhdin, N. I., Shapiro, N. I., Petrova, O. N. & Kitaeva, O. N. (1955). In *Symp. Radiobiology*, p. 113. Moscow: Academy of Sciences of the U.S.S.R., Institute of Genetics and Institute of Biophysics (p. 95, English translation by the Consultants Bureau).
- Oakey, R. E., Stitch, S. R. & Eccles, S. S. (1963). *J. Chromat.* (in the Press).
- Parkes, A. S. (1926). *Proc. Roy. Soc. B*, **100**, 172.
- Parkes, A. S. (1927a). *Proc. Roy. Soc. B*, **101**, 71.
- Parkes, A. S. (1927b). *Proc. Roy. Soc. B*, **101**, 421.
- Parkes, A. S. (1927c). *Proc. Roy. Soc. B*, **102**, 51.
- Parkes, A. S. & Brambell, F. W. R. (1927). *J. R. micr. Soc.* **47**, 97.
- Popják, G. J. (1956). In *Progress in Radiobiology: Proc. 4th int. Conf. Radiobiology*, p. 71. Ed. by Mitchell, J. S., Holmes, B. E. & Smith, C. L. Edinburgh & London: Oliver and Boyd.
- Shellabarger, C. J., Aponte, G. E., Cronkite, E. P. & Bond, V. P. (1962). *Radiat. Res.* **17**, 492.
- Slate, W. G. & Bradbury, J. T. (1962). *Endocrinology*, **70**, 1.
- Smith, O. W. & Emerson, K., jun. (1954). *Proc. Soc. exp. Biol., N.Y.*, **85**, 264.
- Stitch, S. R., Oakey, R. E. & Eccles, S. S. (1963). *Biochem. J.* **88**, 70.

Biochem. J. (1963) **88**, 80

Changes in the Phospholipids of Mouse Skin after Ultraviolet Irradiation *in vitro*

By A. TICKNER

Croydon Group Laboratories, Mayday Hospital, Thornton Heath, Surrey

(Received 21 February 1963)

If a mouse is exposed to ultraviolet radiation, changes occur in the skin that ultimately lead to malignancy (see Blum, 1959, for a complete discussion). In this phenomenon a primary photochemical change may be postulated to occur from which other observed changes derive. Such a photochemical change would be observed with irradiation of the skin *in vitro*, when complications such as inflammation and repair are eliminated. Further, if such irradiation *in vitro* is carried out at low temperature, accompanying or succeeding thermal reactions are correspondingly decreased.

The literature on photochemical changes in nucleic acids and their derivatives (e.g. Hollaender, 1941; Rapport & Canzanelli, 1950) and the known efficiency of ultraviolet light in producing mutations (Hollaender, 1955) suggest that the nucleic acids are the most likely component of the skin to show photochemical change. Such changes, however, were not observed in our experiments, but there were striking changes in the phospholipids. Significant changes were found to be limited to phospholipids containing ethanolamine and were most marked in the ethanolamine plasmalogen fraction.

Preliminary reports of some of the present work have been given (Tickner, 1959, 1960, 1961).

MATERIALS AND METHODS

Chemicals used throughout were of A.R. or, where available, M.A.R. grade.

Adult mice of strain A of either sex were used. The animal was killed by a blow on the head, the hair was shaved off with a scalpel and the mouse was flayed. The skin was freed of subcutaneous tissue by blunt dissection and divided into quarters. Alternate quarters were pooled and weighed. One of the weighed halves was spread in a quartz Petri dish covered with a quartz cover; the other half was similarly spread out in a glass Petri dish that was then enclosed in a closely fitting metal container. The two dishes were placed side by side in a deep-freeze cabinet (Frigidaire JCR 35) at -20° and irradiated with an un-screened medium-pressure mercury lamp (Hanovia, model 10) for 16 hr. at a distance of 50 cm. Under these conditions the temperature of the specimen did not exceed 0° , as judged by the use of a thermometer in apposition to the Petri dishes and also by the persistence of ice in the dish itself. For some experiments, more skin was required than could be obtained from one mouse. In these experiments individual skins were quartered as before and alternate quarters assigned to experimental or control groups.

Separation into and measurement of the main phosphorus-containing groups (acid-soluble phosphorus, phospholipids, RNA phosphorus, protein phosphorus and DNA phosphorus) was carried out by modifications of methods of Schmidt & Thannhauser (1945) and Logan, Mannell & Rossiter (1952). Skin was dispersed in ice-cold water with