of these children failed to reveal any evidence of unusual sensitivity of their skin to sunlight or drugs, or any occurrence of skin rashes or allergic manifestations. The skin of the affected children showed no defect in keratinization and under ultraviolet light it did not differ from that of normal children. The only clinical finding of significance was that both these histidinaemic children had a speech defect (La Du *et al.* 1962). One of the two affected sibs reported by Ghadimi *et al.* (1962) and the histidinaemic child studied by Auerbach *et al.* (1962) also had defects in speech.

SUMMARY

1. A spectrophotometric method to measure histidine α -deaminase in the stratum corneum of normal human skin is described.

2. Normal human stratum corneum lacks urocanase.

3. The skin of two sibs with histidinaemia does not contain any detectable histidine α -deaminase.

4. This lack of histidine α -deaminase would appear to be the biochemical abnormality responsible for histidinaemia.

5. The distribution of histidine α -deaminase in several tissues of the guinea pig has also been examined.

REFERENCES

- Auerbach, V. H., DiGeorge, A. M., Baldrige, R. C., Tourtellotte, C. D. & Brigham, M. P. (1961). *Clin. Res.* **9**, 334.
 Auerbach, V. H., DiGeorge, A. M., Baldrige, R. C., Tourtellotte, C. D. & Brigham, M. P. (1962). *J. Pediat.* **60**, 487.
 Ghadimi, H., Partington, M. W. & Hunter, A. (1961). *New Engl. J. Med.* **265**, 221.
 Ghadimi, H., Partington, M. W. & Hunter, A. (1962). *Pediatrics, N.Y.*, **29**, 714.
 Hais, I. M. & Zenisek, A. (1959). *Amer. Perfum.* **73**, 26.
 La Du, B. N., Howell, R. R., Jacoby, G. A., Seegmiller, J. E. & Zannoni, V. G. (1962). *Biochem. biophys. Res. Commun.* **7**, 398.
 Lowry, O. H., Rosebrough, N. J., Farr, A. L. & Randall, R. J. (1951). *J. biol. Chem.* **193**, 265.
 Schwarz, E. (1961). *Biochem. Z.* **334**, 415.
 Tabor, H. & Mehler, A. H. (1955). In *Methods in Enzymology*, vol. 2, p. 228. Ed. by Colowick, S. P. & Kaplan, N. O. New York: Academic Press Inc.
 Zenisek, A., Kral, J. A. & Hais, I. M. (1955). *Biochim. biophys. Acta*, **18**, 589.

Biochem. J. (1963) **88**, 162

The Distribution of Peptidases in Subcellular Fractions from the Mucosa of the Small Intestine of the Rat

By G. B. ROBINSON*

Department of Medical Biochemistry and Pharmacology, The Medical School, Birmingham

(Received 26 November 1962)

The studies of Newey & Smyth (1959, 1960) established that dipeptides entered the intestinal mucosa and were hydrolysed to free amino acids within the mucosa. G. B. Robinson (unpublished work) subsequently obtained evidence that intestinal peptidases were located primarily within the epithelial cells. Thus hydrolysis of absorbed dipeptides presumably occurs during transfer of the dipeptides across the epithelial cells. The intracellular distribution of intestinal peptidases has now been investigated to obtain information about the site of peptide hydrolysis within the epithelial cells.

A number of workers have reported difficulties in the isolation of subcellular components from small-intestinal mucosa. Schmidt, Bessman & Tann-

hauser (1957) found that mucus, released from the mucosal cells on homogenization, interfered with the fractionation procedure. Baker (1958) and Busch, Davis & Anderson (1958) also reported that mucus interfered with fractionation. To overcome this problem, different homogenizing media have been used. Schmidt *et al.* (1957) used an iso-osmotic salt solution and Buell & Reiser (1959) used 0.6% sodium chloride. Epstein & Shapiro (1959) and Senior & Isselbacher (1960) used iso-osmotic potassium chloride; the latter authors later reported the use of iso-osmotic mannitol (Senior & Isselbacher (1961). Hübscher, Clark & Webb (1962) have used barium sulphate to remove mucus, and Glover & Green (1957) described a technique for washing the particulate components from the adhering mucus. Allard, de Lamirande & Cantero (1957), Triantaphyllopoulos & Tuba (1959) and Borgström & Dahlquist (1958) have reported

* Present address: Department of Biological Chemistry, University of Illinois College of Medicine, Chicago 12, Ill., U.S.A.

fractionations with 0.25M-sucrose as the homogenizing medium.

In the experiments described below a modification of the method of Glover & Green (1957) has been used for fractionation. This work has been published in brief (Robinson, 1962).

MATERIALS AND METHODS

Animals. Male albino rats, weighing about 250 g., were used. They were allowed only water for 16 hr. before use.

Peptides. These were obtained commercially and examined for impurities by paper chromatography. No further purification was found necessary.

Isolation of subcellular fractions from intestinal mucosa. In initial fractionation experiments it was found that much of the subcellular particulate material adhered to the mucus and was sedimented with it at low speeds. Repeated homogenization of the mucus in large volumes of 0.25M-sucrose, followed by low-speed centrifuging, had the effect of washing the particulate material from the mucus. The soluble material released from the cells on homogenization prevented satisfactory sedimentation of the mucus at low centrifuge speeds. It was found that well-defined mucus sediments could be obtained at low centrifuge speeds if this soluble material was previously removed by high-speed centrifuging. The method of fractionation adopted was based on these findings.

Rats were killed by a blow on the head, the whole small intestine was removed, chilled at 0°, and washed with 0.25M-sucrose containing 0.5% of 0.2M-tris-hydrochloric acid buffer, pH 7.4. The intestine was cut into short lengths, which were then cut longitudinally to expose the mucosa. The mucosa was removed by scraping with a scalpel blade.

The mucosal scrapings were homogenized with a Potter-Elvehjem homogenizer in 20 vol. of 0.25M-sucrose. The homogenate was centrifuged at 3.6×10^4 g-min. to sediment all the particulate material together with the mucus. This

sediment was homogenized in 40 vol. of sucrose and centrifuged at 10^4 g-min. This centrifuging resulted in a cloudy supernatant with a well-defined gelatinous residue. The residue was homogenized twice more in 40 vol. of 0.25M-sucrose and centrifuged at 10^4 g-min. The final gelatinous residue was taken as the mucus fraction.

The supernatants from centrifuging at 10^4 g-min. were combined and centrifuged at 1.5×10^5 g-min. to sediment the mitochondrial fraction and at 3.6×10^6 g-min. to sediment the microsomal fraction. The supernatant from this differential centrifuging was combined with supernatant from the initial 3.6×10^4 g-min. centrifuging to give the final supernatant fraction. The fractionation procedure is shown in Fig. 1.

All centrifuging was in a Spinco model L ultracentrifuge equipped with rotor no. 40.

Determination of peptidase activity. The methods followed were those described by Robinson & Shaw (1960). The activities of enzymes hydrolysing L-leucylglycine and glycylglycine were determined. Optimum conditions were determined for the hydrolysis of the substrates by each subcellular fraction and the distribution of enzymes was measured under these conditions. In all experiments the initial concentration of substrate in the reaction mixture was 25 mM. When metal ions were used for activation, the molarity referred to was the molarity of the metal in the reaction mixture. Both dipeptides were hydrolysed under zero-order conditions. In all cases the rate of reaction was proportional to enzyme concentration. One unit of enzyme activity represents the hydrolysis of 1 μ mole of substrate in 10 min.

Determination of cytochrome-oxidase activity. The method used was that of Cooperstein & Lazarow (1951), as modified by Applemans, Wattiaux & de Duve (1955). The rate of reaction was proportional to enzyme concentration for all fractions.

Determination of alkaline-phosphatase activity. The method adopted was that described by Hepple (1958). The substrate was sodium β -glycerophosphate and 1 μ M-Mg²⁺

Fig. 1. Scheme for the subcellular fractionation of intestinal mucosa.

ions were present in the reaction mixture. The rate of reaction was measured under zero-order conditions and the rate was proportional to enzyme concentration.

Determination of protein. Total nitrogen was determined by the micro-Kjeldahl technique; protein was taken as $N \times 6.25$.

RESULTS

Characterization of subcellular fractions

Since a novel method for the subcellular fractionation of intestinal mucosa was used in these experiments, it was essential to characterize the fractions obtained.

Electron microscopy showed that all the fractions were heterogeneous. The mucus fraction was found to contain many mitochondria, although little microsomal material could be seen. The mitochondrial fraction, besides containing many mitochondria, contained some microsomal material. The microsomal fraction contained a great deal of microsomal material and also some fragmented mitochondria together with unidentified debris.

Cytochrome oxidase was shown to be confined to the mitochondria in rat liver (Schneider & Hogeboom, 1950; Hogeboom, Schneider & Stiebich, 1952). Although this association had not been shown for intestinal mucosa, it seemed likely that the distribution of this enzyme might serve as an index of mitochondria in subcellular fractions of mucosa. The results of distribution experiments are given in Table 1. The enzyme activity was almost equally divided between the mucus fraction (44 %) and the mitochondrial fraction (47 %). The microsomal fraction contained 8.8 % of the activity and no activity was detected in the supernatant fraction. The overall recovery of activity was 69.5 %. Comparison of these results with those obtained by electron microscopy indicated that the distribution of cytochrome oxidase followed the distribution of mitochondria in the subcellular fractions. Since nearly 50 % of cytochrome-oxidase activity was retained in the mucus fraction, the

washing procedure was only partially successful in removing mitochondrial material from the mucus. The efficiency of the procedure also varied from experiment to experiment: in Expt. 2, 75 % of the cytochrome-oxidase activity appeared in the mitochondrial fraction; in Expt. 4 little activity was recovered in this fraction. A more protracted washing procedure may have resulted in better recoveries of mitochondria. However, this would have made the volume of homogenizing medium too cumbersome to handle. The recovery of cytochrome-oxidase activity (69.5 %) was quite low, presumably as a result of the protracted homogenization used in the technique. Further homogenization to improve the recoveries of mitochondria might have resulted in further enzyme destruction.

The distribution of alkaline phosphatase in the intestine was studied by Morton (1954) in calves and by Triantaphyllopoulos & Tuba (1959) in rats. These workers found that the enzyme was associated primarily with the microsomal fraction. More recently Holt & Miller (1962), using hamsters, and B. Clark & J. W. Porteous (personal communication), using rabbits, have shown that alkaline phosphatase appears to be associated with the epithelial-cell brush border, which sediments at low centrifuge speeds in media containing EDTA. Therefore the distribution of this enzyme is more likely to be indication of the fate of the brush border during fractionation rather than an indication of the distribution of microsomal material.

The results of measurements of the distribution of alkaline phosphatase in the subcellular fractions from mucosa are given in Table 2. The microsomal fraction contained 63.5 % of the alkaline-phosphatase activity; 23 % was recovered in the mitochondrial fraction and 15.5 % in the mucus fraction. No activity could be detected in the supernatant fraction. The overall recovery of activity was 80 %. These results indicated that most of the epithelial-cell brush border was disrupted to form small particles that sedimented mainly with the micro-

Table 1. *Distribution of cytochrome oxidase in mucosal cell fractions*

The subcellular fractions were prepared as described in the text. For the assay of cytochrome-oxidase activity, 0.1 ml. of suitably diluted subcellular fraction was added to 3 ml. of a solution containing $17 \mu\text{M}$ -reduced cytochrome *c* in 0.03 M-sodium phosphate buffer, pH 7.4. The decrease in extinction at $550 \text{ m}\mu$ was followed every 2 min. Assays were performed at room temperature (18°).

Expt. no.	Recovered activity (%)				Overall recovery (%)
	Mucus fraction	Mitochondrial fraction	Microsomal fraction	Supernatant fraction	
1	47.5	46.0	6.5	None detectable	64.0
2	19.5	75.0	5.5	None detectable	73.6
3	30.0	55.2	14.8	None detectable	67.0
4	79.0	13.0	8.0	None detectable	73.5
Mean	44.0	47.2	8.8	—	69.5
Protein (%)	14.4	9.2	13.4	63.0	103

Table 2. *Distribution of alkaline phosphatase in mucosal cell fractions*

The subcellular fractions were prepared as described in the text. Alkaline-phosphatase activity was determined by measuring the release of inorganic phosphate from sodium β -glycerophosphate. The reaction mixture contained 1 ml. of 0.1M-sodium β -glycerophosphate, 1 ml. of ethanolamine-hydrochloric acid buffer, pH 9.5, 1 ml. of 5 mM-magnesium acetate, 1.8 ml. of water and 0.2 ml. of suitably diluted subcellular fraction. The reaction mixture was incubated for 20 min. at 37° and the degree of hydrolysis determined.

Expt. no.	Recovered activity (%)				Overall recovery (%)
	Mucus fraction	Mitochondrial fraction	Microsomal fraction	Supernatant fraction	
1	10.1	23.4	66.5	None detectable	77.5
2	8.0	28.0	64.0	None detectable	73.0
3	13.2	27.7	67.5	None detectable	89.0
4	30.5	12.9	56.6	None detectable	79.5
Mean	15.5	23.0	63.5	—	80.0
Protein (%)	14.4	9.2	13.4	63.0	103

somal fraction and partly with the mitochondrial fraction. Carnie & Porteous (1962), on the basis of studies with invertase, have suggested that the distribution of brush-border fragments in subcellular fractions depends on the vigour of homogenization. In the fractionation procedure used here the mucosa was subjected to vigorous homogenization, which caused considerable destruction of the brush border. Electron-microscopic examination of the fractions showed no recognizable brush-border fragments.

Characteristics of peptidases of subcellular fractions

Optimum conditions had been previously established for the hydrolysis of L-leucylglycine and glycylglycine by whole homogenates of rat-intestinal mucosa (Robinson & Shaw, 1960). It was possible, however, that the optimum conditions for hydrolysis by peptidases associated with subcellular fractions differed from those found for whole homogenates. Accordingly, optimum conditions for hydrolysis were determined for each subcellular fraction.

L-Leucylglycine. The optimum pH for the hydrolysis of L-leucylglycine by whole homogenates was pH 8.5 (Robinson & Shaw, 1960). The pH optimum for all the subcellular fractions was found to be pH 8.3. No activation of the enzyme activity of subcellular fractions was obtained in the presence of 1 mM-Mn²⁺ or 1 mM-Mg²⁺ ions. Measurements of the distribution of enzymes hydrolysing leucylglycine were made at pH 8.3 without added metal ions.

Glycylglycine. Robinson & Shaw (1960) noted that whole homogenates of rat-intestinal mucosa hydrolysed glycylglycine at two pH optima, pH 7.4 and pH 8.2. The activity at pH 7.4 was enhanced by Co²⁺ ions but was slightly reduced by Mn²⁺ ions; that at pH 8.2 was increased by both

Table 3. *Loss of activity after aging of enzymes from rat-ileal mucosa that hydrolyse glycylglycine at pH 7.4 and 8.4*

Enzyme extracts from rat-ileal mucosa were assayed for glycylglycine-hydrolysing activity before and after periods of aging at 37° and 0°. Assays were conducted at pH 7.4 and pH 8.2 in the presence of 1 mM-Co²⁺ ions with the reaction mixture cited in Table 4. All assays were carried out at 37° and the extent of dipeptide hydrolysis was determined at 15 min. intervals during a 1 hr. incubation period. The percentage loss in activity after aging was calculated.

Expt. no.	Temp. of inactivation	Period of inactivation (hr.)	Percentage of activity lost	
			pH 7.4	pH 8.2
1	37°	2	30	41
2	37	2	0	31
3	37	3	0	20
	37	3½	0	37
4	37	3½	5	30
5	0	18	7	35
6	0	24	20	50

Co²⁺ and by Mn²⁺ ions. These results suggested that two enzymes might be present in rat-intestinal extracts.

This possibility was investigated by studying the stabilities of the enzymes at 37° and at 0°. Enzyme extracts were assayed at both pH values in the presence of 1 mM-Co²⁺ ions, before and after aging; the loss of activity at both pH optima was calculated and the results are shown in Table 3. The activity at pH 7.4 was more stable than that at pH 8.2 and it was concluded that there were probably two separate enzymes present in the homogenates.

The pH optima for hydrolysis by the mucus, microsomal and supernatant fractions were pH 7.5 and pH 8.0. The pH optima for the mitochondrial fractions were pH 7.3 and pH 7.8. The enzyme activity of all the fractions was optimally

activated by 1 mM- Co^{2+} ions. The activity of all fractions was increased by 1 mM- Mn^{2+} ions at pH 8.0 but was inhibited at pH 7.5. Measurements of the activity of the fractions were made at the optimum pH values in the presence of 1 mM- Co^{2+} ions.

Subcellular distribution of peptidase activity

The results of the distribution of leucylglycine-hydrolysing activity are given in Table 4. The results for glycylglycine-hydrolysing activity at pH 7.5 are given in Table 5 and at pH 8.2 in

Table 4. *Protein content and L-leucylglycine-hydrolysing activity of cell fractions from intestinal mucosa*

The subcellular fractions were prepared as described in the text. Peptidase activity was measured by determining the increase in ninhydrin colour as the dipeptide underwent hydrolysis. The reaction mixture contained 0.08 ml. of tris-hydrochloric acid buffer at the appropriate pH value, 0.08 ml. of 62.5 mM-dipeptide and 0.04 ml. of suitably diluted subcellular fraction. The assays were conducted at 37°. The original homogenate was assayed at pH 8.5 and the subcellular fractions were assayed at pH 8.3. One unit of activity represents the hydrolysis of 1 μ mole of substrate in 10 min. Results are means of six experiments.

Fraction	Protein (mg./g. of wet tissue)	Protein (%)	Activity (units/g. of wet tissue)	Activity (%)	Sp. activity (units/mg. of protein)
Mucus	19.0 $\pm$ 6.0	16.3 $\pm$ 5.4	11.5* $\pm$ 5.3	1.18 $\pm$ 1.2	0.59
Mitochondrial	6.65 $\pm$ 1.4	5.65 $\pm$ 1.1	18.3 $\pm$ 3.5	2.0 $\pm$ 1.5	2.75
Microsomal	13.7 $\pm$ 4.3	11.3 $\pm$ 2.1	53.5 $\pm$ 13.2	6.65 $\pm$ 1.9	3.9
Supernatant	81.2 $\pm$ 27.0	67.0 $\pm$ 7.3	870.0 $\pm$ 354	90.0 $\pm$ 3.5	11.7

96.2 $\pm$ 13.2 % of the original protein was recovered; 94.0 $\pm$ 13.6 % of the original activity was recovered.

* No activity found in three out of six experiments.

Table 5. *Protein content and glycylglycine-hydrolysing activity (pH 7.5) of cell fractions from intestinal mucosa*

The subcellular fractions were prepared as described in the text, and the reaction mixture was made up as for Table 4. The original homogenate was assayed at pH 7.4; the subcellular fractions were assayed at pH 7.5, with the exception of the mitochondrial fraction, which was assayed at pH 7.3. In all cases the reaction mixture contained 1 mM- Co^{2+} ions. One unit of activity represents the hydrolysis of 1 μ mole of substrate in 10 min. Results are means of four experiments.

Fraction	Protein (mg./g. of wet tissue)	Protein (%)	Activity (units/g. of wet tissue)	Activity (%)	Sp. activity (units/mg. of protein)
Mucus	30.0 $\pm$ 15.6	21.1 $\pm$ 6.2	0	—	—
Mitochondrial	6.75 $\pm$ 1.6	4.8 $\pm$ 1.8	36.0 $\pm$ 14.3	6.35 $\pm$ 1.5	5.2
Microsomal	11.9 $\pm$ 1.4	9.1 $\pm$ 2.4	50.0 $\pm$ 16.7	9.15 $\pm$ 2.5	4.2
Supernatant	87.0 $\pm$ 18.7	64.5 $\pm$ 5.8	492.0 $\pm$ 230	84.2 $\pm$ 4.0	5.6

97.8 $\pm$ 14.5 % of the original protein was recovered; 104 $\pm$ 18.2 % of the original activity was recovered.

Table 6. *Protein content and glycylglycine-hydrolysing activity (pH 8.0) of cell fractions from intestinal mucosa*

The subcellular fractions were prepared as described in the text, and the reaction mixture was made up as for Table 4. The original homogenate was assayed at pH 8.2; the subcellular fractions were assayed at pH 8.0, with the exception of the mitochondrial fraction, which was assayed at pH 7.8. In all cases the reaction mixture contained 1 mM- Co^{2+} ions. One unit of activity represents the hydrolysis of 1 μ mole of substrate in 10 min. Results are means of five experiments.

Fraction	Protein (mg./g. of wet tissue)	Protein (%)	Activity (units/g. of wet tissue)	Activity (%)	Sp. activity (units/mg. of protein)
Mucus	18.1 $\pm$ 8.5	21.0 $\pm$ 5.7	0	—	—
Mitochondrial	6.5 $\pm$ 1.5	5.2 $\pm$ 1.1	36.8 $\pm$ 13.3	5.9 $\pm$ 1.2	5.6
Microsomal	11.2 $\pm$ 1.9	10.6 $\pm$ 3.8	53.8 $\pm$ 14.6	9.4 $\pm$ 2.9	4.8
Supernatant	81.0 $\pm$ 22.7	65.0 $\pm$ 5.0	556 $\pm$ 264	84.5 $\pm$ 3.9	6.9

102.1 $\pm$ 14.6 % of the original protein was recovered; 92.5 $\pm$ 11.0 % of the original activity was recovered.

Table 6. In all cases between 80 and 90 % of the enzyme activity was recovered in the supernatant fraction. The remaining activity was distributed between the microsomal and mitochondrial fractions; in all experiments the microsomal fraction was more active than the mitochondrial fraction. In only three experiments, with leucylglycine as substrate, was any activity detected in the mucus fraction. The overall recoveries of activity were between 90 and 100 %.

It was possible that the peptidase activity associated with the particulate fractions resulted from adsorption from the soluble fraction. This likelihood was examined by suspending both mitochondrial and microsomal fractions in the supernatant for approximately 1 hr.; the particulate material was isolated again and the enzyme activity against leucylglycine was assayed. The fractions were then subjected to three successive washes in 0.25 M-sucrose and the enzyme activity was assayed after each washing. The fractions showed no change in enzyme activity as a result of these treatments. These experiments did not dismiss the possibility that the enzymic activity of the particulate fractions resulted from adsorption, since experiments with cytochrome *c* have shown that adsorption on to particulate cell components occurred immediately on homogenization and that adsorbed cytochrome *c* was not removed by repeated washing (Beinert, 1951). Suspending the particulate fractions in water for several hours did not cause the release of peptidase activity.

DISCUSSION

A number of workers have studied the subcellular distribution of peptidases. Maver & Greco (1951) reported that the enzymes hydrolysing leucinamide in rat liver, spleen, kidney and hepatoma were located primarily in the supernatant fraction. Rademaker & Soons (1957) reported that enzymes hydrolysing glycylglycine and triglycine were found in the supernatant fraction from rat liver. Hanson & Blech (1959) also reported that several peptidases were located in the supernatant from rat liver, whereas the peptidases of rat kidney appeared in the mitochondrial and microsomal fractions as well as the supernatant fraction, the microsomal fraction being most active. Price, Meister, Gilbert & Greenstein (1949) reported that a particulate fraction from hog kidney contained enzymes hydrolysing glycylalanine. Matheson & Hanes (1959) have also found peptidase activity associated with the microsomal fraction from liver and kidney. The subcellular distribution of peptidase activity in the small intestinal mucosa reported here seems similar to the distributions of peptidases found in rat liver (Maver & Greco,

1951; Rademaker & Soons, 1957; Hanson & Blech, 1959).

Since a major part of the peptidase activity of the intestinal mucosa was found in the supernatant fraction, no conclusions can be made about the site at which these enzymes act within the epithelial cells. It has been reported that in the hamster maltase and invertase are localized in the epithelial-cell brush borders, which were isolated in media containing EDTA (Miller & Crane, 1961); presumably, hydrolysis of absorbed disaccharides occurs in this region of the cell. When pig-intestinal mucosa was fractionated in the absence of EDTA, maltase and invertase appeared in the microsomal fraction (Borgström & Dahlquist, 1958), indicating that fractionation in the absence of EDTA caused disruption of the brush border to smaller fragments. Similar results have been obtained with alkaline phosphatase and are cited above. If the peptidases had been associated with the brush border the distribution of these enzymes might have been expected to be similar to the distribution of alkaline phosphatase in the subcellular fractions. It is possible that the peptidases may be very loosely associated with the brush border *in vivo* and are subsequently released into the supernatant fraction on homogenization. The small amount of peptidase activity found in the microsomal and mitochondrial fractions could be associated with the brush border since alkaline-phosphatase activity appeared in both these fractions.

Holt & Miller (1962) have reported that enzymes hydrolysing leucyl- β -naphthylamide are located in the brush border of hamster-intestinal epithelial cells. Although it was initially thought that this substrate was hydrolysed by leucine aminopeptidase (Green, Tsou, Bressler & Seligman, 1955; Nachlas, Crawford & Seligman, 1957), Glenner, Burstone & Meyer (1959) concluded that the enzyme hydrolysing this substrate was not leucine aminopeptidase but that it had aminopeptidase activity. A comparison of the pH optima of enzymes hydrolysing leucyl- β -naphthylamide and of those hydrolysing leucylglycine also suggests that separate enzymes may be involved. The pH optimum for the hydrolysis of leucyl- β -naphthylamide by enzymes from human serum was pH 7.2 (Goldbarg & Rutenberg, 1958), and that for enzymes from rat liver was pH 8.0 (Nachlas, Goldstein & Seligman, 1962); the pH optimum for dipeptide hydrolysis by enzymes from human serum was pH 7.5 in the presence of Co^{2+} ions and pH 8.0 in the presence of Mn^{2+} ions (Fleisher, 1954), and that for enzymes from rat liver was pH 9.4 in the presence of Mn^{2+} ions (Maver & Greco, 1951). Thus there is some doubt about the nature of the enzymes that catalyse the hydrolysis of leucyl- β -naphthylamide.

SUMMARY

1. A method for the subcellular fractionation of intestinal mucosa is described. The subcellular fractions were characterized by electron microscopy and by determining the distributions of cytochrome oxidase and alkaline phosphatase in the subcellular fractions.

2. The subcellular distribution of peptidases hydrolysing L-leucylglycine and glycylglycine has been determined in intestinal mucosa. Most of the peptidase activity was located in the supernatant fraction. The remaining activity was found mainly in the microsomal fraction, with some in the mitochondrial fraction.

I am indebted to Dr B. Shaw and Professor A. C. Frazer for their help and support during this work, and to Mrs B. Matthews for excellent technical assistance. I am grateful to the University of Birmingham for financial support.

REFERENCES

- Allard, C., de Lamirande, G. & Cantero, A. (1957). *Cancer Res.* **17**, 862.
- Applemans, F., Wattiaux, R. & de Duve, C. (1955). *Biochem. J.* **59**, 438.
- Baker, R. V. (1958). *J. Physiol.* **142**, 563.
- Beinert, H. (1951). *J. biol. Chem.* **190**, 287.
- Borgström, B. & Dahlquist, A. (1958). *Acta chem. scand.* **12**, 1997.
- Buell, G. C. & Reiser, R. (1959). *J. biol. Chem.* **234**, 217.
- Busch, H., Davis, J. R. & Anderson, D. C. (1958). *Cancer Res.* **18**, 916.
- Carnie, J. A. & Porteous, J. W. (1962). *Biochem. J.* **85**, 620.
- Cooperstein, S. J. & Lazarow, A. (1951). *J. biol. Chem.* **189**, 665.
- Epstein, B. & Shapiro, B. (1959). *Biochem. J.* **71**, 615.
- Fleisher, G. A. (1954). *J. biol. Chem.* **206**, 637.
- Glenner, C. G., Burstone, M. S. & Meyer, D. B. (1959). *J. nat. Cancer Inst.* **23**, 857.
- Glover, J. & Green, C. (1957). *Biochem. J.* **67**, 308.
- Goldberg, J. A. & Rutenberg, A. M. (1958). *Cancer*, **11**, 283.
- Green, M. N., Tsou, K. C., Bressler, R. & Seligman, A. M. (1955). *Arch. Biochem. Biophys.* **57**, 458.
- Hanson, H. & Blech, W. (1959). *Hoppe-Seyl. Z.* **315**, 191.
- Hepple, L. A. (1958). In *Methods in Enzymology*, vol. 2, p. 530. Ed. by Colowick, S. P. & Kaplan, N. O. New York: Academic Press Inc.
- Hogeboom, G. H., Schneider, W. C. & Stiebach, M. J. (1952). *J. biol. Chem.* **196**, 111.
- Holt, J. H. & Miller, D. (1962). *Biochim. biophys. Acta*, **58**, 239.
- Hübscher, G., Clark, B. & Webb, M. (1962). *Biochem. J.* **84**, 23P.
- Matheson, A. T. & Hanes, C. S. (1959). *Biochim. biophys. Acta*, **33**, 292.
- Maver, M. E. & Greco, A. E. (1951). *J. nat. Cancer Inst.* **12**, 37.
- Miller, D. & Crane, R. K. (1961). *Biochim. biophys. Acta*, **52**, 293.
- Morton, R. K. (1954). *Biochem. J.* **57**, 595.
- Nachlas, M. M., Crawford, D. T. & Seligman, A. M. (1957). *J. Histochem. Cytochem.* **5**, 264.
- Nachlas, M. M., Goldstein, T. P. & Seligman, A. M. (1962). *Arch. Biochem. Biophys.* **97**, 223.
- Newey, H. & Smyth, D. H. (1959). *J. Physiol.* **145**, 48.
- Newey, H. & Smyth, D. H. (1960). *J. Physiol.* **152**, 367.
- Price, V. E., Meister, A., Gilbert, J. B. & Greenstein, J. P. (1949). *J. biol. Chem.* **181**, 535.
- Rademaker, W. & Soons, J. B. J. (1957). *Biochim. biophys. Acta*, **24**, 451.
- Robinson, G. B. (1962). *Biochem. J.* **84**, 35P.
- Robinson, G. B. & Shaw, B. (1960). *Biochem. J.* **77**, 351.
- Schmidt, G., Bessman, M. J. & Tannhauser, S. J. (1957). *Biochim. biophys. Acta*, **23**, 127.
- Schneider, W. C. & Hogeboom, G. H. (1950). *J. biol. Chem.* **183**, 123.
- Senior, J. R. & Isselbacher, K. J. (1960). *Biochim. biophys. Acta*, **44**, 399.
- Senior, J. R. & Isselbacher, K. J. (1961). *Fed. Proc.* **20**, 245.
- Triantaphyllopoulos, E. & Tuba, J. (1959). *Canad. J. Biochem. Physiol.* **37**, 699.

Biochem. J. (1963) **88**, 168

Enzymic Formation of Oestrone 3-Glucuronide by Rabbit-Liver Microsomes

BY E. R. SMITH* AND H. BREUER

Chemische Abteilung, Chirurgische Universitätsklinik, Bonn, Germany

(Received 29 January 1963)

It has been known for many years that phenolic steroids are excreted in a conjugated form in urine. An oestriol conjugate was first isolated from human pregnancy urine by Cohen & Marrian

* Present address: Courtauld Institute of Biochemistry, The Middlesex Hospital Medical School, London, W. 1.

(1936), and it was found that one glucuronic acid residue was attached to each steroid molecule in ring D (Cohen, Marrian & Odell, 1936; Grant & Marrian, 1950). More recently, Carpenter & Kellie (1962) have shown, by comparison with authentic derivatives, that pregnancy urine contains oestriol