a latent form. The activation of these enzymes was compared with that of glutamic dehydrogenase.

3. In graded activation experiments the activation curves for the three dehydrogenases followed each other closely, but they were distinct from the curves for acid phosphatase.

4. A marked irreversible osmotic activation occurred at sucrose concentrations below 0.025 M. At higher sucrose concentrations only a small degree of reversible activation of glutamic dehydrogenase could be observed.

5. Activation led to a liberation of glutamic and malic dehydrogenases into solution, whereas β -hydroxybutyric dehydrogenase remained bound to the particle residue.

6. Activation is interpreted as being due to an increased permeability of the mitochondrial membrane. It is suggested that two types of activation may be distinguished, depending on whether the increase is sufficiently large to allow enzyme molecules to leak out, or only to allow substrate molecules to penetrate. The former kind is irreversible whereas the latter can possibly be reversed.

These investigations have been supported by grants from the Rockefeller Foundation and the Lilly Research Laboratories. One of us (D.S.B.) is indebted to the Belgian Government for a scholarship and to the Administrators of the H.E. Durham Fund, King's College, Cambridge, for a grant.

REFERENCES

- Appelmans, F. & de Duve, C. (1955). *Biochem. J.* **59**, 426.
 Appelmans, F., Wattiaux, R. & de Duve, C. (1955). *Biochem. J.* **59**, 438.
 Baudhuin, P. (1959). *Arch. int. Physiol. Biochim.* **67**, 106.
 Beaufay, H., Bendall, D. S., Baudhuin, P. & de Duve, C. (1959). *Biochem. J.* **73**, 623.
 Beaufay, H., Bendall, D. S., Baudhuin, P., Wattiaux, R. & de Duve, C. (1959). *Biochem. J.* **73**, 628.
 Beaufay, H. & de Duve, C. (1959). *Biochem. J.* **73**, 604.
 Bendall, D. S. (1958). *Abstr. 4th int. Congr. Biochem., Vienna*, p. 57.
 Berthet, J., Berthet, L., Appelmans, F. & de Duve, C. (1951). *Biochem. J.* **50**, 182.
 Berthet, J. & de Duve, C. (1951). *Biochem. J.* **50**, 174.
 Borgström, B., Sudduth, H. C. & Lehninger, A. L. (1955). *J. biol. Chem.* **215**, 571.
 Cooper, C. & Lehninger, A. L. (1956). *J. biol. Chem.* **219**, 489.
 de Duve, C., Pressman, B. C., Gianetto, R., Wattiaux, R. & Appelmans, F. (1955). *Biochem. J.* **60**, 604.
 Dickman, S. R. & Speyer, J. F. (1954). *J. biol. Chem.* **206**, 67.
 Ernster, L. & Navazio, F. (1956). *Acta chem. scand.* **10**, 1038.
 Gianetto, R. & de Duve, C. (1955). *Biochem. J.* **59**, 433.
 Greville, G. D. & Chappell, J. B. (1959). *Biochim. biophys. Acta*, **33**, 267.
 Hogeboom, G. H. & Schneider, W. C. (1953). *J. biol. Chem.* **204**, 233.
 Johnson, D. & Lardy, H. (1958). *Nature, Lond.*, **181**, 701.
 Kielley, W. W. & Bronk, J. R. (1958). *J. biol. Chem.* **230**, 521.
 Kielley, W. W. & Kielley, R. K. (1951). *J. biol. Chem.* **191**, 485.
 Lehninger, A. L. (1951). *J. biol. Chem.* **190**, 345.
 Lehninger, A. L. & Greville, G. D. (1953). *Biochim. biophys. Acta*, **12**, 188.
 McMurray, W. C., Maley, G. F. & Lardy, H. A. (1958). *J. biol. Chem.* **230**, 219.
 Mahler, H. R., Sarkar, N. K., Vernon, L. P. & Alberty, R. A. (1952). *J. biol. Chem.* **199**, 585.
 Simon, E. W. (1958). *Biochem. J.* **69**, 67.
 Tedeschi, H. & Harris, D. L. (1955). *Arch. Biochem. Biophys.* **53**, 52.
 Wattiaux, R. & de Duve, C. (1956). *Biochem. J.* **63**, 606.

Tissue Fractionation Studies

15. INTRACELLULAR DISTRIBUTION AND PROPERTIES OF β -N-ACETYLGLUCOSAMINIDASE AND β -GALACTOSIDASE IN RAT LIVER

By O. Z. SELLINGER*, H. BEAUFAY†, P. JACQUES‡, ANDREE DOYEN AND C. DE DUVE
Department of Physiological Chemistry, University of Louvain, Belgium

(Received 5 August 1959)

Investigations of the last few years have disclosed that mammalian tissues contain other β -glycosidases besides the well-known and widely studied β -glucuronidase. According to a recent survey by

Conchie, Findlay & Levvy (1959), the most active of these is β -N-acetylglucosaminidase, followed by β -galactosidase. The intracellular distribution of these enzymes has not yet been studied in detail, but Pugh, Leaback & Walker (1957) have published some preliminary observations indicating that the β -N-acetylglucosaminidase of rat kidney is partly associated with subcellular particles and can be released from them by treatment in a Waring Blender.

* Post-doctoral fellow of the U.S. National Institutes of Health. Present address: Istituto Superiore di Sanita, Rome, Italy.

† Chargé de recherches du F.N.R.S.

‡ Aspirant du F.N.R.S.

On the other hand, it is known since the investigations of Walker (1952) that liver β -glucuronidase is largely attached to cytoplasmic particles, where it exhibits phenomena of structure-linked latency and activation similar to those observed in this Laboratory for acid phosphatase (Berthet & de Duve, 1951; Berthet, Berthet, Appelmans & de Duve, 1951). Subsequent studies have shown that, except for the existence of a special microsomal component of β -glucuronidase, the two enzymes are closely related and belong, together with several other acid hydrolases, to a distinct group of cytoplasmic particles which have been called lysosomes (for a review, see de Duve, 1959).

In view of these facts, it seemed of interest to investigate the other β -glycosidases of rat liver with the techniques previously worked out for the study of the lysosomal hydrolases. The present paper describes the results obtained in these studies. Some of them have been briefly reported before (Sellinger & Doyen, 1959).

EXPERIMENTAL

Tissue fractionations

The tissue fractionations were performed according to de Duve, Pressman, Gianetto, Wattiaux & Appelmans (1955). The mitochondrial fractions used for the activation experiments were isolated as described by Appelmans & de Duve (1955); they corresponded essentially to a combination of fractions M and L (see Table 1).

Enzyme assays

Reference enzymes. Cytochrome oxidase was assayed according to Appelmans, Wattiaux & de Duve (1955), but at 25°, glucose 6-phosphatase by the technique of de Duve *et al.* (1955). Total acid phosphatase was measured by the method of Wattiaux & de Duve (1956) in the presence of 0.1% of Triton X-100 as releasing agent or, alternatively, in the presence of 0.025% of digitonin, which, as shown in the present paper, also liberates the enzyme completely; free acid phosphatase was assayed as described by Gianetto & de Duve (1955). Nitrogen was determined by a micro-Kjeldahl procedure.

β -N-Acetylglucosaminidase. In the distribution experiments, total β -N-acetylglucosaminidase activity was measured according to a method derived from that of Conchie (1954) for β -glucosidase. The enzyme preparation was incubated for 30 min. at 37° in a total volume of 2 ml. containing 0.05 M-sodium citrate-HCl buffer, pH 5.0, 0.1% of Triton X-100 or 0.025% of digitonin, and 4 mM-*o*-nitrophenyl-N-acetyl- β -D-glucosaminide. The reaction was stopped by the addition of 3 ml. of 2.75% (w/v) trichloroacetic acid and the mixture was filtered. To 2 ml. of the filtrate were added 0.75 ml. of 0.5 N-NaOH and 1.25 ml. of 0.25 M-glycine-Na₂CO₃ buffer, pH 10.0. The *o*-nitrophenol content of this mixture was measured colorimetrically in a Klett-Summerson photoelectric colorimeter, with a violet light filter (no. 42). Tissue and substrate blanks were run concurrently; they were always very low. It was verified that under these conditions the measured activities were proportional both to enzyme concentration and to incubation time.

In the activation experiments, the free activities of β -N-acetylglucosaminidase and of acid phosphatase were measured simultaneously by incubating the enzyme preparation for 10 min. at 37° in a total volume of 2 ml. containing 0.25 M-sucrose, 0.05 M-pyridine-HCl buffer, pH 5.0, 0.05 M-sodium β -glycerophosphate, pH 5.0 and 4 mM-*o*-nitrophenyl-N-acetyl- β -D-glucosaminide. The reaction was stopped by the addition of 4 ml. of 2.75% (w/v) trichloroacetic acid. Inorganic phosphate and *o*-nitrophenol were determined on separate portions of the filtrate. Total activities were measured similarly in the presence of an emulsifying agent (usually 0.025% of digitonin). Control experiments showed that the activities measured by this procedure were identical with those determined separately by the methods described above. The pyridine buffer was chosen because it was found to inhibit neither of the two enzymes, whereas the aminidase was inhibited by acetate and the phosphatase by citrate.

β -Galactosidase. Free and total β -galactosidase activities were assayed similarly by incubating the enzyme preparation at 37° in a total volume of 2 ml. containing 0.25 M-sucrose, 0.05 M-pyridine-HCl buffer, pH 5.0, and 2.5 mM-*o*-nitrophenyl- β -D-galactoside. The amount of *o*-nitrophenol liberated was measured as in the assay of β -N-acetylglucosaminidase. When total activities were measured, the medium contained in addition 0.1% of Triton X-100 and the incubation time was 2 hr. The incubation time in the free activity assays was 10 min.

Materials

o-Nitrophenyl-N-acetyl- β -D-glucosaminide was prepared by the method of Leaback & Walker (1957), the terminal deacetylation being carried out according to Westphal & Schmidt (1952). *o*-Nitrophenyl β -D-galactoside was purchased from the Sigma Chemical Co. (St Louis, Mo., U.S.A.). Digitonin was a Merck product and the other materials used were those mentioned in previous papers of this series.

RESULTS

Intracellular distribution

In Table 1 are shown the results of the fractionation experiments. With respect to the reference enzymes, cytochrome oxidase, acid phosphatase and glucose 6-phosphatase, the results obtained are very similar to those of de Duve *et al.* (1955), except that the nuclear fraction was more heavily contaminated with cytoplasmic material. The distribution of the glycosidases paralleled that of acid phosphatase, without, however, being entirely identical with it.

These results are represented graphically in Fig. 1. The acid phosphatase data obtained in the experiments on each glycosidase are shown separately in this graph, in order to illustrate better the correlation between the hydrolases. Slight differences in the manner in which the decantations were performed are responsible for the difference in the specific activities of the hydrolases in the two groups of experiments.

Table 1. Intracellular distribution of enzymes

Absolute values are given in mg./g. for nitrogen; in units*/g. (fresh wt.) of liver for enzymes. Statistics refer to means $\pm$ s.d. These are given, rather than s.e.m., since they are a measure of individual variability, the most interesting information in the present case. E, Cytoplasmic extract; N, nuclear fraction; M, heavy mitochondrial fraction; L, light mitochondrial fraction; P, microsomal fraction; S, final supernatant.

Enzyme	No. of expt.	Absolute values E + N	Percentage values					Recovery
			E + N	N	M	L	P	S
Nitrogen	5	31.5 $\pm$ 5.6	100	20.8 $\pm$ 5.0	17.4 $\pm$ 7.2	7.4 $\pm$ 4.0	19.8 $\pm$ 2.9	37.1 $\pm$ 9.2
Cytochrome oxidase	6	27.9 $\pm$ 7.8	100	13.5 $\pm$ 5.7	55.2 $\pm$ 12.4	20.5 $\pm$ 13.4	3.4 $\pm$ 0.9	92.6 $\pm$ 8.4
Acid phosphatase	6	4.88 $\pm$ 0.8	100	9.9 $\pm$ 4.1	20.8 $\pm$ 3.7	37.1 $\pm$ 5.1	24.4 $\pm$ 3.0	11.9 $\pm$ 3.7
β -N-Acetylglucosaminidase	4	6.90 $\pm$ 1.1	100	12.4 $\pm$ 4.6	33.4 $\pm$ 16.4	39.4 $\pm$ 6.9	12.4 $\pm$ 1.5	5.8 $\pm$ 0.9
β -Galactosidase	2	0.34 $\pm$ 0.08	100	7.6 $\pm$ 5.6	29.4 $\pm$ 0.2	30.8 $\pm$ 1.4	10.9 $\pm$ 1.1	10.8 $\pm$ 3.0
Glucose 6-phosphatase	6	14.5 $\pm$ 2.6	100	15.2 $\pm$ 6.5	3.9 $\pm$ 1.5	6.0 $\pm$ 1.6	68.2 $\pm$ 7.6	3.7 $\pm$ 3.1

* Units are those defined by de Duve, Pressman, Gianetto, Wattiaux & Appelmans (1955).

Fig. 1. Distribution patterns of enzymes. Fractions are plotted in the order of their isolation, i.e. (from left to right) N, M, L, P and S (see Table 1). Each fraction is represented separately in the ordinate scale by its own relative specific activity. In the abscissa scale, each fraction is represented (cumulatively from left to right) by its percentage of nitrogen (Table 1).

Activation and release

As preliminary experiments had shown that the two glycosidases are largely masked in fresh particulate fractions incubated under the conditions of the free activity measurements, a comparative investigation of the release of these enzymes and of acid phosphatase was undertaken in the line of previous studies of a similar kind (Gianetto & de Duve, 1955; de Duve *et al.* 1955; Wattiaux & de Duve, 1956).

Comparison of β -N-acetylglucosaminidase with acid phosphatase

In Fig. 2 are shown the results of an experiment in which the free enzymic activities were measured in the presence of increasing concentrations of digitonin. The release of both enzymes occurred in an essentially identical manner, and was complete in the presence of 0.01 % of digitonin. This experiment justifies the use of digitonin as releasing agent in the determination of total activities and shows that it is several times more effective in this respect than Triton X-100 (Wattiaux & de Duve, 1956). Digitonin also has the advantage that it does not interfere with the determination of inorganic phosphate as does Triton X-100. Unfortunately, it has a very low solubility in water and occasionally fails to effect complete activation owing to precipitation.

As shown in Fig. 3, repeated freezing and thawing also caused a parallel liberation of the two

enzymes. In this experiment, as in those of Bendall & de Duve (1960), freezing and thawing proved much more effective on the release of acid phosphatase than had been previously found (Gianetto & de Duve, 1955). This is presumably due to the fact that freezing was performed at a lower temperature (-80° compared with -15° in the earlier experiments).

Through the courtesy of Measuring and Scientific Instruments Ltd. we had an opportunity to submit a particulate preparation to ultrasonic vibrations. The results of this experiment, shown in Fig. 4, indicate that acid phosphatase and β -N-acetylglucosaminidase were both liberated by this treatment and that the phenomenon followed approximately the same course for both enzymes.

As shown in Figs. 5 and 6, β -N-acetylglucosaminidase was also released, as was acid phosphatase, in particulate preparations exposed to media of low osmotic pressure or pre-incubated at pH 5.0 and 37° .

The results obtained in the experiments described above and in a few others of the same type are summarized in Fig. 7, which illustrates the highly significant correlation existing between the free activities of the two enzymes in preparations subjected to graded damage. Fig. 7 also shows that the free β -N-acetylglucosaminidase activity of fresh untreated preparations tends to be lower than their free acid phosphatase activity.

Fig. 2. Release of β -N-acetylglucosaminidase and of acid phosphatase by increasing concentrations of digitonin. Free activities were measured in the presence of digitonin at the concentration indicated, with an amount of particles corresponding to 0.05 g. of original liver/ml. Results are expressed as percentages of highest observed activity. ●, Acid phosphatase; ○, β -N-acetylglucosaminidase.

Fig. 3. Release of β -N-acetylglucosaminidase and of acid phosphatase by repeated freezing and thawing. Free and total activities were measured on 0.2 ml. samples of a suspension of particles in 0.25 M-sucrose (0.5 g. of original liver/ml.) frozen by dipping for 2 min. into propan-2-ol-solid carbon dioxide and then transferred first to ice-water and subsequently to a water bath at 37° until thawing was just completed. Free activities are expressed as percentages of mean total activity. ●, Acid phosphatase; ○, β -N-acetylglucosaminidase.

Fig. 4. Release of β -N-acetylglucosaminidase and of acid phosphatase by ultrasonic treatment. Free activities were measured on 0.2 ml. samples of a suspension of particles in 0.25 M-sucrose (0.5 g. of original liver/ml.) subjected at 0° and for time indicated to 20 kcyc./sec. in a MSE-Mullard ultrasonic disintegrator (60 w). Results are expressed as percentages of total activity before treatment. ●, Acid phosphatase; ○, β -N-acetylglucosaminidase.

A few β -*N*-acetylglucosaminidase assays were also carried out on homogenates and high-speed supernatants from liver lobes subjected to ischaemic necrosis (de Duve & Beaufay, 1959). In these experiments, which were designed to study the reversibility of the biochemical lesions induced by ischaemia, ligatures were placed on the left liver lobe and later removed after 2, 4 and 6 hr. respectively. The animals were killed 15 hr. later and the damaged and undamaged parts of the liver were homogenized and treated as described by de Duve & Beaufay (1959). The results obtained are listed in Table 2. They show that removal of the ligature after 2 hr. or more did not prevent the inactivation of cytochrome oxidase and glucose 6-phosphatase nor the release of acid phosphatase from proceeding almost to completion, but caused an appreciable loss of the latter enzyme, which did not occur when the ligature was maintained until the time of killing (de Duve & Beaufay, 1959). β -*N*-Acetylglucosaminidase was also released to a fairly considerable extent, but, unlike that of acid phosphatase, its total activity did not decrease significantly.

β -Galactosidase. The release of this glycosidase was studied according to an abridged experimental scheme allowing several treatments to be investigated simultaneously on the same preparation. Conditions were chosen so as to provide significant points on each activation curve. Free and total activities of β -galactosidase and of acid phosphatase

were measured in all cases, but on separate samples, since the disparity in the activities of the two enzymes made it necessary to use a larger concentration of particles for β -galactosidase than for acid phosphatase. The results obtained are shown in

Fig. 6. Release of β -*N*-acetylglucosaminidase and of acid phosphatase by incubation at pH 5.0 and 37°. Free and total activities were measured after pre-incubation of the particles (0.1 g. of original liver/ml.) at 37° and for the time indicated in 0.25 M-sucrose containing 0.05 M-pyridine-HCl buffer. Free activities are expressed as percentages of mean total activities. ●, Acid phosphatase; ○, β -*N*-acetylglucosaminidase.

Fig. 5. Release of β -*N*-acetylglucosaminidase and of acid phosphatase by osmotic disruption. Free and total activities were measured on particles kept for 15 min. at 0° in the concentration of sucrose indicated (0.05 g. of original liver/ml.) and then brought back into 0.25 M-sucrose by a 50% dilution with a sucrose solution of suitable concentration. Results are expressed as percentages of mean total activity. ●, Acid phosphatase; ○, β -*N*-acetylglucosaminidase.

Fig. 7. Correlation between free β -*N*-acetylglucosaminidase and acid phosphatase activities of untreated particulate preparations (x) and of preparations exposed to media of lowered sucrose concentration (●), incubated at pH 5.0 and 37° (○), treated with digitonin (■), frozen and thawed (□) and exposed to ultrasonic vibrations (▲). Correlation coefficient, 0.99.

Table 2. Influence of ischaemia on enzymic activities of liver

The left liver lobe was ligated as described by de Duve & Beaufay (1959) and the ligature was later removed at times indicated. The animals were killed 15 hr. after removal of the ligature. The injured lobe and an undamaged part of the liver were homogenized in 0.25 M-sucrose; part of the homogenates were centrifuged for 30 min. at 100 000 g in the Spinco model L centrifuge with rotor no. 40. Total and free activities were measured on the homogenates, unsedimentable (unsed.) activities on the high-speed supernatants. All activities are related to wet weight of tissue. AGase, β -N-Acetylglucosaminidase.

Expt. no.	Treatment of lobe	Percentage of total activity of unligated part				Percentage of corresponding total activity		
		Cytochrome oxidase	Glucose 6-phosphatase	Total acid phosphatase	Total AGase	Free acid phosphatase	Unsed. acid phosphatase	Unsed. AGase
1	Unligated	100	100	100	100	21	5	1
	Ligated (2 hr.)	11	16	68	95	71	19	30
2	Unligated	100	100	100	100	18	6	2
	Ligated (4 hr.)	6	11	50	114	89	31	29
3	Unligated	100	100	100	100	19	7	2
	Ligated (6 hr.)	3	9	69	99	88	46	27

Table 3. Influence of incubation medium on free β -galactosidase and acid phosphatase activities

Fresh particulate preparation were assayed for enzymic activities as described under Methods (separate substrates), or similarly in a substrate mixture containing 0.05 M- β -glycerophosphate together with 2.5 mM- β -galactoside (combined substrates). Activities are expressed as percentages of the total activities measured with separate substrates.

Substrates	Enzymic activities			
	Acid phosphatase		β -Galactosidase	
	Total	Free	Total	Free
Separate	100	9.5	100	19.2
Combined	89.2	9.6	102.5	20.7

summarized form in Fig. 8. They indicate that the release of β -galactosidase under the influence of the applied treatments paralleled that of acid phosphatase, but that the untreated preparations had more free β -galactosidase than acid phosphatase activity. As shown by the data of Table 3, this difference cannot have been due to the difference in the assay conditions, since it was observed also when both enzymes were assayed in media of identical composition.

DISCUSSION

Both by their intracellular distribution and by their structure-linked latency and mode of release under various conditions, the two glycosidases studied in the present investigation have been found to resemble acid phosphatase closely, thus duplicating a pattern of behaviour previously shown to characterize also β -glucuronidase, cathepsin and the two acid nucleases (Gianetto & de Duve, 1955; de Duve *et al.* 1955; Wattiaux & de Duve, 1956), as well as sulphatases A and B (Viala & Gianetto, 1955; Roy, 1958) and phosphoprotein phosphatase (Paigen & Griffiths, 1959). These results suggest strongly that

Fig. 8. Correlation between free β -galactosidase and acid phosphatase activities of untreated particulate preparations ($\times$) and of preparations exposed to 0.175, 0.125 and 0.075 M-sucrose ($\bullet$), incubated at pH 5.0 and 37° for 10, 20, 40 and 60 min. ($\circ$), exposed to digitonin concentrations of 0.025, 0.05, 0.1 and 0.25 mg./ml. at a particle concentration of 0.25 g. of original liver/ml. ($\blacksquare$), frozen and thawed once, twice and four times ($\square$), and treated in a MSE blender (as described by Bendall & de Duve, 1960) for 25, 60 and 180 sec. ($\triangle$). Free enzymic activities were measured on separate samples of the treated suspensions and are expressed as percentages of the corresponding total activities. Correlation coefficient, 0.98.

the newly studied enzymes belong to the same group of particles as acid phosphatase and the other acid hydrolases. They also reinforce the concept, embodied in the name lysosome, that these particles are mainly concerned with acid hydrolysis.

As with the other enzymes of the lysosomal group, the distributions found for the glycosidases

are not identical with that of acid phosphatase, nor, for that matter, with that of any of the other acid hydrolases. This heterogeneity, which in the present case appears to be linked with a difference in the degree of latency of the enzymes, has been commented upon before. It seems to be a general property of lysosomes, which cannot be interpreted unequivocally at the present time (see, for instance, the discussion in de Duve, 1959). It should be noted that the glycosidases do not appear to show a second, microsomal component, as was found for β -glucuronidase, the enzyme most closely related to them functionally (de Duve *et al.* 1955).

The results described here confirm the intimate relationship between β -*N*-acetylglucosaminidase, β -galactosidase and β -glucuronidase established by Conchie *et al.* (1959). These authors, who have made observations very similar to those described in the present paper (G. A. Levvy, personal communication), have made the interesting discovery that tissues which contain the β -glycosidases also contain an α -mannosidase. According to a few preliminary experiments (performed with a sample of substrate kindly provided by Dr G. A. Levvy), α -mannosidase, another soluble acid hydrolase, appears to belong also to the lysosomes.

Conchie *et al.* (1959) have expressed the opinion that these various glycosidases may exert a concerted type of action in the catabolism of mucopolysaccharides and glycoproteins. This hypothesis is in line with the general lytic function attributed to the lysosomes. As discussed by de Duve (1959), the segregation of the hydrolases within impermeable particles probably represents a fundamental control mechanism of their lytic activity.

SUMMARY

1. The intracellular distribution of β -*N*-acetylglucosaminidase and of β -galactosidase has been investigated in rat liver according to a fractionation scheme in which the mitochondrial fraction is divided into two subfractions. It was found to differ distinctly from those of the mitochondrial cytochrome oxidase and of the microsomal glucose 6-phosphatase and to parallel closely that of the lysosomal acid phosphatase.

2. Like acid phosphatase, both enzymes occurred largely in latent form in fresh particulate preparations. Treatments which caused a release of acid phosphatase liberated the β -glycosidases in approximately the same proportion.

3. In both types of experiments, small but significant differences were observed in the behaviour of the three hydrolases.

4. It is concluded from these results that the two β -glycosidases are associated with the lysosomes and the differences observed are attributed to the heterogeneity of this class of particles, already revealed by previous investigations.

These investigations have been supported by grants from the Centre National de Recherches Enzymologiques, the Rockefeller Foundation and the Lilly Research Laboratories.

REFERENCES

- Appelmans, F. & de Duve, C. (1955). *Biochem. J.* **59**, 426.
 Appelmans, F., Wattiaux, R. & de Duve, C. (1955). *Biochem. J.* **59**, 438.
 Bendall, D. S. & de Duve, C. (1960). *Biochem. J.* **74**, 444.
 Berthet, J., Berthet, L., Appelmans, F. & de Duve, C. (1951). *Biochem. J.* **50**, 182.
 Berthet, J. & de Duve, C. (1951). *Biochem. J.* **50**, 174.
 Conchie, J. (1954). *Biochem. J.* **58**, 552.
 Conchie, J., Findlay, F. & Levvy, G. A. (1959). *Biochem. J.* **71**, 318.
 de Duve, C. (1959). In *Subcellular Particles*, p. 128. Ed. by Hayashi, T. New York: Ronald Press Co.
 de Duve, C. & Beaufay, H. (1959). *Biochem. J.* **73**, 610.
 de Duve, C., Pressman, B. C., Gianetto, R., Wattiaux, R. & Appelmans, F. (1955). *Biochem. J.* **60**, 604.
 Gianetto, R. & de Duve, C. (1955). *Biochem. J.* **59**, 433.
 Leaback, D. H. & Walker, P. G. (1957). *J. chem. Soc.* p. 4754.
 Paigen, K. & Griffiths, S. K. (1959). *J. biol. Chem.* **234**, 299.
 Pugh, D., Leaback, D. H. & Walker, P. G. (1957). *Biochem. J.* **65**, 464.
 Roy, A. B. (1958). *Biochem. J.* **68**, 519.
 Sellinger, O. Z. & Doyen, A. (1959). *Arch. int. Physiol. Biochem.* **67**, 519.
 Viala, R. & Gianetto, R. (1955). *Canad. J. Biochem. Physiol.* **33**, 839.
 Walker, P. G. (1952). *Biochem. J.* **51**, 223.
 Wattiaux, R. & de Duve, C. (1956). *Biochem. J.* **63**, 606.
 Westphal, O. & Schmidt, H. (1952). *Liebigs Ann.* **575**, 84.